

11/21/00
JC890 U.S. PTO

11.22.00 Box-88Q

DOCKET NO.: 22221/1030 (RU-339)
EXPRESS MAIL NO.: EL709322449US

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

UTILITY PATENT APPLICATION TRANSMITTAL FORM
(only for new nonprovisional applications under 37 CFR 1.53(b))

ASSISTANT COMMISSIONER FOR PATENTS
Washington, D.C. 20231
BOX: PATENT APPLICATION

JC815 U.S. PTO
09/716964
11/21/00

SIR:

Transmitted herewith for filing is the patent application (including Specification, Claims, and Abstract, 130 pages) of:

Inventors : **Michael E. O'Donnell, Alexander Yuzhakov, Olga Yurieva, David Jeruzalmi, Irina Bruck, and John Kuriyan**

For : **ENZYMES DERIVED FROM THERMOPHILIC ORGANISMS THAT FUNCTION AS A CHROMOSOMAL REPLICASE, PREPARATION AND USE THEREOF**

***If a CONTINUING APPLICATION, please mark where appropriate and supply the requisite information below and in a preliminary amendment:*

☐ continuation ☐ divisional ☒ Continuation-In-Part (CIP)
of prior application Serial No. _____

Prior application information: Examiner :
Art Unit :

Enclosed are:

☒ 83 sheets of Formal drawings.

☐ **Signed** Combined Declaration and Power of Attorney (_____ pages).

☐ **Copy** of **signed** Combined Declaration and Power of Attorney (_____ pages) from a prior application (1.63(d) (for continuation/divisional).

☐ **Signed** statement deleting inventor(s) named in prior application (_____ pages) (1.63(d)(2) and 1.33(b)).

☐ **Incorporation By Reference:** The entire disclosure of the prior application, from which a **copy** of the oath or declaration is supplied herewith, is considered as being part of the disclosure of the enclosed application and is hereby incorporated by reference therein.

☐ Assignment (_____ pages) of the invention to _____.

☐ Certified copy of a foreign priority document.

☐ Associate power of attorney.

☒ Applicants claim small entity status. (See 37 CFR 1.27.)

- ☐ Preliminary Amendment (_____ pages).
- ☐ Information Disclosure Statement, form PTO-1449 (_____ pages) and _____ references.
- ☒ **UNSIGNED** Combined Declaration and Power of Attorney (3 pages).
- ☒ Statement in Accordance with 37 CFR § 1.821(f) and computer readable 3.5" Diskette.
- ☒ Sequence Listing (165 pages).
- ☒ A self-addressed, prepaid postcard acknowledging receipt.
- ☐ Other:


The Filing fee has been calculated as shown below:

(Col. 1)		(Col. 2)	SMALL ENTITY		OR	LARGE ENTITY	
FOR:	NO. FILED	NO. EXTRA	RATE	FEE		RATE	FEE
BASIC FEE	XXXXXXXX	XXXXXXXX	XXXX	\$355	OR	XXXX	\$710
TOTAL CLAIMS	70 - 20 =	50	x 9 =	\$450	OR	x 18 =	\$
INDEP CLAIMS	2 - 3 =	0	x 40 =	\$0	OR	X80 =	\$
<input type="checkbox"/> MULTIPLE DEPENDENT CLAIM PRESENTED			x135 =	\$	OR	x270 =	\$
*If the Total Claims are less than 20 and Indep. Claims are less than 3, enter "0" in Col. 2			TOTAL	\$805	OR	TOTAL	\$

- ☐ Please charge my Deposit Account No. _____ in the amount of \$ _____. **A duplicate copy of this sheet is enclosed.**
- ☒ A check in the amount of **\$805.00** to cover the filing fee is enclosed.
- ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 14-1138. **A duplicate copy of this sheet is enclosed.**
- ☒ Address all future communications to:

Michael L. Goldman
NIXON PEABODY LLP
Clinton Square, P.O. Box 31051
Rochester, New York 14603

Date: November 21, 2000


Edwin V. Merkel
Registration No. 40,087

NIXON PEABODY LLP
Clinton Square, P.O. Box 31051
Rochester, New York 14603
Telephone: (716) 263-1128
Facsimile: (716) 263-1600

JC815 U.S. PTO
09/716964
11/21/00

EXPRESS MAIL CERTIFICATE

DOCKET NO. : 22221/1030 (RU-339)

APPLICANTS : Michael E. O'Donnell, Alexander Yuzhakov, Olga Yurieva, David Jeruzalmi, Irina Bruck, and John Kuriyan

TITLE : ENZYMES DERIVED FROM THERMOPHILIC ORGANISMS
THAT FUNCTION AS A CHROMOSOMAL REPLICASE,
PREPARATION AND USE THEREOF

Certificate is attached to the **Patent Application Including Specification, Claims, and Abstract (130 pages), Unsigned Combined Declaration and Power of Attorney (3 pages), and Sequence Listing (165 pages)** of the above-named application.

"EXPRESS MAIL" NUMBER: EL709322449US
DATE OF DEPOSIT: November 21, 2000

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231, **Box: Patent Application.**

Jo Ann Whalen

(Typed or printed name of person mailing
paper or fee)

Jo Ann Whalen

(Signature of person mailing paper or fee)

09716964-112100

TITLE: **ENZYMES DERIVED FROM THERMOPHILIC
ORGANISMS THAT FUNCTION AS A
CHROMOSOMAL REPLICASE,
PREPARATION AND USE THEREOF**

INVENTORS: **Michael E. O'Donnell, Alexander Yuzhakov, Olga
Yurieva, David Jeruzalmi, Irina Bruck, and John
Kuriyan**

DOCKET NO.: **22221/1030 (RU-339)**

ENZYMES DERIVED FROM THERMOPHILIC ORGANISMS THAT FUNCTION AS A CHROMOSOMAL REPLICASE

The present application is a continuation-in-part of U.S. Patent
5 Application Serial No. 09/057,416 filed April 8, 1998, which claims the benefit of
U.S. Patent Application Serial No. 08/823,407 filed April 8, 1997, and U.S.
Provisional Patent Application Serial No. 60/143,202 filed April 8, 1997, all of which
are hereby incorporated by reference.

The present invention was made with funding from National Institutes
10 of Health Grant No. GM38839. The United States Government may have certain
rights in this invention.

FIELD OF THE INVENTION

15 The present invention relates to thermostable DNA polymerases and,
more particularly, to such polymerases as can serve as chromosomal replicases and
are derived from thermophilic bacteria. More particularly, the invention extends to
DNA polymerase III-type enzymes from thermophilic bacteria, including *Aquifex*
aeolicus, *Thermus thermophilus*, *Thermotoga maritima*, and *Bacillus*
20 *stearothermophilus*, as well as purified, recombinant or non-recombinant subunits
thereof and their use, and to isolated DNA coding for such polymerases and their
subunits. Such DNA is obtained from the respective genes (e.g., *dnaX*, *holA*, *holB*,
dnaA, *dnaN*, *dnaQ*, *dnaE*, *ssb*, etc.) of various thermophilic eubacteria, including but
not limited to *Thermus thermophilus*, *Aquifex aeolicus*, *Thermotoga maritima*, and
25 *Bacillus stearothermophilus*.

BACKGROUND OF THE INVENTION

Thermostable DNA polymerases have been disclosed previously as set
30 forth in U.S. Patent No. 5,192,674 to Oshima et al., U.S. Patent Nos. 5,322,785 and
5,352,778 to Comb et al., U.S. Patent No. 5,545,552 to Mathur, and others. All of the
noted references recite the use of polymerases as important catalytic tools in the
practice of molecular cloning techniques such as polymerase chain reaction (PCR).
Each of the references states that a drawback of the extant polymerases are their

limited thermostability, and consequent useful life in the participation in PCR. Such limitations also manifest themselves in the inability to obtain extended lengths of nucleotides, and in the instance of *Taq* polymerase, the lack of 3' to 5' exonuclease activity, and the drawback of the inability to excise misinserted nucleotides (Perrino, 5 1990).

More generally, such polymerases, including those disclosed in the referenced patents, are of the Polymerase I variety as they are often 90-95kDa in size and may have 5' to 3' exonuclease activity. They define a single subunit with concomitant limits on their ability to hasten the amplification process and to promote 10 the rapid preparation of longer strands of DNA.

Chromosomal replicases are composed of several subunits in all organisms (Kornberg and Baker, 1992). In keeping with the need to replicate long chromosomes, replicases are rapid and highly processive multiprotein machines. Cellular replicases are classically comprised of three components: a clamp, a clamp 15 loader, and the DNA polymerase (reviewed in Kelman and O'Donnell, 1995; McHenry, 1991). For purposes of the present invention, the foregoing components also serve as a broad definition of a "Pol III-type enzyme".

DNA polymerase III holoenzyme (Pol III holoenzyme) is the multi-subunit replicase of the *E. coli* chromosome. Pol III holoenzyme is 20 distinguished from Pol I type DNA polymerases by its high processivity (>50 kbp) and rapid rate of synthesis (750 nts/s) (reviewed in Kornberg and Baker, 1992; Kelman and O'Donnell, 1995). The high processivity and speed is rooted in a ring shaped subunit, called β , that encircles DNA and slides along it while tethering the Pol III holoenzyme to the template (Stukenberg et al., 1991; Kong et al., 1992). The 25 ring shaped β clamp is assembled around DNA by the multisubunit clamp loader, called γ complex. The γ complex couples the energy of ATP hydrolysis to the assembly of the β clamp onto DNA. This γ complex, which functions as a clamp loader, is an integral component of the Pol III holoenzyme particle. A brief overview of the organization of subunits within the holoenzyme and their function follows.

30 Pol III holoenzyme consists of 10 different subunits, some of which are present in multiple copies for a total of 18 polypeptide chains (Onrust et al., 1995). The organization of these subunits in the holoenzyme particle is illustrated in Fig. 1. As depicted in the diagram, the subunits of the holoenzyme can be grouped

09716964.12100

functionally into three components: 1) the DNA polymerase III core is the catalytic unit and consists of the α (DNA polymerase), ϵ (3'-5' exonuclease), and θ subunits (McHenry and Crow, 1979), 2) the β "sliding clamp" is the ring shaped protein that secures the core polymerase to DNA for processivity (Kong et al., 1992), and 3) the 5
protein γ complex ($\gamma\delta\delta'\chi\psi$) is the "clamp loader" that couples ATP hydrolysis to assembly of β clamps around DNA (O'Donnell, 1987; Maki et al., 1988). A dimer of the τ subunit acts as a "macromolecular organizer" holding together two molecules of core (Studwell-Vaughan and O'Donnell, 1991; Low et al., 1976) and one molecule of γ complex forming the Pol III* subassembly (Onrust et al., 1995). This organizing
10 role of τ to form Pol III* is indicated in the center of Fig. 1. Two β dimers associate with the two cores within Pol III* to form the holoenzyme, which is capable of replicating both strands of duplex DNA simultaneously (Maki et al., 1988).

The DNA polymerase III holoenzyme assembles onto a primed template in two distinct steps. In the first step, the γ complex assembles the β clamp
15 onto the DNA. The γ complex and the core polymerase utilize the same surface of the β ring and they cannot both utilize it at the same time (Naktinis et al., 1996). Hence, in the second step the γ complex moves away from β thus allowing access of the core polymerase to the β clamp for processive DNA synthesis. The γ complex and core remain attached to each other during this switching process by the τ subunit organizer.

20 The γ complex consists of 5 different subunits ($\gamma_{2-4}\delta_1\delta'_1\chi_1\psi_1$). An overview of the mechanism of the clamp loading process follows. The δ subunit is the major touch point to the β clamp and leads to ring opening, but δ is buried within γ complex such that contact with β is prevented (Naktinis et al., 1995). The γ subunit is the ATP interactive protein but is not an ATPase by itself (Tsuchihashi and
25 Kornberg, 1989). The δ' subunit bridges the δ and γ subunits resulting in a $\gamma\delta\delta'$ complex that exhibits DNA dependent ATPase activity and is competent to assemble clamps on DNA (Onrust et al., 1991). Upon binding of ATP to γ , a change in the conformation of the complex exposes δ for interaction with β (Naktinis et al., 1995). The function of the smaller subunits, χ and ψ , is to contact SSB (through χ) thus
30 promoting clamp assembly and high processivity during replication (Kelman and O'Donnell, 1995).

09716964.112100

The three component Pol III-type enzyme in eukaryotes contains a clamp that has the same shape as *E. coli* β , but instead of a homodimer it is a heterotrimer. This heterotrimeric ring, called PCNA (proliferating cell nuclear antigen), has 6 domains like β , but instead of each PCNA monomer being composed of 3 domains and dimerizing to form a 6 domain ring (e.g., like β), the PCNA monomer has 2 domains and it trimerizes to form a 6 domain ring (Krishna et al., 1994; Kuriyan and O'Donnell, 1993). The chain fold of the domains are the same in prokaryotes (β) and eukaryotes (PCNA); thus, the rings have the same overall 6-domain ring shape. The clamp loader of the eukaryotic Pol III-type replicase is called RFC (Replication factor C) and it consists of subunits having homology to the γ and δ' subunits of the *E. coli* γ complex (Cullmann et al., 1995). The eukaryotic DNA polymerase III-type enzyme contains either of two DNA polymerases, DNA polymerase δ and DNA polymerase ϵ (Bambara and Jessee, 1991; Linn, 1991; Sugino, 1995). It is entirely conceivable that yet other types of DNA polymerases can function with either a PCNA or β clamp to form a Pol III-type enzyme (for example, DNA polymerase II of *E. coli* functions with the β subunit placed onto DNA by the γ complex clamp loader) (Hughes et al., 1991; Bonner et al., 1992). The bacteriophage T4 also utilizes a Pol III-type 3-component replicase. The clamp is a homotrimer like PCNA, called gene 45 protein (Young et al., 1992). The gene 45 protein forms the same 6-domain ring structure as β and PCNA (Moarefi et al., 2000). The clamp loader is a complex of two subunits called the gene 44/62 protein complex. The DNA polymerase is the gene 43 protein and it is stimulated by the gene 45 sliding clamp when it is assembled onto DNA by the 44/62 protein clamp loader. The Pol III-type enzyme may be either bound together into one particle (e.g., *E. coli* Pol III holoenzyme), or its three components may function separately (like the eukaryotic Pol III-type replicases).

There is an early report on separation of three DNA polymerases from *T.th.* cells, however each polymerase form was reminiscent of the preexisting types of DNA polymerase isolated from thermophiles in that each polymerase was in the 110,000-120,000 range and lacked 3'-5' exonuclease activity (Ruttimann et al., 1985). These are well below the molecular weight of Pol III-type complexes that contain in addition to the DNA polymerase subunit, other subunits such as γ and τ . Although the three polymerases displayed some differences in activity (column elution behavior,

and optimum divalent cation, template, and temperatures) it seems likely that these three forms were either different repair type polymerases or derivatives of one repair enzyme (e.g., Pol I) that was modified by post translational modification(s) that altered their properties (e.g. phosphorylation, methylation, proteolytic clipping of residues that alter activity, or association with different ligands such as a small protein or contaminating DNA). Despite this previous work, it remained to be demonstrated that thermophiles harbor a Pol III-type enzyme that contain multiple subunits such as γ and/or τ , functioned with a sliding clamp accessory protein, or could extend a primer rapidly and processively over a long stretch (>5kb) of ssDNA (Ruttimann et al., 1985).

Previously, it was not known what polymerase thermophilic bacteria used to replicate their chromosome since only Pol I type enzymes have been reported from thermophiles. By distinction, chromosomal replicases, such as Polymerase III, identified in *E. coli*, if available in a thermostable bacterium, with all its accessory subunits, could provide a great improvement over the Polymerase I type enzymes, in that they are generally much more efficient – about 5 times faster – and much more highly processive. Hence, one may expect faster and longer chain production in PCR, and higher quality of DNA sequencing ladders. Clearly, the ability to practice such synthetic techniques as PCR would be enhanced by these methods disclosed for how to obtain genes and subunits of DNA polymerase III holoenzyme from thermophilic sources.

The present invention is directed to achieving these objectives and overcoming the various deficiencies in the art.

SUMMARY OF THE INVENTION

In accordance with the present invention, DNA Polymerase III-type enzymes as defined herein are disclosed that may be isolated and purified from a thermophilic bacterial source, that display rapid synthesis characteristic of a chromosomal replicase, and that possesses all of the structural and processive advantages sought and recited above. More particularly, the invention extends to thermostable Polymerase III-type enzymes derived from thermophilic bacteria that exhibit the ability to extend a primer over a long stretch (>5kb) of ssDNA at elevated

temperature, the ability to be stimulated by a cognate sliding clamp (e.g., β) of the type that is assembled on DNA by a 'clamp' loader (e.g., γ complex), and have clamp loading subunits that show DNA stimulated ATPase activity at elevated temperature and/or ionic strength. Representative thermophile polymerases include those isolated from the thermophilic eubacteria *Aquifex aeolicus* (*A.ae.* polymerase) and other members of the *Aquifex* genus; *Thermus thermophilus* (*T.th.* polymerase), *Thermus favus* (*Tfl/Tub* polymerase), *Thermus ruber* (*Tru* polymerase), *Thermus brockianus* (DYNAZYME™ polymerase), and other members of the *Thermus* genus; *Bacillus stearothermophilus* (*B.st.* polymerase) and other members of the *Bacillus* genus; *Thermoplasma acidophilum* (*Tac* polymerase) and other members of the *Thermoplasma* genus; and *Thermotoga neapolitana* (*Tne* polymerase; see WO 96/10640 to Chatterjee et al.), *Thermotoga maritima* (*Tma* polymerase; see U.S. Patent No. 5,374,553 to Gelfand et al.), and other species of the *Thermotoga* genus (*Tsp* polymerase). In a preferred embodiment, the thermophilic bacteria comprise species of *Aquifex*, *Thermus*, *Bacillus*, and *Thermotoga*, and particularly *A.ae.*, *T.th.*, *B.st.*, and *Tma*.

A particular Polymerase III-type enzyme in accordance with the invention may include at least one of the following sub-units:

- A. a γ subunit having an amino acid sequence corresponding to SEQ. ID. Nos. 4 or 5 (*T.th.*);
- B. a τ subunit having an amino acid sequence corresponding to SEQ. ID. No. 2 (*T.th.*), SEQ. ID. No. 120 (*A.ae.*), SEQ. ID. No. 142 (*T.ma.*) or SEQ. ID. No. 182 (*B.st.*);
- C. a ϵ subunit having an amino acid sequence corresponding to SEQ. ID. No. 95 (*T.th.*), SEQ. ID. No. 128 (*A.ae.*), or SEQ. ID. No. 140 (*T.ma.*);
- D. a α subunit including an amino acid sequence corresponding to SEQ. ID. No. 87 (*T.th.*), SEQ. ID. No. 118 (*A.ae.*), SEQ. ID. No. 138 (*T.ma.*), or SEQ. ID. Nos. 184 (PolC which has both α and ϵ activity, *B.st.*);
- E. a β subunit having an amino acid sequence corresponding to SEQ. ID. No. 107 (*T.th.*), SEQ. ID. No. 122 (*A.ae.*), SEQ. ID. No. 144 (*T.ma.*), or SEQ. ID. No. 174 (*B.st.*);

F. a δ subunit having an amino acid sequence corresponding to SEQ. ID. No. 158 (*T.th.*), SEQ. ID. No. 124 (*A.ae.*), SEQ. ID. No. 146 (*T.ma.*) or SEQ. ID. No. 178 (*B.st.*);

G. a δ' subunit having an amino acid sequence corresponding to
5 SEQ. ID. No. 156 (*T.th.*), SEQ. ID. No. 126 (*A.ae.*), SEQ. ID. No. 148 (*T.ma.*) or
SEQ. ID. No. 180 (*B.st.*);

variants, including allelic variants, muteins, analogs and fragments of any of subparts (A) through (G), and compatible combinations thereof, capable of functioning in DNA amplification and sequencing.

10 The invention also extends to the genes that correspond to and can code on expression for the subunits set forth above, and accordingly includes the following: *dnaX*, *holA*, *holB*, *dnaQ*, *dnaE*, *dnaN*, and *ssb*, as well as conserved variants and active fragments thereof.

Accordingly, the Polymerase III-type enzyme of the present invention
15 comprises at least one gene encoding a subunit thereof, which gene is selected from the group consisting of *dnaX*, *holA*, *holB*, *dnaQ*, *dnaE* and *dnaN*, and combinations thereof. More particularly, the invention extends to the nucleic acid molecule encoding the γ and τ subunits, and includes the *dnaX* gene which has a nucleotide
20 analogs thereof. Likewise, the nucleotide sequences encoding the α subunit (*dnaE* gene), the ϵ subunit (*dnaQ* gene), the β subunit (*dnaN* gene), the δ subunit (*holA* gene), and the δ' subunit (*holB* gene) each comprise the nucleotide sequences as set forth herein, as well as conserved variants, active fragments and analogs thereof.
Those nucleotide sequences for *T.th.* are as follows: *dnaX* (SEQ. ID. No. 3), *dnaE*
25 (SEQ. ID. No. 86), *dnaQ* (SEQ. ID. No. 94), *dnaN* (SEQ. ID. No. 106), *holA* (SEQ. ID. No. 157), and *holB* (SEQ. ID. No. 155). Those nucleotide sequences for *A.ae.* are as follows: *dnaX* (SEQ. ID. No. 119), *dnaE* (SEQ. ID. No. 117), *dnaQ* (SEQ. ID. No. 127), *dnaN* (SEQ. ID. No. 121), *holA* (SEQ. ID. No. 123), and *holB* (SEQ. ID. No. 125). Those nucleotide sequences for *T.ma.* are as follows: *dnaX* (SEQ. ID. No. 141),
30 *dnaE* (SEQ. ID. No. 137), *dnaQ* (SEQ. ID. No. 139), *dnaN* (SEQ. ID. No. 143), *holA* (SEQ. ID. No. 145), and *holB* (SEQ. ID. No. 147). Those nucleotide sequences for *B.st.* are as follows: *dnaX* (SEQ. ID. No. 181), *polC* (SEQ. ID. Nos. 183), *dnaN* (SEQ. ID. No. 173), *holA* (SEQ. ID. No. 177), and *holB* (SEQ. ID. No. 179).

The invention also provides methods and products for identifying, isolating and cloning DNA molecules which encode such accessory subunits encoded by the recited genes of the DNA polymerase III-type enzyme hereof.

Yet further, the invention extends to Polymerase III-type enzymes
5 prepared by the purification of an extract taken from, e.g., the particular thermophile under examination, treated with appropriate solvents and then subjected to chromatographic separation on, e.g., an anion exchange column, followed by analysis of long chain synthetic ability or Western analysis of the respective peaks against antibody to at least one of the anticipated enzyme subunits to confirm presence of Pol
10 III, and thereafter, peptide sequencing of subunits that co purify and amplification to obtain the putative gene and its encoded enzyme.

The present invention also relates to recombinant γ , τ , ϵ , α (as well as PolC), δ , δ' and β subunits and SSB from thermophiles. In the instance of the γ and τ subunits of *T.th.*, the invention includes the characterization of a frameshifting
15 sequence that is internal to the gene and specifies relative abundance of the γ and τ gene products of *T.th. dnaX*. From this characterization, expression of either one of the subunits can be increased at the expense of the other (i.e. mutant frameshift could make all τ , simple recloning at the end of the frameshift could make exclusively γ and no τ).

In a further aspect of the present invention, DNA probes can be
20 constructed from the DNA sequences coding for, e.g., the *T.th.*, *A.ae.*, *T.ma.*, or *B.st. dnaX*, *dnaQ*, *dnaE*, *dnaA*, *dnaN*, *holA*, *holB*, and *ssb* genes, conserved variants and active fragments thereof, all as defined herein, and may be used to identify and isolate the corresponding genes coding for the subunits of DNA polymerase III holoenzyme
25 from other thermophiles, such as those listed earlier herein. Accordingly, all chromosomal replicases (DNA Polymerase III-type) from thermophilic sources are contemplated and included herein.

The invention also extends to methods for identifying Polymerase III-type enzymes by use of the techniques of long-chain extension and elucidation of
30 subunits with antibodies, as described herein and with reference to the examples.

The invention further extends to the isolated and purified DNA Polymerase III from *T.th.*, *A.ae.*, *T.ma.*, and *B.st.*, the amino acid sequences of the γ , τ , ϵ , α (as well as PolC), δ , δ' , and β subunits and SSB, as set forth herein, and the

09746964.1.2100

nucleotide sequences of the corresponding genes from *T.th.*, *A.ae.*, *T.ma.*, or *B.st.* set forth herein, as well as to active fragments thereof, oligonucleotides and probes prepared or derived therefrom and the transformed cells that may be likewise prepared. Accordingly, the invention comprises the individual subunits enumerated
5 above and hereinafter, corresponding isolated polynucleotides and respective amino acid sequences for each of the γ , τ , ϵ , α (as well as PolC), δ , δ' , and β subunits and SSB, and to conserved variants, fragments, and the like, as well as to methods of their preparation and use in DNA amplification and sequencing. In a particular embodiment, the invention extends to vectors for the expression of the subunit genes
10 of the present invention.

The invention also includes methods for the preparation of the DNA Polymerase III-type enzymes and the corresponding subunit genes of the present invention, and to the use of the enzymes and constructs having active fragments thereof, in the preparation, reconstitution or modification of like enzymes, as well as
15 in amplification and sequencing of DNA by methods such as PCR, and like protocols, and to the DNA molecules amplified and sequenced by such methods. In this regard, a Pol III-type enzyme that is reconstituted in the absence of ϵ , or using a mutated ϵ with less 3'-5' exonuclease activity, may be a superior enzyme in either PCR or DNA sequencing applications, (e.g. Tabor et al., 1995).

20 The invention is directed to methods for amplifying and sequencing a DNA molecule, particularly via the polymerase chain reaction (PCR), using the present DNA polymerase III-type enzymes or complexes. In particular, the invention extends to methods of amplifying and sequencing of DNA using thermostable pol III-type enzyme complexes isolated from thermophilic bacteria such as *Thermotoga* and
25 *Thermus* species, or recombinant thermostable enzymes. The invention also provides amplified DNA molecules made by the methods of the invention, and kits for amplifying or sequencing a DNA molecule by the methods of the invention.

In this connection, the invention extends to methods for amplification of DNA that can achieve long chain extension of primed DNA, as by the application
30 and use of Polymerase III-type enzymes of the present invention. An illustration of such methods is presented in Examples 15 and 16, *infra*.

Likewise, kits for amplification and sequencing of such DNA molecules are included, which kits contain the enzymes of the present invention,

including subunits thereof, together with other necessary or desirable reagents and materials, and directions for use. The details of the practice of the invention as set forth above and later on herein, and with reference to the patents and literature cited herein, are all expressly incorporated herein by reference and made a part hereof.

5 As stated, and in accordance with a principal object of the present invention, Polymerase III-type enzymes and their sub-units are provided that are derived from thermophiles and that are adapted to participate in improved DNA amplification and sequencing techniques, and the consequent ability to prepare larger DNA strands more rapidly and accurately.

10 It is a further object of the present invention to provide DNA molecules that are amplified and sequenced using the Polymerase III-type enzymes hereof.

 It is a still further object of the present invention to provide enzymes and corresponding methods for amplification and sequencing of DNA that can be
15 practiced without the participation of the clamp-loading component of the enzyme.

 It is a still further object of the present invention to provide kits and other assemblies of materials for the practice of the methods of amplification and sequencing as aforesaid, that include and use the DNA polymerase III-type enzymes herein as part thereof.

20 One goal of this invention is to fully reconstitute the rapid and processive replicase from an extreme thermophilic eubacterium from fully recombinant protein subunits. One might think that the extreme heat in which these bacteria grow may have resulted in a completely different solution to the problem of chromosome replication. Prior to filing of the previously-identified priority
25 applications, it is believed that Pol III had not been identified in any thermophile until the present inventors found that *Thermus thermophilus*, which grows at a rather high temperature of 70-80°C, would appear to contain a Pol III. Subsequent to this invention, the genome sequence of *A. aeolicus* was published which shows *dnaE*, *dnaN*, and *dnaX* genes. However, previous work did not fully reconstitute the
30 working replication machinery from fully recombinant subunits. A *holA* gene and *holB* has not been identified previously in *T. thermophilus* or *A. aeolicus*, and studies in the *E. coli* system show that delta and delta prime, encoded by *holA* and *holB*, respectively, are essential to loading the beta clamp onto DNA and, thus, is essential

for rapid and processive holoenzyme function (U.S. Patent Nos. 5,583,026 and 5,668,004 to O'Donnell, which are hereby incorporated by reference).

This invention fully reconstitutes a functional DNA polymerase III holoenzyme from the extreme thermophiles *Thermus thermophilus* and *Aquifex*
5 *aeolicus*. *Aquifex aeolicus* grows at an even higher temperature than *Thermus thermophilus*, up to 85°C. In this invention, the genes of *Thermus thermophilus*, *Aquifex aeolicus*, *Thermotoga maritima*, and *Bacillus stearothermophilus* that are necessary to reconstitute the complete DNA polymerase III machinery, which acts as a rapid and processive polymerase, are identified. Indeed, a delta prime (*holB*) and
10 delta (*holA*) subunits are needed.

The *dnaE*, *dnaN*, *dnaX*, *dnaQ*, *holA*, and *holB* genes are used to express and purify the protein "gears", and the proteins are used to reassemble the replication machine. The *T.th.* Pol III is similar to *E. coli*. The *A.ae.* Pol III is slightly dissimilar from the machinery of previously studied replicases. The *A.ae. dnaX* gene
15 encoded only one protein, tau, and in this fashion is similar to the *dnaX* of the gram positive organism, *Staphylococcus aureus*. In contrast, the *dnaX* of the gram negative cell, *E. coli*, produces two proteins. The *Aquifex aeolicus* polymerase subunit, alpha (encoded by *dnaE*) does not contain the 3'-5' proofreading exonuclease. In this regard, *A. aeolicus* is similar to *E. coli*, but dissimilar to the replicase of the gram
20 positive organisms. In Gram positive organisms, the PolC polymerase subunit of the replicase contains the exonuclease activity in the same polypeptide chain as the polymerase (Low et al., 1976; Barnes et al., 1994; Pacitti et al., 1995). Further, the polymerase III of thermophilic bacteria retains activity at high temperature.

Thermostable rapid and processive three component DNA polymerases
25 can be applied to several important uses. DNA polymerases currently in use for DNA sequencing and DNA amplification use enzymes that are much slower and thus could be improved upon. This is especially true of amplification as the three component polymerase is capable of speed and high processivity making possible amplification of very long (tens of Kb to Mb) lengths of DNA in a time-efficient manner. These
30 three component polymerases also function in conjunction with a replicative helicase (DnaB), and thus are capable of amplification at a single temperature, using the helicase to melt the DNA duplex. This property could be useful in some methods of amplification, and in polymerase chain reaction (PCR) methodology. For example, the $\alpha\tau\delta\delta'/\beta$ form of the *E. coli* DNA polymerase III holoenzyme has been shown to

function in both DNA sequencing and PCR (U.S. Patent Nos. 5,583,026 and 5,668,004 to O'Donnell).

Other objects and advantages will become apparent from a review of the ensuing description which proceeds with reference to the following illustrative
5 drawings.

DESCRIPTION OF THE DRAWINGS

FIGURE 1 is a schematic depiction of the structure and components of
10 enzymes of the general family to which the enzymes of the present invention belong.

FIGURE 2 is an alignment of the N-terminal regions of *E. coli* (SEQ. ID. No. 19) and *B. subtilis* (SEQ. ID. No. 20) *dnaX* gene product. Asterisks indicate identities. The ATP binding consensus sequence is indicated. The two regions used for PCR primer design are shown in bold.

15 FIGURE 3 is an image showing the Southern analysis of *T. thermophilus* genomic DNA. Genomic DNA was analyzed for presence of the *dnaZ* gene using the PCR radiolabeled probe. Enzymes used for digestion are shown above each lane. The numbering to the right corresponds to the length of DNA fragments (kb).

20 FIGURES 4A and 4B depict the full sequence of the *dnaX* gene of *T. thermophilus*. DNA sequence (upper case, and corresponding to SEQ ID No. 1) and predicted amino acid sequence (lower case, and corresponding to SEQ ID No. 2) yields a 529 amino acid protein (τ) of 58.0 kDa. A putative frameshifting sequence containing several A residues 1478-1486 (underlined) may produce a smaller protein
25 (γ) of 49.8 kDa. The potential Shine-Dalgarno (S.D.) signal is bold and underlined. The start codon is in bold, and the stop codon for τ is marked by an asterisk. The potential stop codon for γ is shown in bold after the frameshift site, and two potential Shine-Dalgarno sequences upstream of the frameshift site are indicated. Sequences of the primers used for PCR are shown in italics above the nucleotide sequence of *dnaX*.
30 The ATP binding site is indicated, and the asterisks above the four Cys residues near the ATP site indicate the putative Zn^{2+} finger. The proline rich area is indicated above the sequence. Numbering of the nucleotide sequence is presented to the right. Numbering of the amino acid sequence of τ is shown in parenthesis to the right.

FIGURE 4C depicts the isolated DNA coding sequence for the *dnaX* gene (also present in FIGURES 3A and 3B) in accordance with the invention, which corresponds to SEQ. ID. No. 3.

FIGURE 4D depicts the polypeptide sequence of the γ subunit of the Polymerase III of the present invention, which corresponds to SEQ. ID. No. 4.

FIGURE 4E depicts the polypeptide sequence of the γ subunit of the Polymerase III of the present invention defined by a -1 frameshift, which corresponds to SEQ. ID. No. 4.

FIGURE 4F depicts the polypeptide sequence of the γ subunit of the Polymerase III of the present invention defined by a -2 frameshift, which corresponds to SEQ. ID. No. 5.

FIGURES 5A-B are alignments of the γ/τ ATP binding domains for different bacteria. Dots indicate those residues that are identical to the *E. coli dnaX* sequence. The ATP consensus site is underlined, and the conserved cysteine residues that form the zinc finger are indicated with asterisks. *E. coli*, *Escherichia coli* (SEQ. ID. No. 21); *H. inf.*, *Haemophilus influenzae* (SEQ. ID. No. 22); *B. sub.*, *Bacillus subtilis* (SEQ. ID. No. 23); *C. cres.*, *Caulobacter crescentus* (SEQ. ID. No. 24); *M. gen.*, *Mycoplasma genitalium* (SEQ. ID. No. 25); *T.th.*, *Thermus thermophilus* (SEQ. ID. No. 26). Alignments were produced using Clustal.

FIGURE 6 is a diagram indicating a signal for ribosomal frameshifting in *T.th. dnaX*. The diagram shows part of the sequence of the RNA (SEQ. ID. No. 27) around the frameshifting site (SEQ. ID. No. 28), including the suspected slippery sequence A9 (bold italic). The stop codon in the -2 reading frame is indicated. Also indicated are potential step loop structures and the nearest stop codons in the -1 reading frame.

FIGURE 7 is an image showing a Western analysis of γ and τ in *T.th.* cells. Whole cells were lysed in SDS and electrophoresed on a 10 % SDS polyacrylamide gel then transferred to a membrane and probed with polyclonal antibody against *E. coli* γ/τ as described in Experimental Procedures. Positions of molecular weight size markers are shown to the left. Putative *T.th.* γ and τ are indicated to the right.

FIGURES 8A-B are images of *E. coli* colonies expressing *T.th. dnaX* -1 and -2 frameshifts. The region of the *dnaX* gene slippery sequence was cloned into

the *lacZ* gene of pUC19 in three reading frames, then transformed into *E. coli* cells and plated on LB plates containing X-gal. The slippery sequence was also mutated by inserting two G residues into the A9 sequence and then cloned into pUC19 in all three reading frames. Color of colonies observed are indicated by the plus signs. The picture shows the colonies, the type of frameshift required for readthrough (blue color) is indicted next to the sector.

FIGURE 9 shows the construction of the *T.th.* γ/τ expression vector. A genomic fragment containing a partial sequence of *dnaX* was cloned into pALTER-1. This fragment was subcloned into pUC19 (pUC19_*dnaX*). Then the N-terminal section of *dnaX* was amplified such that the fragment was flanked by NdeI (at the initiating codon) and the internal BamHI site. This fragment was inserted to form the entire coding sequence of the *dnaX* gene in pUC19 (pUC19*dnaX*). The *dnaX* gene was then cloned behind the polyhistidine leader in the T7 based expression vector pET16 to give pET16*dnaX*. Details are in "Experimental Procedures".

FIGURES 10A-C illustrate the purification of recombinant *T.th.* γ and τ subunits. *T.th.* γ and τ subunits were expressed in *E. coli* harboring pET16*dnaX*. Molecular size markers are shown to the left of the gels, and the two induced proteins are labeled as g and t to the right of the gel. Panel A) 10% SDS gel of *E. coli* whole cell lysates before and after induction with IPTG. Panel B) 8% SDS gel of the purification two steps after cell lysis. First lane: the lysate was applied to a HiTrap Nickel chromatography column. Second lane: the *T.th.* γ/τ subunits were further purified on a Superose 12 gel filtration column. Third lane, the *E. coli* γ and τ subunits. Panel C) Western analysis of the pure *T.th.* γ and τ subunits (first lane) and *E. coli* γ and τ subunits (second lane).

FIGURES 11A-B show the gel filtration of *T.th.* γ and τ . *T.th.* γ and τ were gel filtered on a Superose 12 column. Column fractions were analyzed for ATPase activity and in a Coomassie Blue stained 10% SDS polyacrylamide gel. Positions of molecular weight markers are shown to the left of the gel. The elution position of size standards analyzed in a parallel Superose 12 column under identical conditions are indicated above the gel. Thyroglobin (670 kDa), bovine gamma globin (150 kDa), chicken ovalbumin (44 kDa), equine myoglobin (17 kDa).

FIGURES 12A-C illustrate the characterization of the *T.th.* γ and τ ATPase activity. The *T.th.* γ/τ and *E. coli* τ subunits are compared in their ATPase

activity characteristics. Due to the greater activity of *E. coli* τ , the values are plotted as percent for ease of comparison. Actual specific activities for 100 % values are given below as pmol ATP hydrolyzed/30 min./pmol *T.th.* γ/τ (or pmol *E. coli* τ).

Panel A) *T.th.* γ and τ ATPase is stimulated by the presence of ssDNA. *T.th.* γ/τ was incubated at 65°C. Specific activity was: 11.5 (+DNA); 2.5 (-DNA); *E. coli* τ was assayed at 37°C. Specific activity values were: 112.5 (+DNA); (7.3-DNA). Panel B) Temperature stability of DNA stimulated ATPase activity. *T.th.* γ/τ , 11.3 (65°C); *E. coli* τ , 97.5 (37°C). Panel C) Stability of *T.th.* γ/τ ATPase to NaCl. *T.th.* γ/τ , 8.1 (100 mM added NaCl and 65°C); *E. coli* τ , 52.7 (0 M added NaCl and 37°C).

FIGURES 13A-13C are graphs that summarize the purification of the DNA polymerase III from *T.th.* extracts. Panel A) shows the activity and total protein in column fractions from the Heparin Agarose column. Peak 1 fractions were chromatographed on ATP agarose. Panel B) depicts the ATP-agarose column step, and Panel C) shows the total protein and DNA polymerase activity eluted from the MonoQ column.

FIGURES 14A-B are SDS polyacrylamide gels of *T.th.* subunits. Fig. 14A is a 12% SDS polyacrylamide gel stained with Coomassie Blue of the MonoQ column. Load stands for the material loaded onto the column (ATP agarose bound fractions). FT stands for protein that flowed through the MonoQ column. Fractions are indicated above the gel. *T.th.* subunits in fractions 17-19 are indicated by the labels placed between fractions 18 and 19. Additional small subunits may be present but difficult to visualize, or may have run off the gel. *E. coli* γ,δ shows a mixture of the α , γ , and δ subunits of DNA polymerase III holoenzyme (they are labeled to the right in the figure). Fig. 14B shows the Western results of an SDS gel of the MonoQ fractions probed with rabbit antiserum raised against the *E. coli* α subunit. Load and FT are as described in Panel A. Fraction numbers are shown above the gel. The band that comigrates with *E. coli* α , and the band in the Coomassie Blue stained gel in Panel A, is marked with an arrow. This band was analyzed for microsequence and the results are shown in Fig. 15.

FIGURES 15A-B show the alignments of the peptides obtained from *T.th.* α subunit, TTH1 (shown in A) and TTH2 (shown in B) with the amino acid sequences of the α subunits of other organisms. The amino acid number of these regions within each respective protein sequence are shown to the right. The

abbreviations of the organisms are as follows. *E.coli* - *Escherichia coli*, *V.chol.* - *Vibrio cholerae*, *H.inf.* - *Haemophilus influenzae*, *R.prow.* - *Rickettsia prowazekii*, *H.pyl.* - *Helicobacter pylori*, *S.sp.* - *Synechocystis sp.*, *M.tub.* - *Mycobacterium tuberculosis*, *T.th.* - *Thermus thermophilus*.

5 FIGURES 16A-C show a nucleotide (Panels A-B, SEQ. ID. No. 86) and amino acid (Panel C, SEQ. ID. No. 87) sequence of the *dnaE* gene encoding the α subunit of DNA polymerase III replication enzyme.

FIGURE 17 shows an alignment of the amino acid sequence of ϵ subunits encoded by *dnaQ* of several organisms. The amino acid sequence of the
10 *Thermus thermophilus* ϵ subunit of *dnaQ* is also shown. *T.th.*, *Thermus thermophilus* (SEQ. ID. No. 88); *D.rad.*, *Deinococcus radiodurans* (SEQ. ID. No. 89); *Bac.sub.*, *Bacillus subtilis* (SEQ. ID. No. 90); *H.inf.*, *Haemophilus influenzae* (SEQ. ID. No. 91); *E.c.*, *Escherichia coli* (SEQ. ID. No. 92); *H.pyl.*, *Helicobacter pylori* (SEQ. ID. No. 93). The regions used to obtain the inner part of the *dnaQ* gene are shown in
15 bold. The starts used for expression of the *T.th.* ϵ subunit are marked.

FIGURES 18A-B show the nucleotide (Panel A, SEQ. ID. No. 94) and amino acid (Panel B, SEQ. ID. No. 95) sequence of the *dnaQ* gene encoding the ϵ subunit of DNA polymerase III replication enzyme.

FIGURES 19A-B show an alignment of the DnaA protein of several
20 organisms. The amino acid sequence of the *Thermus thermophilus* DnaA protein is also shown. *P.mar.*, *Pseudomonas marcesans* (SEQ. ID. No. 96); *Syn.sp.*, *Synechocystis sp.* (SEQ. ID. No. 97); *Bac.sub.*, *Bacillus subtilis* (SEQ. ID. No. 98); *M.tub*; *Mycobacterium tuberculosis* (SEQ. ID. No. 99); *T.th.*, *Thermus thermophilus* (SEQ. ID. No. 100); *E.coli.*, *Escherichia coli* (SEQ. ID. No. 101); *T. mar.*,
25 *Thermatoga maritima* (SEQ. ID. No. 102); and *H.pyl.*, *Helicobacter pylori* (SEQ. ID. No. 103).

FIGURES 20A-B show the nucleotide (Panel A, SEQ. ID. No. 104) and amino acid (Panel B, SEQ. ID. No. 105) sequence of the *dnaA* gene of *Thermus thermophilus*.

30 FIGURES 21A-B show the nucleotide (Panel A, SEQ. ID. No. 106) and amino acid (Panel B, SEQ. ID. No. 107) sequence of the *dnaN* gene encoding the β subunit of DNA polymerase III replication enzyme.

FIGURES 22A-B show an alignment of the β subunit of *T.th.* to the β subunits of other organisms. *T.th.*, *Thermus thermophilus* (SEQ. ID. No. 108); *E. coli*, *Escherichia coli* (SEQ. ID. No. 109); *P. mirab.*, *Proteus mirabilis* (SEQ. ID. No. 110); *H. infl.*, *Haemophilus influenzae* (SEQ. ID. No. 111); *P. put.*, *Pseudomonas putida* (SEQ. ID. No. 112); and *B. cap.*, *Buchnera aphidicola* (SEQ. ID. No. 113).

FIGURE 23 is a map of the pET24:dnaN plasmid. The functional regions of the plasmid are indicated by arrows and italic, restriction sites are marked with bars and symbols. The hatched parts in the plasmid correspond to *T.th. dnaN*.

FIGURES 24A-B show the induction of *T.th. β* in *E. coli* cells harboring the *T.th. β* expression vector. Panel A is the cell induction. The first lane shows molecular weight markers (MW). The second lane shows uninduced *E. coli* cells, and the third lane shows induced *E. coli*. The induced *T.th. β* is indicated by the arrow shown to the left. Induced cells were lysed then treated with heat and the soluble portion was chromatographed on MonoQ. Panel B shows the results of MonoQ purification of *T.th. β* .

FIGURE 25A is a schematic depiction of the use of the use of the enzymes of the present invention in accordance with an alternate embodiment hereof. In this scheme the clamp (β or PCNA) slides over the end of linear DNA to enhance the polymerase (Pol III-type such as Pol III, Pol β or Pol δ .) In this fashion the clamp loader activity is not needed.

FIGURE 25B graphically demonstrates the results of the practice of the alternate embodiment of the invention described and set forth in Example 15, *infra*. Lane 1, *E. coli* Pol III without β ; Lane 2, *E. coli* with β ; Lane 3, human Pol δ without PCNA; Lane 4, human Pol δ with PCNA; Lane 5, *T.th.* Pol III without *T.th. β* ; Lane 6, *T.th.* Pol III with *T.th. β* . The respective pmol synthesis in lanes 1-6 are: 6, 35, 2, 24, 0.6 and 1.9.

FIGURES 26A-B show the use of *T.th.* Pol III in extending singly primed M13mp18 to an RFII form. The scheme in Fig. 26A shows the primed template in which a DNA 57mer was annealed to the M13mp18 ssDNA circle. Then *T.th. β* subunit (produced recombinantly) and *T.th.* Pol III were added to the DNA in the presence of radioactive nucleoside triphosphates. In Fig. 26B, the products of the reaction were analyzed in a 0.8% native agarose gel. The position of ssDNA starting

material, the RFII product, and of intermediate species, are shown to the sides of the gel. Lane 1, use of Pol III. Lane 2, use of the non-Pol III DNA polymerase.

FIGURE 27 is an SDS polyacrylamide gel of the proteins of the *A. aeolicus* replication machinery.

5 FIGURE 28 is an SDS polyacrylamide gel analysis of the MonoQ fractions of the method used to reconstitute and purify the *A. aeolicus* $\tau\delta\delta'$ complex.

FIGURE 29 is an SDS polyacrylamide gel analysis of the gel filtration column fractions used in the preparation of the *A. aeolicus* $\alpha\tau\delta\delta'$ complex. The bottom gel analysis shows the profile obtained using the *A. aeolicus* α subunit (polymerase) in the absence of the other subunits.

10

FIGURE 30 is an alkaline agarose gel analysis of reaction products for extension of a single primer around a 7.2 kb M13mp18 circular ssDNA genome that has been coated with *A. aeolicus* SSB. The time course on the left are produced by $\alpha\tau\delta\delta'/\beta$, and the time course on the right is produced by $\alpha\tau\delta\delta'$ in the absence of β .

15 FIGURE 31 is a graph illustrating the optimal temperature for activity of the alpha subunit of *Thermus* replicase using a calf thymus DNA replication assay. Reactions were shifted to the indicated temperature for 5 minutes before detecting the level of DNA synthesis activity.

FIGURE 32 is a graph illustrating the optimal temperature for activity of the alpha subunit of the *Aquifex* replicase using a calf thymus DNA replication assay. Reactions were shifted to the indicated temperature for 5 minutes before detecting the level of DNA synthesis activity.

20

FIGURES 33A-E illustrate the heat stability of *Aquifex* components. Assays of either α (Fig. 33A), β (Fig. 33B), $\tau\delta\delta'$ complex (Fig. 33C), SSB (Fig. 33D) and $\alpha\tau\delta\delta'$ complex (Fig. 33E) were performed after heating samples at the indicated temperatures. Components were heated in buffer containing the following: 0.1% Triton X-100 (filled diamonds); 0.05% Tween-20 and 0.01% NP-40 (filled circles); 4 mM CaCl_2 (filled triangles); 40% Glycerol (inverted filled triangles); 0.01% Triton X-100, 0.05% Tween-20, 0.01% NP-40, 4 mM CaCl_2 (half-filled square); 40% Glycerol, 0.1% Triton X-100 (open diamonds); 40% Glycerol, 0.05% Tween-20, 0.01% NP-40 (open circles); 40% Glycerol, 4 mM CaCl_2 (open triangles); 40% Glycerol, 0.01% Triton X-100, 0.05% Tween-20, 0.01% NP-40, 4 mM CaCl_2 (half-filled diamonds).

25

30

FIGURES 34A-B show the nucleotide sequence (SEQ. ID. No. 117) of the *dnaE* gene of *A. aeolicus*.

FIGURE 35 shows the amino acid sequence (SEQ. ID. No. 118) of the α subunit of *A. aeolicus*.

5 FIGURE 36 shows the nucleotide sequence (SEQ. ID. No. 119) of the *dnaX* gene of *A. aeolicus*.

FIGURE 37 shows the amino acid sequence (SEQ. ID. No. 120) of the tau subunit of *A. aeolicus*.

10 FIGURE 38 shows the nucleotide sequence (SEQ. ID. No. 121) of the *dnaN* gene of *A. aeolicus*.

FIGURE 39 shows the amino acid sequence (SEQ. ID. No. 122) of the β subunit of *A. aeolicus*.

FIGURE 40 shows the partial nucleotide sequence (SEQ. ID. No. 123) of the *holA* gene of *A. aeolicus*.

15 FIGURE 41 shows the partial amino acid sequence (SEQ. ID. No. 124) of the δ subunit of *A. aeolicus*.

FIGURE 42 shows the nucleotide sequence (SEQ. ID. No. 125) of the *holB* gene of *A. aeolicus*.

20 FIGURE 43 shows the amino acid sequence (SEQ. ID. No. 126) of the δ' subunit of *A. aeolicus*.

FIGURE 44 shows the nucleotide sequence (SEQ. ID. No. 127) of the *dnaQ* of *A. aeolicus*.

FIGURE 45 shows the amino acid sequence (SEQ. ID. No. 128) of the ϵ subunit of *A. aeolicus*.

25 FIGURE 46 shows the nucleotide sequence (SEQ. ID. No. 129) of the *ssb* gene of *A. aeolicus*.

FIGURE 47 shows the amino acid sequence (SEQ. ID. No. 130) of the single-strand binding protein of *A. aeolicus*.

30 FIGURE 48 shows the nucleotide sequence (SEQ. ID. No. 131) of the *dnaB* gene of *A. aeolicus*.

FIGURE 49 shows the amino acid sequence (SEQ. ID. No. 132) of the DnaB helicase of *A. aeolicus*.

0916964-112100

FIGURE 50 shows the nucleotide sequence (SEQ. ID. No. 133) of the *dnaG* gene of *A. aeolicus*.

FIGURE 51 shows the amino acid sequence (SEQ. ID. No. 134) of the DnaG primase of *A. aeolicus*.

5 FIGURE 52 shows the nucleotide sequence (SEQ. ID. No. 135) of the *dnaC* gene of *A. aeolicus*.

FIGURE 53 shows the amino acid sequence (SEQ. ID. No. 136) of the DnaC protein of *A. aeolicus*.

10 FIGURE 54A-B shows the nucleotide sequence (SEQ. ID. No. 137) of the *dnaE* gene of *T. maritima*.

FIGURE 55 shows the amino acid sequence (SEQ. ID. No. 138) of the α subunit of *T. maritima*.

FIGURE 56 shows the nucleotide sequence (SEQ. ID. No. 139) of the *dnaQ* gene of *T. maritima*.

15 FIGURE 57 shows the amino acid sequence (SEQ. ID. No. 140) of the ϵ subunit of *T. maritima*.

FIGURE 58 shows the nucleotide sequence (SEQ. ID. No. 141) of the *dnaX* gene of *T. maritima*.

20 FIGURE 59 shows the amino acid sequence (SEQ. ID. No. 142) of the tau subunit of *T. maritima*.

FIGURE 60 shows the nucleotide sequence (SEQ. ID. No. 143) of the *dnaN* gene of *T. maritima*.

FIGURE 61 shows the amino acid sequence (SEQ. ID. No. 144) of the β subunit of *T. maritima*.

25 FIGURE 62 shows the nucleotide sequence (SEQ. ID. No. 145) of the *holA* gene of *T. maritima*.

FIGURE 63 shows the amino acid sequence (SEQ. ID. No. 146) of the δ subunit of *T. maritima*.

30 FIGURE 64 shows the nucleotide sequence (SEQ. ID. No. 147) of the *holB* gene of *T. maritima*.

FIGURE 65 shows the amino acid sequence (SEQ. ID. No. 148) of the δ' subunit of *T. maritima*.

FIGURE 66 shows the nucleotide sequence (SEQ. ID. No. 149) of the *ssb* gene of *T. maritima*.

FIGURE 67 shows the amino acid sequence (SEQ. ID. No. 150) of the single-strand binding protein of *T. maritima*.

5 FIGURE 68 shows the nucleotide sequence (SEQ. ID. No. 151) of the *dnaB* gene of *T. maritima*.

FIGURE 69 shows the amino acid sequence (SEQ. ID. No. 152) of the DnaB helicase of *T. maritima*.

10 FIGURE 70 shows the nucleotide sequence (SEQ. ID. No. 153) of the *dnaG* gene of *T. maritima*.

FIGURE 71 shows the amino acid sequence (SEQ. ID. No. 154) of the DnaG primase of *T. maritima*.

FIGURE 72 shows the nucleotide sequence (SEQ. ID. No. 155) of the *holB* gene of *T. thermophilus*.

15 FIGURE 73 shows the amino acid sequence (SEQ. ID. No. 156) of the δ' subunit of *T. thermophilus*.

FIGURE 74 shows the nucleotide sequence (SEQ. ID. No. 157) of the *holA* gene of *T. thermophilus*.

20 FIGURE 75 shows the amino acid sequence (SEQ. ID. No. 158) of the δ subunit of *T. thermophilus*.

FIGURE 76 shows the nucleotide sequence (SEQ. ID. No. 171) of the *ssb* gene of *T. thermophilus*.

FIGURE 77 shows the amino acid sequence (SEQ. ID. No. 172) of the single-strand binding protein of *T. thermophilus*.

25 FIGURE 78 shows the partial nucleotide sequence (SEQ. ID. No. 173) of the *dnaN* gene of *B. stearothermophilus*.

FIGURE 79 shows the partial amino acid sequence (SEQ. ID. No. 174) of the β subunit of *B. stearothermophilus*.

30 FIGURE 80 shows the nucleotide sequence (SEQ. ID. No. 175) of the *ssb* gene of *B. stearothermophilus*.

FIGURE 81 shows the amino acid sequence (SEQ. ID. No. 176) of the single-strand binding protein of *B. stearothermophilus*.

FIGURE 82 shows the nucleotide sequence (SEQ. ID. No. 177) of the *hola* gene of *B. stearothermophilus*.

FIGURE 83 shows the amino acid sequence (SEQ. ID. No. 178) of the δ subunit of *B. stearothermophilus*.

5 FIGURE 84 shows the nucleotide sequence (SEQ. ID. No. 179) of the *holB* gene of *B. stearothermophilus*.

FIGURE 85 shows the amino acid sequence (SEQ. ID. No. 180) of the δ' subunit of *B. stearothermophilus*.

10 FIGURES 86A-B show the partial nucleotide sequence (SEQ. ID. No. 181) of the *dnaX* gene of *B. stearothermophilus*.

FIGURE 87 shows the partial amino acid sequence (SEQ. ID. No. 182) of the tau subunit of *B. stearothermophilus*.

15 FIGURES 88A-B show the nucleotide sequence (SEQ. ID. No. 183) of the *polC* gene of *B. stearothermophilus*.

FIGURE 89 shows the amino acid sequence (SEQ. ID. No. 184) of the PolC or α -large subunit of *B. stearothermophilus*.

DETAILED DESCRIPTION OF THE INVENTION

20 In accordance with the present invention there may be employed conventional molecular biology, microbiology, and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature. See, e.g., Sambrook et al., "Molecular Cloning: A Laboratory Manual" (1989); "Current Protocols in Molecular Biology" Volumes I-III (Ausubel, R. M., ed.) (1994); "Cell
25 Biology: A Laboratory Handbook" Volumes I-III (Celis, J.E., ed.) (1994); "Current Protocols in Immunology" Volumes I-III (Coligan, J.E., ed.) (1994); "Oligonucleotide Synthesis" (M.J. Gait, ed.) (1984); "Nucleic Acid Hybridization" (B.D. Hames & S.J. Higgins, eds.) (1985); "Transcription And Translation" (B.D. Hames & S.J. Higgins, eds.) (1984); "Animal Cell Culture" (R.I. Freshney, ed.) (1986);
30 "Immobilized Cells And Enzymes" (IRL Press) (1986); B. Perbal, "A Practical Guide To Molecular Cloning" (1984), each of which is hereby incorporated by reference.

Therefore, if appearing herein, the following terms shall have the definitions set out below.

09716964.112100

09716964.112100

The terms "DNA Polymerase III," "Polymerase III-type enzyme(s)",
"Polymerase III enzyme complex(s)", "*T.th.* DNA Polymerase III", "*A.ae.* DNA
Polymerase III", "*T.ma.* DNA Polymerase III", and any variants not specifically listed,
5 are also γ complex, clamp loader, and RFC, as used throughout the present application
and claims refer to proteinaceous material including single or multiple proteins, and
extends to those proteins having the amino acid sequence data described herein and
presented in the Figures and corresponding Sequence Listing entries, and the
corresponding profile of activities set forth herein and in the Claims. Accordingly,
10 proteins displaying substantially equivalent or altered activity are likewise
contemplated. These modifications may be deliberate, for example, such as
modifications obtained through site-directed mutagenesis, or may be accidental, such
as those obtained through mutations in hosts that are producers of the complex or its
named subunits. Also, the terms "DNA Polymerase III," "*T.th.* DNA Polymerase III,"
15 and " γ and τ subunits", " β subunit", " α subunit", " ϵ subunit", " δ subunit", " δ'
subunit", "SSB protein", "sliding clamp" and "clamp loader" are intended to include
within their scope proteins specifically recited herein as well as all substantially
homologous analogs and allelic variations. As used herein γ complex refers to a
particular type of clamp loader that includes a γ subunit.

20 Also as used herein, the term "thermolabile enzyme" refers to a DNA
polymerase which is not resistant to inactivation by heat. For example, T5 DNA
polymerase, the activity of which is totally inactivated by exposing the enzyme to a
temperature of 90°C for 30 seconds, is considered to be a thermolabile DNA
polymerase. As used herein, a thermolabile DNA polymerase is less resistant to heat
25 inactivation than in a thermostable DNA polymerase. A thermolabile DNA
polymerase typically will also have a lower optimum temperature than a thermostable
DNA polymerase. Thermolabile DNA polymerases are typically isolated from
mesophilic organisms, for example mesophilic bacteria or eukaryotes, including
certain animals.

30 As used herein, the term "thermostable enzyme" refers to an enzyme
which is stable to heat and is heat resistant and catalyzes (facilitates) combination of
the nucleotides in the proper manner to form the primer extension products that are
complementary to each nucleic acid strand. Generally, the synthesis will be initiated

at the 3' end of each primer and will proceed in the 5' direction along the template strand, until synthesis terminates, producing molecules of different lengths.

The thermostable enzyme herein must satisfy a single criterion to be effective for the amplification reaction, i.e., the enzyme must not become irreversibly denatured (inactivated) when subjected to the elevated temperatures for the time necessary to effect denaturation of double-stranded nucleic acids. Irreversible denaturation for purposes herein refers to permanent and complete loss of enzymatic activity. The heating conditions necessary for denaturation will depend, e.g., on the buffer salt concentration and the length and nucleotide composition of the nucleic acids being denatured, but typically range from about 90°C to about 96°C for a time depending mainly on the temperature and the nucleic acid length, typically about 0.5 to four minutes. Higher temperatures may be tolerated as the buffer salt concentration and/or GC composition of the nucleic acid is increased. Preferably, the enzyme will not become irreversibly denatured at about 90°-100°C.

The thermostable enzymes herein preferably have an optimum temperature at which they function that is higher than about 40°C, which is the temperature below which hybridization of primer to template is promoted, although, depending on (1) magnesium and salt concentrations and (2) composition and length of primer, hybridization can occur at higher temperature (e.g., 45°-70°C). The higher the temperature optimum for the enzyme, the greater the specificity and/or selectivity of the primer-directed extension process. However, enzymes that are active below 40°C, e.g., at 37°C, are also within the scope of this invention provided they are heat-stable. Preferably, the optimum temperature ranges from about 50° to about 90°C, more preferably about 60° to about 80°C. In this connection, the term "elevated temperature" as used herein is intended to cover sustained temperatures of operation of the enzyme that are equal to or higher than about 60°C.

The term "template" as used herein refers to a double-stranded or single-stranded DNA molecule which is to be amplified, synthesized, or sequenced. In the case of a double-stranded DNA molecule, denaturation of its strands to form a first and a second strand is performed before these molecules may be amplified, synthesized or sequenced. A primer, complementary to a portion of a DNA template is hybridized under appropriate conditions and the DNA polymerase of the invention may then synthesize a DNA molecule complementary to said template or a portion thereof. The newly synthesized DNA molecule, according to the invention, may be

equal or shorter in length than the original DNA template. Mismatch incorporation during the synthesis or extension of the newly synthesized DNA molecule may result in one or a number of mismatched base pairs. Thus, the synthesized DNA molecule need not be exactly complementary to the DNA template.

5 The term "incorporating" as used herein means becoming a part of a DNA molecule or primer.

As used herein "amplification" refers to any *in vitro* method for increasing the number of copies of a nucleotide sequence, or its complimentary sequence, with the use of a DNA polymerase. Nucleic acid amplification results in
10 the incorporation of nucleotides into a DNA molecule or primer thereby forming a new DNA molecule complementary to a DNA template. The formed DNA molecule and its template can be used as templates to synthesize additional DNA molecules. As used herein, one amplification reaction may consist of many rounds of DNA replication. DNA amplification reactions include, for example, polymerase chain
15 reactions (PCR). One PCR reaction may consist of about 20 to 100 "cycles" of denaturation and synthesis of a DNA molecule. In this connection, the use of the term "long stretches of DNA" as it refers to the extension of primer along DNA is intended to cover such extensions of an average length exceeding 7 kilobases. Naturally, such length will vary, and all such variations are considered to be included within the scope
20 of the invention.

As used herein, the term "holoenzyme" refers to a multi-subunit DNA polymerase activity comprising and resulting from various subunits which each may have distinct activities but which when contained in an enzyme reaction operate to carry out the function of the polymerase (typically DNA synthesis) and enhance its
25 activity over use of the DNA polymerase subunit alone. For example, *E. coli* DNA polymerase III is a holoenzyme comprising three components of one or more subunits each: (1) a core component consisting of a heterotrimer of α , ϵ and θ subunits; (2) a β component consisting of a β subunit dimer; and (3) a γ complex component consisting of a heteropentamer of γ , δ , δ' , χ and ψ subunits (see Studwell and O'Donnell, 1990).
30 These three components, and the various subunits of which they consist, are linked non-covalently to form the DNA polymerase III holoenzyme complex. However, they also function when not linked in solution.

09716964-112100

As used herein, "enzyme complex" refers to a protein structure consisting essentially of two or more subunits of a replication enzyme, which may or may not be identical, noncovalently linked to each other to form a multi-subunit structure. An enzyme complex according to this definition ideally will have a particular enzymatic activity, up to and including the activity of the replication enzyme. For example, a "DNA pol III enzyme complex" as used herein means a multi-subunit protein activity comprising two or more of the subunits of the DNA pol III replication enzyme as defined above, and having DNA polymerizing or synthesizing activity. Thus, this term encompasses the native replication enzyme, as well as an enzyme complex lacking one or more of the subunits of the replication enzyme (e.g., DNA pol III exo-, which lacks the ϵ subunit).

The amino acid residues described herein are preferred to be in the "L" isomeric form. However, residues in the "D" isomeric form can be substituted for any L-amino acid residue, as long as the desired functional property of immunoglobulin-binding is retained by the polypeptide. NH_2 refers to the free amino group present at the amino terminus of a polypeptide. COOH refers to the free carboxy group present at the carboxy terminus of a polypeptide. In keeping with standard polypeptide nomenclature, *J. Biol. Chem.*, 243:3552-59 (1969), abbreviations for amino acid residues are shown in the following Table of Correspondence:

TABLE OF CORRESPONDENCE

<u>1-Letter</u>	<u>SYMBOLS</u>	<u>3-Letter</u>	<u>AMINO ACID</u>
Y		Tyr	tyrosine
G		Gly	glycine
F		Phe	phenylalanine
M		Met	methionine
A		Ala	alanine
S		Ser	serine
I		Ile	isoleucine
L		Leu	leucine
T		Thr	threonine
V		Val	valine
P		Pro	proline
K		Lys	lysine
H		His	histidine
Q		Gln	glutamine
E		Glu	glutamic acid
W		Trp	tryptophan

R	Arg	arginine
D	Asp	aspartic acid
N	Asn	asparagine
C	Cys	cysteine

It should be noted that all amino-acid residue sequences are represented herein by formulae whose left and right orientation is in the conventional direction of amino-terminus to carboxy-terminus. Furthermore, it should be noted that a dash at the beginning or end of an amino acid residue sequence indicates a peptide bond to a further sequence of one or more amino-acid residues. The above Table is presented to correlate the three-letter and one-letter notations which may appear alternately herein.

A "replicon" is any genetic element (e.g., plasmid, chromosome, virus) that functions as an autonomous unit of DNA replication *in vivo*; i.e., capable of replication under its own control.

A "vector" is a replicon, such as plasmid, phage or cosmid, to which another DNA segment may be attached so as to bring about the replication of the attached segment.

A "DNA molecule" refers to the polymeric form of deoxyribonucleotides (adenine, guanine, thymine, or cytosine) in its either single stranded form, or a double-stranded helix. This term refers only to the primary and secondary structure of the molecule, and does not limit it to any particular tertiary forms. Thus, this term includes double-stranded DNA found, *inter alia*, in linear DNA molecules (e.g., restriction fragments), viruses, plasmids, and chromosomes. In discussing the structure of particular double-stranded DNA molecules, sequences may be described herein according to the normal convention of giving only the sequence in the 5' to 3' direction along the nontranscribed strand of DNA (i.e., the strand having a sequence homologous to the mRNA).

An "origin of replication" refers to those DNA sequences that participate in DNA synthesis.

A DNA "coding sequence" is a double-stranded DNA sequence which is transcribed and translated into a polypeptide *in vivo* when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxyl) terminus. A coding sequence can include, but is not limited to, prokaryotic sequences, cDNA from eukaryotic mRNA, genomic DNA sequences

from eukaryotic (e.g., mammalian) DNA, and even synthetic DNA sequences. A polyadenylation signal and transcription termination sequence will usually be located 3' to the coding sequence.

Transcriptional and translational control sequences are DNA regulatory sequences, such as promoters, enhancers, polyadenylation signals, terminators, and the like, that provide for the expression of a coding sequence in a host cell.

A "promoter sequence" is a DNA regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. For purposes of defining the present invention, the promoter sequence is bounded at its 3' terminus by the transcription initiation site and extends upstream (5' direction) to include the minimum number of bases or elements necessary to initiate transcription at levels detectable above background. Within the promoter sequence will be found a transcription initiation site (conveniently defined by mapping with nuclease S1), as well as protein binding domains (consensus sequences) responsible for the binding of RNA polymerase. Eukaryotic promoters will often, but not always, contain "TATA" boxes and "CAT" boxes. Prokaryotic promoters contain Shine-Dalgarno sequences in addition to the -10 and -35 consensus sequences.

An "expression control sequence" is a DNA sequence that controls and regulates the transcription and translation of another DNA sequence. A coding sequence is "under the control" of transcriptional and translational control sequences in a cell when RNA polymerase transcribes the coding sequence into mRNA, which is then translated into the protein encoded by the coding sequence.

A "signal sequence" can be included before the coding sequence. This sequence encodes a signal peptide, N-terminal to the polypeptide, that communicates to the host cell to direct the polypeptide to the cell surface or secrete the polypeptide into the media, and this signal peptide is clipped off by the host cell before the protein leaves the cell. Signal sequences can be found associated with a variety of proteins native to prokaryotes and eukaryotes.

The term "oligonucleotide," as used generally herein, such as in referring to probes prepared and used in the present invention, is defined as a molecule comprised of two or more (deoxy)ribonucleotides, preferably more than

00716964-112100

three. Its exact size will depend upon many factors which, in turn, depend upon the ultimate function and use of the oligonucleotide.

The term "primer" as used herein refers to an oligonucleotide, whether occurring naturally as in a purified restriction digest or produced synthetically, which is capable of acting as a point of initiation of synthesis when placed under conditions in which synthesis of a primer extension product, which is complementary to a nucleic acid strand, is induced, i.e., in the presence of nucleotides and an inducing agent such as a DNA polymerase and at a suitable temperature and pH. The primer may be either single-stranded or double-stranded and must be sufficiently long to prime the synthesis of the desired extension product in the presence of the inducing agent. The exact length of the primer will depend upon many factors, including temperature, source of primer and use of the method. For example, for diagnostic applications, depending on the complexity of the target sequence, the oligonucleotide primer typically contains 15-25 or more nucleotides, although it may contain fewer nucleotides.

The primers herein are selected to be "substantially" complementary to different strands of a particular target DNA sequence. This means that the primers must be sufficiently complementary to hybridize with their respective strands. Therefore, the primer sequence need not reflect the exact sequence of the template. For example, a non-complementary nucleotide fragment may be attached to the 5' end of the primer, with the remainder of the primer sequence being complementary to the strand. Alternatively, non-complementary bases or longer sequences can be interspersed into the primer, provided that the primer sequence has sufficient complementarity with the sequence of the strand to hybridize therewith and thereby form the template for the synthesis of the extension product.

As used herein, the terms "restriction endonucleases" and "restriction enzymes" refer to bacterial enzymes, each of which cut double-stranded DNA at or near a specific nucleotide sequence.

A cell has been "transformed" by exogenous or heterologous DNA when such DNA has been introduced inside the cell. The transforming DNA may or may not be integrated (covalently linked) into chromosomal DNA making up the genome of the cell. In prokaryotes, yeast, and mammalian cells for example, the transforming DNA may be maintained on an episomal element such as a plasmid. With respect to eukaryotic cells, a stably transformed cell is one in which the

transforming DNA has become integrated into a chromosome so that it is inherited by daughter cells through chromosome replication. This stability is demonstrated by the ability of the eukaryotic cell to establish cell lines or clones comprised of a population of daughter cells containing the transforming DNA. A "clone" is a population of cells
5 derived from a single cell or common ancestor by mitosis. A "cell line" is a clone of a primary cell that is capable of stable growth *in vitro* for many generations.

Two DNA sequences are "substantially homologous" when at least about 75% (preferably at least about 80%, and most preferably at least about 90 or 95%) of the nucleotides match over the defined length of the DNA sequences.

10 Sequences that are substantially homologous can be identified by comparing the sequences using standard software available in sequence data banks, or in a Southern hybridization experiment under, for example, stringent conditions as defined for that particular system. Suitable conditions include those characterized by a hybridization buffer comprising 0.9M sodium citrate ("SSC") buffer at a temperature of about 37°C
15 and washing in SSC buffer at a temperature of about 37°C; and preferably in a hybridization buffer comprising 20% formamide in 0.9M SSC buffer at a temperature of about 42°C and washing with 0.2x SSC buffer at about 42°C. Stringency conditions can be further varied by modifying the temperature and/or salt content of the buffer, or by modifying the length of the hybridization probe as is known to those
20 of skill in the art. Defining appropriate hybridization conditions is within the skill of the art. See, e.g., Maniatis et al., 1982; Glover, 1985; Hames and Higgins, 1984.

It should be appreciated that also within the scope of the present invention are degenerate DNA sequences. By "degenerate" is meant that a different three-letter codon is used to specify a particular amino acid. It is well known in the
25 art that the following codons can be used interchangeably to code for each specific amino acid:

30	Phenylalanine (Phe or F)	UUU or UUC
	Leucine (Leu or L)	UUA or UUG or CUU or CUC or CUA or CUG
	Isoleucine (Ile or I)	AUU or AUC or AUA
	Methionine (Met or M)	AUG
	Valine (Val or V)	GUU or GUC or GUA or GUG
	Serine (Ser or S)	UCU or UCC or UCA or UCG or AGU or AGC

	Proline (Pro or P)	CCU or CCC or CCA or CCG
	Threonine (Thr or T)	ACU or ACC or ACA or ACG
	Alanine (Ala or A)	GCU or GCG or GCA or GCG
	Tyrosine (Tyr or Y)	UAU or UAC
5	Histidine (His or H)	CAU or CAC
	Glutamine (Gln or Q)	CAA or CAG
	Asparagine (Asn or N)	AAU or AAC
	Lysine (Lys or K)	AAA or AAG
	Aspartic Acid (Asp or D)	GAU or GAC
10	Glutamic Acid (Glu or E)	GAA or GAG
	Cysteine (Cys or C)	UGU or UGC
	Arginine (Arg or R)	CGU or CGC or CGA or CGG or AGA or AGG
	Glycine (Gly or G)	GGU or GGC or GGA or GGG
	Tryptophan (Trp or W)	UGG
15	Termination codon	UAA (ochre) or UAG (amber) or UGA (opal)

It should be understood that the codons specified above are for RNA sequences. The corresponding codons for DNA have a T substituted for U.

Mutations can be made, e.g., in SEQ. ID. No. 1, or any of the nucleic acids set forth herein, such that a particular codon is changed to a codon which codes for a different amino acid. Such a mutation is generally made by making the fewest nucleotide changes possible. A substitution mutation of this sort can be made to change an amino acid in the resulting protein in a non-conservative manner (i.e., by changing the codon from an amino acid belonging to a grouping of amino acids having a particular size or characteristic to an amino acid belonging to another grouping) or in a conservative manner (i.e., by changing the codon from an amino acid belonging to a grouping of amino acids having a particular size or characteristic to an amino acid belonging to the same grouping). Such a conservative change generally leads to less change in the structure and function of the resulting protein. A non-conservative change is more likely to alter the structure, activity or function of the resulting protein. The present invention should be considered to include sequences containing conservative changes which do not significantly alter the activity or binding characteristics of the resulting protein.

The following is one example of various groupings of amino acids:

Amino acids with nonpolar R groups

- Alanine
- Valine
- 5 Leucine
- Isoleucine
- Proline
- Phenylalanine
- Tryptophan
- 10 Methionine

Amino acids with uncharged polar R groups

- Glycine
- Serine
- 15 Threonine
- Cysteine
- Tyrosine
- Asparagine
- Glutamine

20

Amino acids with charged polar R groups (negatively charged at pH 6.0)

- Aspartic acid
- Glutamic acid

25 Basic amino acids (positively charged at pH 6.0)

- Lysine
- Arginine
- Histidine (at pH 6.0)

30 Amino acids with phenyl groups:

- Phenylalanine
- Tryptophan
- Tyrosine

05716964-112100

Another grouping may be according to molecular weight (i.e., size of R groups):

	Glycine	75
	Alanine	89
	Serine	105
5	Proline	115
	Valine	117
	Threonine	119
	Cysteine	121
	Leucine	131
10	Isoleucine	131
	Asparagine	132
	Aspartic acid	133
	Glutamine	146
	Lysine	146
15	Glutamic acid	147
	Methionine	149
	Histidine (at pH 6.0)	155
	Phenylalanine	165
	Arginine	174
20	Tyrosine	181
	Tryptophan	204

Particularly preferred substitutions are:

- Lys for Arg and vice versa such that a positive charge may be maintained;
- 25 - Glu for Asp and vice versa such that a negative charge may be maintained;
- Ser for Thr such that a free -OH can be maintained; and
- Gln for Asn such that a free NH₂ can be maintained.

Amino acid substitutions may also be introduced to substitute an amino acid with a particularly preferable property. For example, a Cys may be
30 introduced into a potential site for disulfide bridges with another Cys. A His may be introduced as a particularly “catalytic” site (i.e., His can act as an acid or base and is the most common amino acid in biochemical catalysis). Pro may be introduced because of its particularly planar structure, which induces β -turns in the protein's structure.

Two amino acid sequences are "substantially homologous" when at least about 70% of the amino acid residues (preferably at least about 80%, and most preferably at least about 90 or 95%) are identical, or represent conservative substitutions.

5 A "heterologous" region of the DNA construct is an identifiable segment of DNA within a larger DNA molecule that is not found in association with the larger molecule in nature. Thus, when the heterologous region encodes a mammalian gene, the gene will usually be flanked by DNA that does not flank the mammalian genomic DNA in the genome of the source organism. Another example
10 of a heterologous coding sequence is a construct where the coding sequence itself is not found in nature (e.g., a cDNA where the genomic coding sequence contains introns, or synthetic sequences having codons different than the native gene). Allelic variations or naturally-occurring mutational events do not give rise to a heterologous region of DNA as defined herein.

15 An "antibody" is any immunoglobulin, including antibodies and fragments thereof, that binds a specific epitope. The term encompasses polyclonal, monoclonal, and chimeric antibodies, the last mentioned described in further detail in U.S. Patent Nos. 4,816,397 to Boss et al. and 4,816,567 to Cabilly et al.

20 An "antibody combining site" is that structural portion of an antibody molecule comprised of heavy and light chain variable and hypervariable regions that specifically binds antigen.

 The phrase "antibody molecule" in its various grammatical forms as used herein contemplates both an intact immunoglobulin molecule and an immunologically active portion of an immunoglobulin molecule. Exemplary
25 antibody molecules are intact immunoglobulin molecules, substantially intact immunoglobulin molecules and those portions of an immunoglobulin molecule that contains the paratope, including those portions known in the art as Fab, Fab', F(ab')₂ and F(v), which portions are preferred for use in the therapeutic methods described herein. Fab and F(ab')₂ portions of antibody molecules are prepared by the proteolytic
30 reaction of papain and pepsin, respectively, on substantially intact antibody molecules by methods that are well-known. See for example, U.S. Patent No. 4,342,566 to Theofilopolous et al. Fab' antibody molecule portions are also well-known and are produced from F(ab')₂ portions followed by reduction of the disulfide bonds linking the two heavy chain portions as with mercaptoethanol, and followed by alkylation of

05716964.12100

the resulting protein mercaptan with a reagent such as iodoacetamide. An antibody containing intact antibody molecules is preferred herein.

The phrase "monoclonal antibody" in its various grammatical forms refers to an antibody having only one species of antibody combining site capable of immunoreacting with a particular antigen. A monoclonal antibody thus typically displays a single binding affinity for any antigen with which it immunoreacts. A monoclonal antibody may therefore contain an antibody molecule having a plurality of antibody combining sites, each immunospecific for a different antigen; e.g., a bispecific (chimeric) monoclonal antibody.

A DNA sequence is "operatively linked" to an expression control sequence when the expression control sequence controls and regulates the transcription and translation of that DNA sequence. The term "operatively linked" includes having an appropriate start signal (e.g., ATG) in front of the DNA sequence to be expressed and maintaining the correct reading frame to permit expression of the DNA sequence under the control of the expression control sequence and production of the desired product encoded by the DNA sequence. If a gene that one desires to insert into a recombinant DNA molecule does not contain an appropriate start signal, such a start signal can be inserted in front of the gene.

The term "standard hybridization conditions" refers to salt and temperature conditions substantially equivalent to 5x SSC and 65°C for both hybridization and wash. However, one skilled in the art will appreciate that such "standard hybridization conditions" are dependent on particular conditions including the concentration of sodium and magnesium in the buffer, nucleotide sequence length and concentration, percent mismatch, percent formamide, and the like. Also important in the determination of "standard hybridization conditions" is whether the two sequences hybridizing are RNA-RNA, DNA-DNA or RNA-DNA. Such standard hybridization conditions are easily determined by one skilled in the art according to well known formulae, wherein hybridization is typically 10-20°C below the predicted or determined T_m with washes of higher stringency, if desired.

In its primary aspect, the present invention concerns the identification of a class of DNA Polymerase III-type enzymes or complexes found in thermophilic bacteria such as *Thermus thermophilus* (*T.th.*), *Aquifex aeolicus* (*A.ae.*), *Thermotoga maritima* (*T.ma.*), *Bacillus stearothermophilus* (*B.st.*) and other eubacteria which exhibit the following characteristics, among their properties: the ability to extend a

00716964-112100

primer over a long stretch of ssDNA at elevated temperature, stimulation by its cognate sliding clamp of the type that is assembled on DNA by a clamp loader, accessory subunits that exhibit DNA-stimulated ATPase activity at elevated temperature and/or ionic strength, and an associated 3'-5' exonuclease activity. In a particular aspect, the invention extends to Polymerase III-type enzymes derived from a broad class of thermophilic eubacteria that include polymerases isolated from the thermophilic bacteria *Aquifex aeolicus* (*A.ae.* polymerase) and other members of the *Aquifex* genus; *Thermus thermophilus* (*T.th.* polymerase), *Thermus favus* (*Tfl/Tub* polymerase), *Thermus ruber* (*Tru* polymerase), *Thermus brockianus* (DYNAZYME™ polymerase) and other members of the *Thermus* genus; *Bacillus stearothermophilus* (*Bst* polymerase) and other members of the *Bacillus* genus; *Thermoplasma acidophilum* (*Tac* polymerase) and other members of the *Thermoplasma* genus; and *Thermotoga neapolitana* (*Tne* polymerase; See WO 96/10640 to Chatterjee et al.), *Thermotoga maritima* (*Tma* polymerase; See U.S. Patent No. 5,374,553 to Gelfand et al.), and other members of the *Thermotoga* genus. The particular polymerase discussed herein by way of illustration and not limitation, is the enzyme derived from *T.th.*, *A.ae.*, *T.ma.*, or *B.st.*

Polymerase III-type enzymes covered by the invention include those that may be prepared by purification from cellular material, as described in detail in the Examples *infra*, as well as enzyme assemblies or complexes that comprise the combination of individually prepared enzyme subunits or components. Accordingly, the entire enzyme may be prepared by purification from cellular material, or may be constructed by the preparation of the individual components and their assembly into the functional enzyme. A representative and non-limitative protocol for the preparation of an enzyme by this latter route is set forth in U.S. Patent No. 5,583,026 to O'Donnell, and the disclosure thereof is incorporated herein in its entirety for such purpose.

Likewise, individual subunits may be modified, e.g. as by incorporation therein of single residue substitutions to create active sites therein, for the purpose of imparting new or enhanced properties to enzymes containing the modified subunits (see, e.g., Tabor, 1995). Likewise, individual subunits prepared in accordance with the invention, may be used individually and for example, may be substituted for their counterparts in other enzymes, to improve or particularize the

properties of the resultant modified enzyme. Such modifications are within the skill of the art and are considered to be included within the scope of the present invention.

Accordingly, the invention includes the various subunits that may comprise the enzymes, and accordingly extends to the genes and corresponding
5 proteins that may be encoded thereby, such as the α (as well as PolC), β , γ , ϵ , τ , δ and δ' subunits, respectively. More particularly, in *Thermus thermophilus* the α subunit corresponds to *dnaE*, the β subunit corresponds to *dnaN*, the ϵ subunit corresponds to *dnaQ*, and the γ and τ subunits correspond to *dnaX*, the δ subunit corresponds to *holA*, and the δ' subunit corresponds to *holB*. In *Aquifex aeolicus* and *Thermotoga*
10 *maritima*, the α subunit corresponds to *dnaE*, the β subunit corresponds to *dnaN*, the ϵ subunit corresponds to *dnaQ*, the τ subunit corresponds to *dnaX*, the δ subunit corresponds to *holA*, and the δ' subunit corresponds to *holB*. In *Bacillus stearothermophilus*, the PolC which has both α and ϵ activities corresponds to *polC*, the β subunit corresponds to *dnaN*, the ϵ subunit corresponds to *dnaQ*, the τ subunit
15 corresponds to *dnaX*, the δ subunit corresponds to *holA*, and the δ' subunit corresponds to *holB*.

Accordingly, the Polymerase III-type enzyme of the present invention comprises at least one gene encoding a subunit thereof, which gene is selected from the group consisting of *dnaX*, *dnaQ*, *dnaE*, *dnaN*, *holA*, *holB*, and combinations
20 thereof. More particularly, the invention extends to the nucleic acid molecule encoding them and their encoded subunits.

In the *T.th.* Pol III enzyme, this includes the following nucleotide sequences: *dnaX* (SEQ. ID. No. 3), *dnaE* (SEQ. ID. No. 86), *dnaQ* (SEQ. ID. No. 94), *dnaN* (SEQ. ID. No. 106), *holA* (SEQ. ID. No. 157), and *holB* (SEQ. ID. No. 155).

25 In the *A.ae.* Pol III enzyme, this includes the following nucleotide sequences: *dnaX* (SEQ. ID. No. 119), *dnaE* (SEQ. ID. No. 117), *dnaQ* (SEQ. ID. No. 127), *dnaN* (SEQ. ID. No. 121), *holA* (SEQ. ID. No. 123), and *holB* (SEQ. ID. No. 125).

In the *T.ma.* Pol III enzyme, this includes the following nucleotide
30 sequences: *dnaX* (SEQ. ID. No. 141), *dnaE* (SEQ. ID. No. 137), *dnaQ* (SEQ. ID. No. 139), *dnaN* (SEQ. ID. No. 143), *holA* (SEQ. ID. No. 145), and *holB* (SEQ. ID. No. 147).

In the *B.st.* Pol III enzyme, this includes the following nucleotide sequences: *dnaX* (SEQ. ID. No. 181), *dnaN* (SEQ. ID. No. 173), *holA* (SEQ. ID. No. 177), *holB* (SEQ. ID. No. 179), and *polC* (SEQ. ID. Nos. 183).

5 In each of the Pol III type enzymes of the present invention, not only are each of the above-identified coding sequences contemplated, but also conserved variants, active fragments and analogs thereof.

A particular *T.th.* Polymerase III-type enzyme in accordance with the invention may include at least one of the following sub-units: a γ subunit having an amino acid sequence corresponding to SEQ. ID. Nos. 4 and 5; a τ subunit having an amino acid sequence corresponding to SEQ. ID. No. 2; a ϵ subunit having an amino acid sequence corresponding to SEQ. ID. No. 95; a α subunit including an amino acid sequence corresponding to SEQ. ID. No. 87; a β subunit having an amino acid sequence corresponding to SEQ. ID. No. 107; a δ subunit having an amino acid sequence corresponding to SEQ. ID. No. 158; a δ' subunit having an amino acid sequence corresponding to SEQ. ID. No. 156; as well as variants, including allelic variants, muteins, analogs and fragments of any of the subunits, and compatible combinations thereof, capable of functioning in DNA amplification and sequencing.

15 A particular *A.ae.* Polymerase III-type enzyme in accordance with the invention may include at least one of the following sub-units: a τ subunit having an amino acid sequence corresponding to SEQ. ID. No. 120; a ϵ subunit having an amino acid sequence corresponding to SEQ. ID. No. 128; a α subunit including an amino acid sequence corresponding to SEQ. ID. No. 118; a β subunit having an amino acid sequence corresponding to SEQ. ID. No. 122; a δ subunit having an amino acid sequence corresponding to SEQ. ID. No. 124; a δ' subunit having an amino acid sequence corresponding to SEQ. ID. No. 126; as well as variants, including allelic variants, muteins, analogs and fragments of any of the subunits, and compatible combinations thereof, capable of functioning in DNA amplification and sequencing.

25 A particular *T.ma.* Polymerase III-type enzyme in accordance with the invention may include at least one of the following sub-units: a τ subunit having an amino acid sequence corresponding to SEQ. ID. No. 142; a ϵ subunit having an amino acid sequence corresponding to SEQ. ID. No. 140; a α subunit including an amino acid sequence corresponding to SEQ. ID. No. 138; a β subunit having an amino acid sequence corresponding to SEQ. ID. No. 144; a δ subunit having an amino acid

sequence corresponding to SEQ. ID. No. 146; a δ' subunit having an amino acid sequence corresponding to SEQ. ID. No. 148; as well as variants, including allelic variants, muteins, analogs and fragments of any of the subunits, and compatible combinations thereof, capable of functioning in DNA amplification and sequencing.

5 A particular *B.st.* Polymerase III-type enzyme in accordance with the invention may include at least one of the following subunits: a τ subunit having a partial amino acid sequence corresponding to SEQ. ID. No. 182; a β subunit having an amino acid sequence corresponding to SEQ ID. No. 174; a δ subunit having an amino acid sequence corresponding to SEQ. ID. No. 178; a δ' subunit having an amino acid sequence corresponding to SEQ. ID. No. 180; a PolC subunit having an amino acid sequence corresponding to SEQ. ID. Nos. 184; as well as variants, including allelic variants, muteins, analogs and fragments of any of the subunits, and compatible combinations thereof, capable of functioning in DNA amplification and sequencing.

15 The invention also includes and extends to the use and application of the enzyme and/or one or more of its components for DNA molecule amplification and sequencing by the methods set forth hereinabove, and in greater detail later on herein.

20 One of the subunits of the invention is the *T.th.* γ/τ subunit encoded by a *dnaX* gene, which frameshifts as much as -2 with high efficiency, and that, upon frameshifting, leads to the addition of more than one extra amino acid residue to the C-terminus (to form the γ subunit). Further, the invention likewise extends to a *dnaX* gene derived from a thermophile such as *T.th.*, that possesses the frameshift defined herein and that codes for expression of the γ and τ subunits of DNA Polymerase III.

25 The present invention provides methods for amplifying or sequencing a nucleic acid molecule comprising contacting the nucleic acid molecule with a composition comprising a DNA polymerase III enzyme (DNA pol III) complex (for sequencing, preferably a DNA pol III complex that is substantially reduced in 3'-5' exonuclease activity). DNA pol III complexes used in the methods of the present invention are thermostable.

30 The invention also provides DNA molecules amplified by the present methods, methods of preparing a recombinant vector comprising inserting a DNA

molecule amplified by the present methods into a vector, which is preferably an expression vector, and recombinant vectors prepared by these methods.

The invention also provides methods of preparing a recombinant host cell comprising inserting a DNA molecule amplified by the present methods into a host cell, which preferably a bacterial cell, most preferably an *Escherichia coli* cell; a yeast cell; or an animal cell, most preferably an insect cell, a nematode cell or a mammalian cell. The invention also provides and recombinant host cells prepared by these methods.

In additional preferred embodiments, the present invention provides kits for amplifying or sequencing a nucleic acid molecule. DNA amplification kits according to the invention comprise a carrier means having in close confinement therein two or more container means, wherein a first container means contains a DNA polymerase III enzyme complex and a second container means contains a deoxynucleoside triphosphate. DNA sequencing kits according to the present invention comprise a multi-protein Pol III-type enzyme complex and a second container means contains a dideoxynucleoside triphosphate. The DNA pol III contained in the container means of such kits is preferably substantially reduced in 5'-3' exonuclease activity, may be thermostable, and may be isolated from the thermophilic cellular sources described above.

DNA pol III-type enzyme complexes for use in the present invention may be isolated from any organism that produced the DNA pol III-type enzyme complexes naturally or recombinantly. Such enzyme complexes may be thermostable, isolated from a variety of thermophilic organisms.

The thermostable DNA polymerase III-type enzymes or complexes that are an important aspect of this invention, may be isolated from a variety of thermophilic bacteria that are available commercially (for example, from American Type Culture Collection, Rockville, Maryland). Suitable for use as sources of thermostable enzymes are the thermophilic eubacteria *Aquifex aeolicus* and other species of the *Aquifex* genus; *Thermus aquaticus*, *Thermus thermophilus*, *Thermus flavus*, *Thermus ruber*, *Thermus Brockianus*, and other species of the *Thermus* genus; *Bacillus stearothermophilus*, *Bacillus subtilis*, and other species of the *Bacillus* genus; *Thermoplasma acidophilum* and other species of the *Thermoplasma* genus; *Thermotoga neapolitana*, *Thermotoga maritima* and other species of the *Thermotoga* genus; and mutants of each of these species. It will be understood by one of ordinary

09716964-112100

skill in the art, however, that any thermophilic microorganism might be used as a source of thermostable DNA pol III-type enzymes and polypeptides for use in the methods of the present invention. Bacterial cells may be grown according to standard microbiological techniques, using culture media and incubation conditions suitable for growing active cultures of the particular thermophilic species that are well-known to one of ordinary skill in the art (see, e.g., Brock et al., 1969; Oshima et al., 1974). Thermostable DNA pol III complexes may then be isolated from such thermophilic cellular sources as described for thermolabile complexes above.

Several methods are available for identifying homologous nucleic acids and protein subunits in other thermophilic eubacteria, either those listed above or otherwise. These methods include the following:

(1) The following procedure was used to obtain the genes encoding *T.th.* ϵ (*dnaQ*), τ/γ (*dnaX*), DnaA (*dnaA*), and β (*dnaN*). Protein sequences encoded by genes of non-thermophilic bacteria (i.e., mesophiles) are aligned to identify highly conserved amino acid sequences. PCR primers at conserved positions are designed using the codon usage of the organism of interest to amplify an internal section of the gene from genomic DNA extracted from the organism. The PCR product is sequenced. New primers are designed near the ends of the sequence to obtain new sequence that flanks the ends using circular PCR (also called inversed PCR) on genomic DNA that has been cut with the appropriate restriction enzyme and ligated into circles. These new PCR products are sequenced. The procedure is repeated until the entire gene sequence has been obtained. Also, *dnaN* (encoding β) is located next to *dnaA* in bacteria and, therefore, *dnaN* can be obtained by cloning DNA flanking the *dnaA* gene by the circular PCR procedure starting within *dnaA*. Once the gene is obtained, it is cloned into an expression vector for protein production.

(2) The following procedure was used to obtain the genes encoding *T.th.* α polymerase (*dnaE* gene). The DNA polymerase III can be purified directly from the organism of interest and amino acid sequence of the subunit(s) obtained directly. In the case of *T.th.*, *T.th.* cells were lysed and proteins were fractionated. An antibody against *E. coli* α was used to probe column fractions by Western analysis, which reacted with *T.th.* α . The *T.th.* α was transferred to a membrane, proteolyzed, and fragments were sequenced. The sequence was used to design PCR primers for

amplification of an internal section of the *dnaE* gene. Remaining flanking sequences are then obtained by circular PCR.

(3) The following procedure can be used to identify published nucleotide sequences which have not yet been identified as to their function. This method was used to obtain *T.th.* δ (*holA*) and δ' (*holB*), although they could presumably also have been obtained via Methods 1 and 2 above. Discovery of *T.th.* *dnaE* (α), *dnaN* (β) and *dnaX* (τ/γ) indicates that thermophiles use a class III type of DNA polymerase (α) that utilize a clamp (β) and must also use a clamp loader since they have τ/γ . Also, the biochemical experiments in the Examples *infra* show that the *T.th.* polymerase functions with the *T.th.* β clamp. Having demonstrated that a thermophile (e.g., *T.th.*) does indeed utilize a class III type of polymerase with a clamp and clamp loader, it can be assumed that they may have δ and δ' subunits needed to form a complex with τ/γ for functional clamp loading activity (i.e., as shown in *E. coli*, δ and δ' bind either τ or γ to form $\tau\delta\delta'$ or $\gamma\delta\delta'$ complex, both of which are functional clamp loaders). The δ subunit is not very well conserved, but does give a match in the sequence databases for *A.ae.*, *T.ma.*, and *T.th.* The *T.th.* database provided limited information on the amino acid sequence of δ subunit, although one can easily obtain the complete sequence of *T.th.* *holA* by PCR and circular PCR as outlined above in Method 1. The *A.ae.* and *T.ma.* databases are complete and, therefore, the entire *holA* sequence from these genomes are identified. Neither database recognized these sequences as δ encoded by *holA*. The δ' subunit (*holB*) is fairly well conserved. Again the incomplete *T.th.* database provided limited δ' sequence, but as with δ , it is a straight forward process for anyone experienced in the area to obtain the rest of the *holB* sequence using PCR and circular PCR as described in Method 1. Neither the *A.ae.* nor *T.ma.* databases recognized *holB* encoding δ' . Nevertheless, *holB* was identified as encoding δ' by searching the databases with δ' sequence. In each case, the *Thermatoga maritima* and *Aquifex aeolicus* *holB* gene and δ' sequence were obtained in their entirety. Neither database had previously annotated *holA* or *holB* encoding δ and δ' .

As stated above and in accordance with the present invention, once nucleic acid molecules have been obtained, they may be amplified according to any of the literature-described manual or automated amplification methods. Such methods includes, but are not limited to, PCR (U.S. Patent No. 4,683,195 to Mullis et al. and U.S. Patent No. 4,683,202 to Mullis), Strand Displacement Amplification (SDA)

(U.S. Patent No. 5,455,166 to Walker), and Nucleic Acid Sequence-Based Amplification (NASBA) (U.S. Patent No. 5,409,818 to Davey et al.; EP 329,822 to Davey et al.). Most preferably, nucleic acid molecules are amplified by the methods of the present invention using PCR-based amplification techniques.

5 In the initial steps of each of these amplification methods, the nucleic acid molecule to be amplified is contacted with a composition comprising a DNA polymerase belonging to the evolutionary "family A" class (e.g., *Taq* DNA pol I or *E. coli* pol I) or the "family B" class (e.g., Vent and *Pfu* DNA polymerases -- see Ito and Braithwaite, 1991). All of these DNA polymerases are present as single subunits and are primarily involved in DNA repair. In contrast, the DNA pol III-type enzymes are multisubunit complexes that mainly function in the replication of the
10 chromosome, and the subunit containing the DNA polymerase activity is in the "family C" class.

 Thus, in amplifying a nucleic acid molecule according to the methods
15 of the present invention, the nucleic acid molecule is contacted with a composition comprising a thermostable DNA pol III-type enzyme complex.

 Once the nucleic acid molecule to be amplified is contacted with the DNA pol III-type complex, the amplification reaction may proceed according to standard protocols for each of the above-described techniques. Since most of these
20 techniques comprise a high-temperature denaturation step, if a thermolabile DNA pol III-type enzyme complex is used in nucleic acid amplification by any of these techniques the enzyme would need to be added at the start of each amplification cycle, since it would be heat-inactivated at the denaturation step. However, a thermostable DNA pol III-type complex used in these methods need only be added
25 once at the start of the amplification (as for *Taq* DNA polymerase in traditional PCR amplifications), as its activity will be unaffected by the high temperature of the denaturation step. It should be noted, however, that because DNA pol III-type enzymes may have a much more rapid rate of nucleotide incorporation than the polymerases commonly used in these amplification techniques, the cycle times may
30 need to be adjusted to shorter intervals than would be standard.

 In an alternative preferred embodiment, the invention provides methods of extending primers for several kilobases, a reaction that is central to amplifying large nucleic acid molecules, by a technique commonly referred to as "long chain PCR" (Barnes, 1994; Cheng, 1994).

09716964-12100

In such a method the target primed DNA can contain a single strand stretch of DNA to be copied into the double strand form of several or tens of kilobases. The reaction is performed in a suitable buffer, preferably Tris, at a pH of between 5.5 - 9.5, preferably 7.5. The reaction also contains $MgCl_2$ in the range 1 mM to 10 mM, preferably 8 mM, and may contain a suitable salt such as NaCl, KCl or sodium or potassium acetate. The reaction also contains ATP in the range of 20 μM to 1 mM, preferably 0.5 mM, that is needed for the clamp loader to assemble the clamp onto the primed template, and a sufficient concentration of deoxynucleoside triphosphates in the range of 50 μM to 0.5 mM, preferably 60 μM for chain extension.

The reaction contains a sliding clamp, such as the β subunit, in the range of 20ng to 200 ng, preferably 100 ng, for action as a clamp to stimulate the DNA polymerase. The chain extension reaction contains a DNA polymerase and a clamp loader, that could be added either separately or as a single Pol III* -like particle, preferably as a Pol III* like particle that contains the DNA polymerase and clamp loading activities.

The Pol III-type enzyme is added preferably at a concentrations of about 0.0002-200 units per milliliter, about 0.002-100 units per milliliter, about 0.2-50 units per milliliter, and most preferably about 2-50 units per milliliter. The reaction is incubated at elevated temperature, preferably 60°C or more, and could include other proteins to enhance activity such as a single strand DNA binding protein.

In another preferred embodiment, the invention provides methods of extending primers on linear templates in the absence of the clamp loader. In this reaction, the primers are annealed to the linear DNA, preferably at the ends such as in standard PCR applications. The reaction is performed in a suitable buffer, preferably Tris, at a pH of between 5.5 - 9.5, preferably 7.5. The reaction also contains $MgCl_2$ in the range of 1 mM to 10 mM, preferably 8 mM, and may contain a suitable salt such as NaCl, KCl or sodium or potassium acetate. The reaction also contains a sufficient concentration of deoxynucleoside triphosphates in the range of 50 μM to 0.5 mM, preferably 60 μM for chain extension. The reaction contains a sliding clamp, such as the β subunit, in the range of 20ng to 20 μg , preferably about 2 μg , for ability to slide on the end of the DNA and associate with the polymerase for action as a clamp to stimulate the DNA polymerase. The chain extension reaction also contains a Pol III-type polymerase subunit such as α , core, or a Pol III* -like particle. The Pol III-type enzyme is added preferably at a concentrations of about 0.0002-200 units per

09716964-112100

milliliter, about 0.002-100 units per milliliter, about 0.2-50 units per milliliter, and most preferably about 2-50 units per milliliter. The reaction is incubated at elevated temperature, preferably 60°C or more, and could include other proteins to enhance activity such as a single strand DNA binding protein.

5 The methods of the present invention thus will provide high-fidelity amplified copies of a nucleic acid molecule in a more rapid fashion than traditional amplification methods using the repair-type enzymes.

 These amplified nucleic acid molecules may then be manipulated according to standard recombinant DNA techniques. For example, a nucleic acid
10 molecule amplified according to the present methods may be inserted into a vector, which is preferably an expression vector, to produce a recombinant vector comprising the amplified nucleic acid molecule. This vector may then be inserted into a host cell, where it may, for example, direct the host cell to produce a recombinant polypeptide encoded by the amplified nucleic acid molecule. Methods for inserting nucleic acid
15 molecules into vectors, and inserting these vectors into host cells, are well-known to one of ordinary skill in the art (see, e.g., Maniatis, 1992).

 Alternatively, the amplified nucleic acid molecules may be directly inserted into a host cell, where it may be incorporated into the host cell genome or may exist as an extrachromosomal nucleic acid molecule, thereby producing a
20 recombinant host cell. Methods for introduction of a nucleic acid molecule into a host cell, including calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection or other methods, are described in many standard laboratory manuals (see, e.g., Davis, 1986).

 For each of the above techniques wherein an amplified nucleic acid
25 molecule is introduced into a host cell via a vector or via direct introduction, preferred host cells include but are not limited to a bacterial cell, a yeast cell, or an animal cell. Bacterial host cells preferred in the present invention are *E. coli*, *Bacillus* spp., *Streptomyces* spp., *Erwinia* spp., *Klebsiella* spp. and *Salmonella typhimurium*. Preferred as a host cell is *E. coli*, and particularly preferred are *E. coli* strains DH10B
30 and Stbl2, which are available commercially (Life Technologies, Inc. Gaithersburg, Maryland). Preferred animal host cells are insect cells, nematode cells and mammalian cells. Insect host cells preferred in the present invention are *Drosophila* spp. cells, *Spodoptera* Sf9 and Sf21 cells, and *Trichoplusa* High-Five cells, each of which is available commercially (e.g., from Invitrogen; San Diego, California).

09716964.112100

Preferred nematode host cells are those derived from *C. elegans*, and preferred mammalian host cells are those derived from rodents, particularly rats, mice or hamsters, and primates, particularly monkeys and humans. Particularly preferred as mammalian host cells are CHO cells, COS cells and VERO cells.

5 By the present invention, nucleic acid molecules may be sequenced according to any of the literature-described manual or automated sequencing methods. Such methods include, but are not limited to, dideoxy sequencing methods such as "Sanger sequencing" (Sanger and Coulson, 1975; Sanger et al., 1977; U.S. Patent No. 4,962,022 to Fleming et al.; and U.S. Patent No. 5,498,523 to Tabor et al.), as well as
10 more complex PCR-based nucleic acid fingerprinting techniques such as Random Amplified Polymorphic DNA (RAPD) analysis (Williams et al., 1990). Arbitrarily Primed PCR (AP-PCR) (Welsh and McClelland, 1990), DNA Amplification Fingerprinting (DAF) (Caetano-Anollés, 1991), microsatellite PCR or Directed Amplification of Minisatellite-region DNA (DAMD) (Heath et al., 1993), and
15 Amplification Fragment Length Polymorphism (AFLP) analysis (EP 534,858 to Vos et al.; Vos et al., 1995; Lin and Kuo, 1995).

As described above for amplification methods, the nucleic acid molecule to be sequenced by these methods is typically contacted with a composition comprising a type A or type B DNA polymerase. By contrast, in sequencing a nucleic
20 acid molecule according to the methods of the present invention, the nucleic acid molecule is contacted with a composition comprising a thermostable DNA pol III-type enzyme complex instead of necessarily using a DNA polymerase of the family A or B classes. As for amplification methods, the DNA pol III-type complexes used in the nucleic acid sequencing methods of the present invention are preferably
25 substantially reduced in 3'-5' exonuclease activity; most preferable for use in the present methods is a DNA polymerase III-type complex which lacks the ϵ subunit. DNA pol III-type complexes used for nucleic acid sequencing according to the present methods are used at the same preferred concentration ranges described above for long chain extension of primers.

30 Once the nucleic acid molecule to be sequenced is contacted with the DNA pol III complex, the sequencing reactions may proceed according to the protocols disclosed in the above-referenced techniques.

As discussed above, the invention extends to kits for use in nucleic acid amplification or sequencing utilizing DNA polymerase III-type enzymes

09716964.1.21.00

according to the present methods. A DNA amplification kit according to the present invention may comprise a carrier means, such as vials, tubes, bottles and the like. A first such container means may contain a DNA polymerase III-type enzyme complex, and a second such container means may contain a deoxynucleoside triphosphate. The amplification kit encompassed by this aspect of the present invention may further comprise additional reagents and compounds necessary for carrying out standard nucleic amplification protocols (See U.S. Patent No. 4,683,195 to Mullis et al. and U.S. Patent No. 4,683,202 to Mullis, which are directed to methods of DNA amplification by PCR).

Similarly, a DNA sequencing kit according to the present invention comprises a carrier means having in close confinement therein two or more container means, such as vials, tubes, bottles and the like. A first such container means may contain a DNA polymerase III-type enzyme complex, and a second such container means may contain a dideoxynucleoside triphosphate. The sequencing kit may further comprise additional reagents and compounds necessary for carrying out standard nucleic sequencing protocols, such as pyrophosphatase, agarose or polyacrylamide media for formulating sequencing gels, and other components necessary for detection of sequenced nucleic acids (See U.S. Patent No. 4,962,020 to Fleming et al. and U.S. Patent No. 5,498,523 to Tabor et al., which are directed to methods of DNA sequencing).

The DNA polymerase III-type complex contained in the first container means of the amplification and sequencing kits provided by the invention is preferably a thermostable DNA polymerase III-type enzyme complex and more preferably a DNA polymerase III-type enzyme complex that is reduced in 3-5' exonuclease activity. Naturally, the foregoing methods and kits are presented as illustrative and not restrictive of the use and application of the enzymes of the invention for DNA molecule amplification and sequencing. Likewise, the applications of specific embodiments of the enzymes, including conserved variants and active fragments thereof are considered to be disclosed and included within the scope of the invention.

As discussed earlier, individual subunits could be modified to customize enzyme construction and corresponding use and activity. For example, the region of α that interacts with β could be subcloned onto another DNA polymerase, thereby causing β to enhance the activity of the recombinant polymerase.

Alternatively, the β clamp could be modified to function with another protein or enzyme thereby enhancing its activity or acting to localize its action to a particular targeted DNA. Finally, the polymerase active site could be modified to enhance its action, for example changing Tyrosine enabling more equal site stoppage with the
5 four ddNTPs (Tabor et al., 1995). This represents a particular non-limiting illustration of the scope and practice of the present invention with reference to the utility of individual subunits hereof.

Accordingly and as stated above, the present invention also relates to a recombinant DNA molecule or cloned gene, or a degenerate variant thereof, which
10 encodes any one or all of the subunits of the DNA Polymerase III-type enzymes of the present invention, or active fragments thereof. In the instance of the τ subunit, a predicted molecular weight of about 58 kD and an amino acid sequence set forth in SEQ ID Nos. 4 or 5 is comprehended; preferably a nucleic acid molecule, in particular a recombinant DNA molecule or cloned gene, encoding the 58 kD subunit of the
15 Polymerase III of the invention, that has a nucleotide sequence or is complementary to a DNA sequence shown in FIGURES 4A and 4B (SEQ ID No. 1), and the coding region for *dnaX* set forth in FIGURE 4C (SEQ ID No. 3). The γ subunit is smaller, and is approximately 50 kD, depending upon the extent of the frameshift that occurs. More particularly, and as set forth in FIGURE 4E (SEQ ID No. 4), the γ subunit
20 defined by a -1 frameshift possesses a molecular weight of 50.8 kD, while the γ subunit defined by a -2 frameshift, set forth in FIGURE 4F (SEQ ID No. 5), possesses a molecular weight of 49.8 kD.

As discussed above, the invention also extends to the genes including *holA*, *holB*, *dnaX*, *dnaQ*, *dnaE*, and *dnaN* from thermophilic eubacteria (i.e., *T.th.* and
25 *A.ae.*) that have been isolated and/or purified, to corresponding vectors for the genes, and particularly, to the vectors disclosed herein, and to host cells including such vectors. In this connection, probes have been prepared which hybridize to the DNA polymerase III-type enzymes of the present invention, and which are selected from the various oligonucleotide probes or primers set forth in the present application.
30 These include, without limitation, the oligonucleotide defined in SEQ ID No. 6 the oligonucleotide defined in SEQ ID No. 8 the oligonucleotide defined in SEQ ID No. 10 the oligonucleotide defined in SEQ ID No. 11 the oligonucleotide defined in SEQ ID No. 12 the oligonucleotide defined in SEQ ID No. 13 the oligonucleotide defined

09716964-112100

in SEQ ID No. 14 the oligonucleotide defined in SEQ ID No. 15, and the oligonucleotide defined in SEQ ID No. 16.

The methods of the invention include a method for producing a recombinant thermostable DNA polymerase III-type enzyme from a thermophilic bacterium, such as *T.th.*, *A.ae.*, *Th.ma.*, or *B.st.* which comprises culturing a host cell transformed with a vector of the invention under conditions suitable for the expression of the present DNA polymerase III. Another method includes a method for isolating a target DNA fragment consisting essentially of a DNA coding for a thermostable DNA polymerase III-type enzyme from a thermophilic bacterium comprising the steps of:

- 10 (a) forming a genomic library from the bacterium;
- (b) transforming or transfecting an appropriate host cell with the library of step (a);
- (c) contacting DNA from the transformed or transfected host cell with a DNA probe which hybridizes to a DNA fragment selected from the group consisting of the DNA fragments defined in SEQ ID No. 6 and the DNA fragments defined in
15 SEQ ID No. 8 or the oligonucleotides set forth above; wherein hybridization is conducted under the following conditions:
 - i) hybridization: 1% crystalline BSA (fraction V) (Sigma), 1 mM EDTA, 0.5 M NaHPO₄ (pH 7.2), 7% SDS at 65°C for 12 hours and;
 - 20 ii) wash: 5 x 20 minutes with wash buffer consisting of 0.5% BSA, fraction V), 1mM Na₂EDTA, 40 mM NaHPO₄ (pH 7.2), and 5% SDS;
- (d) assaying the transformed or transfected cell of step (c) which hybridizes to the DNA probe for DNA polymerase III-type activity; and
- (e) isolating a target DNA fragment which codes for the thermostable
25 DNA polymerase III-type enzyme.

Also, antibodies including both polyclonal and monoclonal antibodies, and the DNA Polymerase III-like enzyme complex and/or their γ and τ subunits, α subunit(s), δ subunit, δ' subunit, β subunit, ϵ subunit may be used in the preparation of the enzymes of the present invention as well as other enzymes of similar
30 thermophilic origin. For example, the DNA Polymerase III-type complex or its subunits may be used to produce both polyclonal and monoclonal antibodies to themselves in a variety of cellular media, by known techniques such as the hybridoma technique utilizing, for example, fused mouse spleen lymphocytes and myeloma cells.

The general methodology for making monoclonal antibodies by hybridomas is well known. Immortal, antibody-producing cell lines can also be created by techniques other than fusion, such as direct transformation of B lymphocytes with oncogenic DNA, or transfection with Epstein-Barr virus. See, e.g.,

5 Schreier et al., 1980; Hammerling et al., 1981; Kennett et al., 1980; see also U.S. Patent No. 4,341,761 to Ganfield et al.; U.S. Patent No. 4,399,121 to Albarella et al.; U.S. Patent No. 4,427,783 to Newman et al.; U.S. Patent No. 4,444,887 to Hoffman; U.S. Patent No. 4,451,570 to Royston et al.; U.S. Patent No. 4,466,917 to Nussenzweig et al.; U.S. Patent No. 4,472,500 to Milstein et al.; U.S. Patent No.

10 4,491,632 to Wands et al.; and U.S. Patent No. 4,493,890 to Morris.

Methods for producing polyclonal anti-polypeptide antibodies are well-known in the art. See U.S. Patent No. 4,493,795 to Nestor et al. A monoclonal antibody, typically containing Fab and/or F(ab')₂ portions of useful antibody molecules, can be prepared using the hybridoma technology described in *Antibodies -*

15 *A Laboratory Manual*, Harlow and Lane, eds., Cold Spring Harbor Laboratory, New York (1988), which is incorporated herein by reference. Briefly, to form the hybridoma from which the monoclonal antibody composition is produced, a myeloma or other self-perpetuating cell line is fused with lymphocytes obtained from the spleen of a mammal hyperimmunized with an elastin-binding portion thereof.

20 A monoclonal antibody useful in practicing the present invention can be produced by initiating a monoclonal hybridoma culture comprising a nutrient medium containing a hybridoma that secretes antibody molecules of the appropriate antigen specificity. The culture is maintained under conditions and for a time period sufficient for the hybridoma to secrete the antibody molecules into the medium. The

25 antibody-containing medium is then collected. The antibody molecules can then be further isolated by well-known techniques.

Media useful for the preparation of these compositions are both well-known in the art and commercially available and include synthetic culture media, inbred mice and the like. An exemplary synthetic medium is Dulbecco's minimal

30 essential medium (DMEM) (Dulbecco et al., 1959) supplemented with 4.5 gm/l glucose, 20 mm glutamine, and 20% fetal calf serum. An exemplary inbred mouse strain is the Balb/c.

Another feature of this invention is the expression of the DNA sequences disclosed herein. As is well known in the art, DNA sequences may be

05716964-112100

expressed by operatively linking them to an expression control sequence in an appropriate expression vector and employing that expression vector to transform an appropriate unicellular host.

Such operative linking of a DNA sequence of this invention to an
5 expression control sequence, of course, includes, if not already part of the DNA sequence, the provision of an initiation codon, ATG, in the correct reading frame upstream of the DNA sequence.

A wide variety of host/expression vector combinations may be employed in expressing the DNA sequences of this invention. Useful expression
10 vectors, for example, may consist of segments of chromosomal, non-chromosomal and synthetic DNA sequences. Suitable vectors include derivatives of SV40 and known bacterial plasmids, e.g., *E. coli* plasmids col El, pCR1, pBR322, pMB9 and their derivatives, plasmids such as RP4; phage DNAs, e.g., the numerous derivatives of phage λ , e.g., NM989, and other phage DNA, e.g., M13 and filamentous single
15 stranded phage DNA; yeast plasmids such as the 2 μ plasmid or derivatives thereof; vectors useful in eukaryotic cells, such as vectors useful in insect or mammalian cells; vectors derived from combinations of plasmids and phage DNAs, such as plasmids that have been modified to employ phage DNA or other expression control sequences; and the like.

20 Any of a wide variety of expression control sequences -- sequences that control the expression of a DNA sequence operatively linked to it -- may be used in these vectors to express the DNA sequences of this invention. Such useful expression control sequences include, for example, the early or late promoters of SV40, CMV, vaccinia, polyoma or adenovirus, the *lac* system, the *trp* system, the
25 *TAC* system, the *TRC* system, the *LTR* system, the major operator and promoter regions of phage λ , the control regions of fd coat protein, the promoter for 3-phosphoglycerate kinase or other glycolytic enzymes, the promoters of acid phosphatase (e.g., Pho5), the promoters of the yeast α -mating factors, and other sequences known to control the expression of genes of prokaryotic or eukaryotic cells
30 or their viruses, and various combinations thereof.

A wide variety of unicellular host cells are also useful in expressing the DNA sequences of this invention. These hosts may include well known eukaryotic and prokaryotic hosts, such as strains of *E. coli*, *Pseudomonas*, *Bacillus*,

00716964 112100

Streptomyces, fungi such as yeasts, and animal cells, such as CHO, R1.1, B-W and L-M cells, African Green Monkey kidney cells (e.g., COS 1, COS 7, BSC1, BSC40, and BMT10), insect cells (e.g., Sf9), and human cells and plant cells in tissue culture.

It will be understood that not all vectors, expression control sequences
5 and hosts will function equally well to express the DNA sequences of this invention. Neither will all hosts function equally well with the same expression system. However, one skilled in the art will be able to select the proper vectors, expression control sequences, and hosts without undue experimentation to accomplish the desired expression without departing from the scope of this invention. For example, in
10 selecting a vector, the host must be considered because the vector must function in it. The vector's copy number, the ability to control that copy number, and the expression of any other proteins encoded by the vector, such as antibiotic markers, will also be considered.

In selecting an expression control sequence, a variety of factors will
15 normally be considered. These include, for example, the relative strength of the system, its controllability, and its compatibility with the particular DNA sequence or gene to be expressed, particularly with regard to potential secondary structures. Suitable unicellular hosts will be selected by consideration of, e.g., their compatibility with the chosen vector, their secretion characteristics, their ability to fold proteins
20 correctly, and their fermentation requirements, as well as the toxicity to the host of the product encoded by the DNA sequences to be expressed, and the ease of purification of the expression products.

Considering these and other factors a person skilled in the art will be able to construct a variety of vector/expression control sequence/host combinations
25 that will express the DNA sequences of this invention on fermentation or in large scale animal culture.

It is further intended that analogs may be prepared from nucleotide sequences of the protein complex/subunit derived within the scope of the present invention. Analogs, such as fragments, may be produced, for example, by pepsin
30 digestion of bacterial material. Other analogs, such as muteins, can be produced by standard site-directed mutagenesis of *dnaX*, *dnaE*, *dnaQ*, *dnaN*, *holA*, or *holB* coding sequences. Especially useful may be a mutation in *dnaE* that provides the polymerase with the ability to incorporate all four ddNTPs with equal efficiency thereby

03716964.12100

producing an even binding pattern in sequencing gels, as discussed above and with reference to Tabor et al., 1995.

As mentioned above, a DNA sequence corresponding to *dnaX*, *dnaQ*, *holA*, *holB*, *dnaE*, or *dnaN*, or encoding the subunits of the DNA Polymerase III of the invention can be prepared synthetically rather than cloned. The DNA sequence can be designed with the appropriate codons for the amino acid sequence of the subunit(s) of interest. In general, one will select preferred codons for the intended host if the sequence will be used for expression. The complete sequence is assembled from overlapping oligonucleotides prepared by standard methods and assembled into a complete coding sequence (Edge, 1981; Nambair et al., 1984; Jay et al., 1984).

Synthetic DNA sequences allow convenient construction of genes which will express DNA Polymerase III analogs or "muteins". Alternatively, DNA encoding muteins can be made by site-directed mutagenesis of native *dnaX*, *dnaQ*, *holA*, *holB*, *dnaE* or *dnaN* genes or their corresponding cDNAs, and muteins can be made directly using conventional polypeptide synthesis.

A general method for site-specific incorporation of unnatural amino acids into proteins is described in Noren et al., 1989. This method may be used to create analogs with unnatural amino acids.

GENERAL DESCRIPTION OF THE INVENTION

As discussed above, the present invention has as one of its characterizing features, that a Polymerase III-type enzyme as defined hereinabove, has been discovered in a thermophile, that has the structure and function of a chromosomal replicase. This structure and function confers significant benefit when the enzyme is employed in procedures such as PCR where speed and accuracy of DNA reconstruction is crucial.

Chromosomal replicases are composed of several subunits in all organisms (Kornberg and Baker, 1992). In keeping with the need to replicate long chromosomes, replicases are rapid and highly processive multiprotein machines. All cellular replicases examined to date derive their processivity from one subunit that is shaped like a ring and completely encircles DNA (Kuriyan and O'Donnell, 1993; Kelman and O'Donnell, 1994). This "sliding clamp" subunit acts as a mobile tether for the polymerase machine (Stukenberg et al., 1991). The sliding clamp does not

assemble onto the DNA by itself, but requires a complex of several proteins, called a “clamp loader” which couples ATP hydrolysis to the assembly of sliding clamps onto DNA (O'Donnell et al., 1992). Hence, Pol III-type cellular replicases are comprised of three components: a clamp, a clamp loader, and the DNA polymerase.

5 An overall goal is to identify and isolate all of the genes encoding the replicase subunits from a thermophile for expression and purification in large quantity. Following this, the replication apparatus can be reassembled from individual subunit components for use in kits, PCR, sequencing and diagnostic applications (Onrust et al., 1995).

10 As a beginning to identify and characterize the replicase of a thermophile, we started by looking for a homologue to the prokaryotic *dnaX* gene which encode subunits (γ and τ) of the replicase. The *dnaX* gene has another homologue, *holB*, which encodes yet another subunit (δ') of the replicase. The amino acid sequence of δ' (encoded by *holA*) and τ/γ subunits (encoded by *dnaX*) are
15 particularly highly conserved in evolution from prokaryotes to eukaryotes (Chen et al., 1992; O'Donnell et al., 1993; Onrust et al., 1993; Carter et al., 1993; Cullman et al., 1995).

One organism chosen for study and exposition herein is the exemplary extreme thermophile *Thermus thermophilus* (*T.th.*). It is understood that other
20 members of the class such as the eubacterium *Thermatoga* are expected to be analogous in both structure and function. Thus, the investigation of *T.th.* proceeded and initially, a *T.th.* homologue of *dnaX* was identified. The gene encodes a full length protein of 529 amino acids. The amino terminal third of the sequence shares over 50% homology to *dnaX* genes as divergent as *E. coli* (gram negative) and *B.*
25 *subtilis* (gram positive). The *T.th. dnaX* gene contains a DNA sequence that provides a translational frameshift signal for production of two proteins from the same gene. Such frameshifting has been documented only in the case of *E. coli* (Tsuchihashi and Kornberg, 1990; Flower and McHenry, 1990; Blinkowa and Walker, 1990). No frameshifting has been documented to occur in the *dnaX* homologues (RFC subunit
30 genes) of yeast and humans (Eukaryotic kingdom).

The presence of a *dnaX* gene that produces two subunits implies that *T.th.* has a clamp loader (γ) and may be organized by τ into a PolIII*-type replicase like the replicative DNA polymerase of *Escherichia coli*, DNA polymerase III

holoenzyme. The *E. coli* DNA polymerase III holoenzyme contains 10 different subunits, some in copies of two or more for a total composition of 18 polypeptide chains (Kornberg and Baker, 1992; Onrust et al., 1995). The holoenzyme is composed of three major activities: the 3-subunit DNA polymerase core ($\alpha\epsilon\theta$), the β subunit DNA sliding clamp, and the 5-subunit γ complex clamp loader ($\gamma\delta\delta'\chi\psi$). This 3 component strategy generalizes to eukaryotes which utilize a clamp (PCNA) and a 5-subunit RFC clamp loader (RFC) which provide processivity to DNA polymerase δ (reviewed in Kelman and O'Donnell, 1994).

In *E. coli*, the polymerase and clamp loader components are organized into one PolIII* particle by the τ subunit, that acts as a “glue” protein (Onrust et al., 1995). One dimer of τ holds together two core polymerases in the particle which are utilized for the coordinated and simultaneous replication of both strands of duplex DNA (McHenry, 1982; Maki et al., 1988; Yuzhakov et al., 1996). The “glue” protein τ subunit also binds one clamp loader (called γ complex) thereby acting as a scaffold for a large superstructure assembly called DNA polymerase III*. The gene encoding τ , called *dnaX*, also encodes the γ subunit of DNA polymerase III. The β subunit then associates with Pol III* to form the DNA polymerase III holoenzyme. The γ subunit is approximately 2/3 the length of τ . γ shares the N-terminus of τ , but is truncated by a translational frameshifting mechanism that, after the shift, encounters a stop codon within two amino acids (Tsuchihashi and Kornberg, 1990; Flower and McHenry, 1990; Blinkowa and Walker, 1990). Hence, γ is the N-terminal 453 amino acids of τ , but contains one unique residue at the C-terminus (the penultimate codon encodes a Lys residue which is the same sequence as if the frameshift did not take place). This frameshift is highly efficient and occurs approximately 50% of the time.

The sequence of the γ and τ subunits encoded by the *dnaX* gene are homologous to the clamp loading subunits in all other organisms extending from gram negative bacteria through gram positive bacteria, the Archeae Kingdom and the Eukaryotic Kingdom from yeast to humans (O'Donnell et al., 1993). All of these organisms utilize a three component replicase (DNA polymerase, clamp and clamp loader) and in these cases the 3 components appear to behave as independent units in solution rather than forming a large holoenzyme superstructure. For example, in eukaryotes from yeast to humans, the clamp loader is the five subunit RFC, the clamp

is PCNA, and the polymerases δ and ϵ are all stimulated by the PCNA clamp assembled onto primed DNA by RFC (reviewed in Kelman and O'Donnell 1994).

The discovery of a *dnaX* gene in *T.th.* provided confidence that thermophilic bacteria would contain a three component Pol III-type enzyme. Hence, we proceeded to identify the *dnaQ* and *dnaN* genes encoding, respectively, the proofreading 3'-5' exonuclease, and the β DNA sliding clamp subunits of a Pol III-type enzyme. Following this, we purified from extracts of *T.th.* cells, a Pol III-type enzyme. This enzyme preparation had the unique property of extending a single primer around a long 7.2 kb single strand DNA genome of M13mp18 bacteriophage. Such a primer extension assay serves as a tool to detect and identify the Pol III-type of enzyme in cell extracts. The enzyme was confirmed to be a Pol III-type enzyme based on its reactivity with antibody directed against the *E. coli* α subunit (the DNA polymerase subunit) and antibody directed against *E. coli* γ subunit. Proteins corresponding to α , τ , γ , δ and δ' were easily visible and lend themselves to identification of the genes through use of peptide microsequencing followed by primer design for PCR amplification. For example, from this DNA pol III-type preparation, the peptide sequence of the α subunit was obtained, which then allowed the *dnaE* gene encoding the α subunit (DNA polymerase) of the Pol III-type enzyme to be obtain.

These methods should be widely applicable to other thermophilic bacteria. Additional antibody reagents against other Pol III-type enzyme components, such as RFC subunits, DNA polymerase delta, epsilon or beta, and the PCNA clamp from known organisms can be made quite easily as polyclonal or monoclonal antibody preparations using as antigen either naturally purified sequence, recombinant sequence, or synthetic peptide sequence. Examples of known sequences of these Pol III-type enzymes are to be found in: DNA polymerases (Braithwaite and Ito, 1993), RFC clamp loaders (Cullman et al., 1995) and PCNA (Kelman and O'Donnell, 1995).

The remaining genes of *T.th.* Pol III needed for efficient extension of primed templates, *holA* and *holB*, are now identified. The *holA* coding sequence (SEQ. ID. No. 157) encodes the δ subunit (SEQ. ID. No. 158) and the *holB* coding sequence (SEQ. ID. No. 155) encodes the δ' subunit (SEQ. ID. No. 156). The *holA* and *holB* coding sequences and the δ and δ' subunits were identified via BLAST search (Altschul et al., 1997), and subsequently isolated following circular PCR.

These genes will provide the subunit preparations through use of standard recombinant techniques and protein purification protocols. The protein subunits can then be used to reconstitute the enzyme complexes as they exist in the cell. This type of reconstitution of Pol III has been demonstrated using the protein subunits of DNA polymerase III holoenzyme from *E. coli* to assemble the entire particle. See, e.g.,
5 U.S. Patent Nos. 5,583,026 and 5,668,004 to O'Donnell; and Onrust et al., 1995. The disclosures of these references are incorporated herein in their entireties.

Another organism chosen for study and exposition herein is the extreme thermophile *Aquifex aeolicus*. Thus, the present invention also relates to
10 various isolated DNA molecules from *Aquifex aeolicus*, in particular the DNA molecules encoding various replication proteins. These include *dnaE*, *dnaX*, *dnaN*, *holA*, *holB*, *ssb* DNA molecules from *A. aeolicus*. These DNA molecules can be inserted into an expression system or used to transform host cells from which isolated proteins can be obtained. The isolated proteins encoded by these DNA molecules are
15 also disclosed.

Unless otherwise indicated below, the *Aquifex aeolicus* sequences were obtained by sequence comparisons using the *Thermus thermophilus* counterparts as query against the genome of *Aquifex aeolicus* (Deckert et al., 1998).

The *A. aeolicus dnaE* gene has a nucleotide coding sequence according
20 to SEQ. ID. No. 117 and encodes the α subunit of the of DNA Polymerase III, which has an amino acid sequence according to SEQ. ID. No. 118. The *A. ae.* α subunit has approximately 41% aa identity to the *T.th.* α subunit.

The *A. aeolicus dnaX* gene has a nucleotide coding sequence according to SEQ. ID. No. 119 and encodes the τ subunit of the of DNA Polymerase III, which
25 has an amino acid sequence according to SEQ. ID. No. 120. The *A. ae.* τ subunit has approximately 51% aa identity to the *T.th.* τ subunit.

The *A. aeolicus dnaN* gene has a nucleotide coding sequence according to SEQ. ID. No. 121 and encodes the β subunit of DNA Polymerase III, which has an amino acid sequence according to SEQ. ID. No. 122. The *A. ae.* β subunit has
30 approximately 27% aa identity to the *T.th.* β subunit.

The *A. aeolicus dnaQ* gene has a nucleotide coding sequence according to SEQ. ID. No. 127 and encodes the ϵ subunit of the of DNA Polymerase

III, which has an amino acid sequence according to SEQ. ID. No. 128. The *A. ae.* ϵ subunit has approximately 26% aa identity to the *T.th.* ϵ subunit.

The *A. aeolicus ssb* gene has a nucleotide coding sequence according to SEQ. ID. No. 129 and encodes the SSB protein, which has an amino acid sequence according to SEQ. ID. No. 130. The *A. ae* SSB protein has approximately 22% aa identity to the *T.th.* SSB protein.

Further, the coding sequences of *A. aeolicus* genes encoding the helicase (*dnaB*), helicase loader (*dnaC*), and primase (*dnaG*) are also disclosed. The *A. aeolicus dnaB* gene has a nucleotide coding sequence according to SEQ. ID. No. 131 and encodes the DnaB protein, which functions as a helicase and has an amino acid sequence according to SEQ. ID. No. 132. The *A. aeolicus dnaG* gene has a nucleotide coding sequence according to SEQ. ID. No. 133 and encodes the DnaG protein, which functions as a primase and has an amino acid sequence according to SEQ. ID. No. 134. The *A. aeolicus dnaC* gene has a nucleotide coding sequence according to SEQ. ID. No. 135 and encodes the DnaC protein, which functions as a helicase loader and has an amino acid sequence according to SEQ. ID. No. 136.

The *A. aeolicus holA* and *holB* genes were previously unidentified by Deckert et al., 1998. Using *Thermus thermophilus* δ' subunit amino acid sequence and the *Thermatoga maritima* δ subunit amino acid sequence (SEQ. ID. No. 146 which itself was obtained using the *T.th.* δ subunit amino acid sequence of SEQ. ID. No. 158) in separate BLAST searches (Altschul et al., 1997), corresponding polypeptide products in *Aquifex aeolicus* were identified. The *A. aeolicus holA* gene has a nucleotide coding sequence according to SEQ. ID. No. 123 and encodes the δ subunit of the of DNA Polymerase III, which has an amino acid sequence according to SEQ. ID. No. 124. The *A. ae.* δ subunit has approximately 21% aa identity to the *T.m.* δ subunit. The *A. aeolicus holB* gene has a nucleotide coding sequence according to SEQ. ID. No. 125 and encodes the δ' subunit of the of DNA Polymerase III, which has an amino acid sequence according to SEQ. ID. No. 126. The *A. ae.* δ' subunit has approximately 24% aa identity to the *T.th.* δ' subunit.

This invention also clones at least the coding regions of a set of *A. aeolicus* genes which encode proteins that assemble into an *A. aeolicus* DNA polymerase III replication enzyme. These genes (*dnaE*, *dnaN*, *dnaX*, *dnaQ*, *holA*, *holB*, *ssb*) were cloned into expression vectors, the proteins were expressed in *E. coli*,

and the corresponding protein subunits were purified (alpha, beta, tau, delta, delta prime, SSB). This invention identifies the major protein-protein contacts among these subunits, shows how these proteins can be assembled into higher order multiprotein complexes, and how to form a rapid and processive DNA polymerase III holoenzyme.

5 In contrast to the *E. coli* and *T. thermophilus dnaX* genes which encode both τ and γ subunits, the *A. aeolicus dnaX* gene produces only the full length τ subunit when expressed in *E. coli*. The *A. aeolicus* τ is intermediate in length between the γ and τ subunits of *E. coli* DNA polymerase III holoenzyme. The *E. coli* τ binds α , the γ subunit does not bind α . Due to the intermediate size of *A. aeolicus* τ ,
10 it was not known whether the *A. aeolicus* τ would bind the α subunit. This invention shows that indeed, the *A. aeolicus* τ binds to α , as well as δ and δ' , thereby forming an *A. aeolicus* $\alpha\tau\delta\delta'$ complex. Until the identification of the δ and δ' subunits by the present invention, their existence, let alone their interaction with τ and α , was not even known.

15 The *A. aeolicus* $\alpha\tau\delta\delta'/\beta$ Pol III can be applied in several useful DNA handling techniques. For example, the thermophilic Pol III will be useful in DNA sequencing, especially at high temperature. Also, use of a thermal resistant rapid and processive Pol III is an important improvement to polymerase chain reaction technology. The ability of the *A. aeolicus* Pol III to extend primers for multiple
20 kilobases makes possible the amplification of very long segments of DNA (long chain PCR).

 Another organism chosen for study and exposition herein is the extreme thermophile *Thermotoga maritima*. Thus, the present invention also relates to various isolated DNA molecules from *Thermotoga maritima*, in particular the DNA
25 molecules encoding various replication proteins. These include *dnaE*, *dnaX*, *dnaN*, *dnaQ*, *holA*, *holB*, *ssb* DNA molecules from *Thermotoga maritima*. These DNA molecules can be inserted into an expression system or used to transform host cells from which isolated proteins can be obtained. The isolated proteins encoded by these DNA molecules are also disclosed.

30 Unless otherwise indicated below, the *Thermotoga maritima* sequences were obtained by sequence comparisons using the *Thermus thermophilus* counterparts as query against the genome of *Thermotoga maritima* (Nelson et al., 1999).

The *T. maritima dnaE* gene has a nucleotide coding sequence according to SEQ. ID. No. 137 and encodes the α subunit of the of DNA Polymerase III, which has an amino acid sequence according to SEQ. ID. No. 138. The *T.m.* α subunit has approximately 33% aa identity to the *T.th.* α subunit.

5 The *T. maritima dnaQ* gene has a nucleotide coding sequence according to SEQ. ID. No. 139 and encodes the ϵ subunit of the of DNA Polymerase III, which has an amino acid sequence according to SEQ. ID. No. 140. The *T.m.* ϵ subunit has approximately 34% aa identity to the *T.th.* ϵ subunit.

 The *T. maritima dnaX* gene has a nucleotide coding sequence
10 according to SEQ. ID. No. 141 and encodes the τ subunit of the of DNA Polymerase III, which has an amino acid sequence according to SEQ. ID. No. 142. The *T.m.* τ subunit has approximately 48% aa identity to the *T.th.* τ subunit.

 The *T. maritima dnaN* gene has a nucleotide coding sequence according to SEQ. ID. No. 143 and encodes the β subunit of DNA Polymerase III,
15 which has an amino acid sequence according to SEQ. ID. No. 144. The *T.m.* β subunit has approximately 28% aa identity to the *T.th.* β subunit.

 The *T. maritima ssb* gene has a nucleotide coding sequence according to SEQ. ID. No. 149 and encodes the SSB protein, which has an amino acid sequence according to SEQ. ID. No. 150. The *T.m.* SSB protein has approximately 18% aa
20 identity to the *T.th.* SSB protein.

 Further, the coding sequences of *T. maritima* genes encoding the helicase (*dnaB*) and primase (*dnaG*) are also disclosed. The *T. maritima dnaB* gene has a nucleotide coding sequence according to SEQ. ID. No. 151 and encodes the DnaB protein, which functions as a helicase and has an amino acid sequence
25 according to SEQ. ID. No. 152. The *T. maritima dnaG* gene has a nucleotide coding sequence according to SEQ. ID. No. 153 and encodes the DnaG protein, which functions as a primase and has an amino acid sequence according to SEQ. ID. No. 154.

 The *T. maritima hola* and *holB* genes were previously unidentified by
30 Nelson et al., 1999). Using the *Thermus thermophilus* δ and δ' subunit amino acid sequences (SEQ. ID. Nos. 158 and 156, respectively) in separate BLAST searches (Altschul et al., 1997), corresponding polypeptide products in *T. maritima* were identified. The *T. maritima hola* gene has a nucleotide coding sequence according to

SEQ. ID. No. 145 and encodes the δ subunit of the of DNA Polymerase III, which has an amino acid sequence according to SEQ. ID. No. 146. The *T.m.* δ subunit has approximately 37% aa identity to the *T.th.* δ subunit. The *T.m. holB* gene has a nucleotide coding sequence according to SEQ. ID. No. 147 and encodes the δ' subunit
5 which has an amino acid sequence according to SEQ. ID. No. 148. The *T.m.* δ' subunit has approximately 25% aa identity to the *T.th.* δ' subunit.

Yet another organism chosen for study and exposition herein is the extreme thermophile *Bacillus stearothermophilus*. Thus, the present invention also relates to various isolated DNA molecules from *Bacillus stearothermophilus*, in
10 particular the DNA molecules encoding various replication proteins. These include *dnaE*, *dnaX*, *dnaN*, *dnaQ*, *holA*, *holB*, *ssb* DNA molecules from *Bacillus stearothermophilus*. These DNA molecules can be inserted into an expression system or used to transform host cells from which isolated proteins can be obtained. The isolated proteins encoded by these DNA molecules are also disclosed.

15 Unless otherwise indicated below, the *Bacillus stearothermophilus* sequences were obtained by searching the database of this organism (at <http://www.genome.ou.edu>).

The *B. stearothermophilus polC* gene has a nucleotide coding sequence according to SEQ. ID. No. 183 and encodes the PolC or α -large subunit of the DNA
20 Polymerase III, which has an amino acid sequence according to SEQ. ID. No. 184. The *B.st.* PolC subunit, like the PolC subunits of other Gram positive organisms, contains both polymerase and 3'-5' exonuclease activity. This subunit, therefore, is essentially a fusion of α and ϵ .

The *B. stearothermophilus dnaX* gene has a partial nucleotide coding
25 sequence according to SEQ. ID. No. 181 and encodes the τ subunit of the of DNA Polymerase III, which has a partial amino acid sequence according to SEQ. ID. No. 182. The *B.st.* τ subunit has approximately 31% aa identity to the *T.th.* τ subunit.

The *B. stearothermophilus dnaN* gene has a partial nucleotide coding
sequence according to SEQ. ID. No. 173 and encodes the β subunit of DNA
30 Polymerase III, which has a partial amino acid sequence according to SEQ. ID. No. 174. The *B.st.* β subunit has approximately 21% aa identity to the *T.th.* β subunit.

The *B. stearothermophilus ssb* gene has a nucleotide coding sequence according to SEQ. ID. No.175 and encodes the SSB protein, which has an amino acid

sequence according to SEQ. ID. No. 176. The *B.st.* SSB protein has approximately 23% aa identity to the *T.th.* SSB protein.

The *B. stearrowthermophilus holA* gene has a nucleotide coding sequence according to SEQ. ID. No. 177 and encodes the δ subunit of DNA Polymerase III, which has an amino acid sequence according to SEQ. ID. No. 178. The *B.st.* δ subunit has approximately 26% aa identity to the *T.th.* δ subunit.

The *B. stearrowthermophilus holB* gene has a nucleotide coding sequence according to SEQ. ID. No. 179 and encodes the δ' subunit of DNA Polymerase III, which has an amino acid sequence according to SEQ. ID. No. 180. The *B.st.* δ' subunit has approximately 25% aa identity to the *T.th.* δ' subunit.

By conducting BLAST searches of unidentified genomic DNA from other thermophilic eubacteria, it is possible to identify coding regions which encode various functional subunits of other Pol III replicative machinery.

Although it is generally appreciated that proteins isolated from a thermophile should retain activity at high temperature, there is no guarantee that they will retain temperature resistance when isolated in pure form. This invention shows that the *A. aeolicus* Pol III, like the *T. thermophilus* Pol III, is resistant to high temperature. It is expected that the *Th. maritima* and *B. stearrowthermophilus* Pol III enzymes will similarly be resistant to high temperature.

The following experiments illustrate the identification and characterization of the enzymes and constructs of the present invention. Accordingly, in Examples 1-8 below, the identification and expression of the γ and τ is presented, as the first step in the elucidation of the *Thermus thermophilus* Polymerase III reflective of the present invention. Examples 9-12 which follow set forth the protocol for the purification of the remainder of the sub-units of the enzyme that represent substantial entirety of the functional replicative machinery of the enzyme. Examples 18-30 demonstrate the preparation of isolated *A. aeolicus* sequences Pol III subunits and their thermostable use.

EXAMPLE 1

EXPERIMENTAL PROCEDURES

5 Materials

DNA modification enzymes were from New England Biolabs.

Labelled nucleotides were from Amersham, and unlabeled nucleotides were from New England Biolabs. The Alter-1 vector was from Promega. pET plasmids and *E. coli* strains, BL21(DE3) and BL21(DE3)pLysS were from Novagen.

- 10 Oligonucleotides were from Operon. Buffer A is 20mM Tris-HCl (pH 7.5), 0.1mM EDTA, 5mM DTT, and 10% glycerol.

Genomic DNA

Thermus thermophilus (strain HB8) was obtained from the American

- 15 Type Tissue Collection. Genomic DNA was prepared from cells grown in 0.1 l of *Thermus* medium N697 (ATCC: 4 g yeast extract, 8.0 g polypeptone (BBL 11910), 2.0 g NaCl, 30.0 g agar, 1.0 L distilled water) at 75°C overnight. Cells were collected by centrifugation at 4°C and the cell pellet was resuspended in 25 ml of 100 mM Tris-HCl (pH 8.0), 0.05 M EDTA, 2 mg/ml lysozyme and incubated at room
- 20 temperature for 10 min. Then 25 ml 0.10 M EDTA (pH 8.0), 6% SDS was added and mixed followed by 60 ml of phenol. The mixture was shaken for 40 min. followed by centrifugation at 10,000 X G for 10 min. at room temperature. The upper phase (50 ml) was removed and mixed with 50 ml of phenol:chloroform (50:50 v/v) for 30 min. followed by centrifugation for 10 min. at room temperature. The upper phase was
- 25 decanted and the DNA was precipitated upon addition of 1/10th volume 3 M sodium acetate (pH 6.5) and 1 volume ethanol. The precipitate was collected by centrifugation and washed twice with 2 ml of 80% ethanol, dried and resuspended in 1 ml T.E. buffer (10mM Tris Hcl (pH 7.5), 1mM EDTA).

30 Cloning of *dnaX*

DNA oligonucleotides for amplification of *T.th.* genomic DNA were as follows. The upstream 32mer (5'-CGCAAGCTTCACGCSTACCTSTTCTCCGGSAC-3', S indicating a mixture of G and C) (SEQ. ID. No. 6) consists of a Hind III site

within the first 9 nucleotides (underlined) followed by codons (SEQ. ID. No. 29) encoding the following amino acid sequence (HAYLFSGT) (SEQ. ID. No. 7). The downstream 34 mer (5'-CGCGAATTCGTGCTCSGGSGGCTCCTCSAGSGTC-3') (SEQ. ID. No. 8) consists of an EcoRI site (underlined) followed by codons (SEQ. ID. No. 30) encoding the sequence KTLEEPPEH (SEQ. ID. No. 9) on the complementary strand. The amplification reactions contained 10 ng *T.th.* genomic DNA, 0.5 mM of each primer, in a volume of 100 µl of Vent polymerase reaction mixture according to the manufacturers instructions (10 µl ThermoPol Buffer, 0.5 mM each dNTP and 0.5 mM MgSO₄). Amplification was performed using the following cycling scheme: 5 cycles of: 30 sec. at 95.5°C, 30 sec. at 40°C, 2 min. at 72°C; 5 cycles of: 30 sec. at 95.5°C, 30 sec. at 45°C, and 2 min. at 72°C; and 30 cycles of: 30 sec. at 95.5°C, 30 sec. at 50°C, and 30 sec. at 72°C. Products were visualized in a 1.5 % native agarose gel.

Genomic DNA was digested with either XhoI, XbaI, StuI, PstI, NcoI, MluI, KpnI, HindIII, EcoRI, EagI, BglI, or BamHI, followed by Southern analysis in a native agarose gel (Maniatis et al., 1982). Approximately 0.5 µg of digest was analyzed in each lane of a 0.8 % native agarose gel followed by transfer to an MSI filter (Micron Separations Inc.). The transfer included the following steps:

1. The agarose gel was soaked in 500 ml of 1% HCl with gentle shaking for 10 min.
2. Then the gel was soaked in 500 ml of 0.5 M NaOH + 1.5 M NaCl for 40 min.
3. After that the gel was soaked in 500 ml of 1M ammonium acetate for 1 h.
4. The DNA was transferred to the MSI filter with the use of blotting paper for 4 h.
5. The filter was kept at 80°C for 15 min. in the oven.
6. The pre-hybridization step was run in 10 ml of Hybridization solution (1% crystalline BSA (fraction V) (Sigma), 1 mM EDTA, 0.5 M NaHPO₄ (pH 7.2), 7% SDS) at 65°C for 30 min.
7. The probe, radiolabelled by the random priming method (see below), was added to the pre-hybridization solution and kept at 65°C for 12 h.
8. The filter was washed with low stringency with 200 ml of the wash buffer (0.5% BSA, fractionV), 1mM Na₂EDTA, 40 mM NaHPO₄ (pH 7.2), 5% SDS with gentle shaking for 20 min. This step was repeated 5 times, followed by exposure to X-ray film (XAR-5, Kodak).

As a probe, the PCR product was radiolabelled by random as follows.

1. 14 ml of the mixture containing 0.2 μ g of PCR product DNA, 1 μ g of the pd(N6) (Promega) and 2.5 ml of the 10X Klenow reaction buffer (100 mM Tris-HCl (pH 7.5), 50 mM $MgCl_2$, 75 mM dithiothreitol) were boiled for 10 min. and then kept at 4°C.
 2. The reaction volume was increased up to 25 μ l, containing in addition 33 μ M of each dNTP, except dATP, 10 μ Ci [α - 32 P] dATP (800 Ci/mM), and 2 units of Klenow enzyme. The reaction mixture was incubated 1.5 h.
 3. 2 mg of sonicated herring sperm DNA (GibcoBRL) was added to the reaction and the volume was increased to 2 ml using hybridization solution. The sample was then boiled for 10 min.
- 10 A genomic library of XbaI digested DNA was prepared upon treating 1 μ g genomic *T.th.* DNA with 10 units of XbaI in 100 μ l of NEBuffer N2 (50 mM NaCl, 10 mM Tris-HCl (pH 7.9), 10 mM $MgCl_2$, 1 mM DTT) for 2 h at 37°C. The digested DNA was purified by phenol chloroform extraction and ethanol precipitation. The Alter-1 vector (0.5 μ g)(Promega) was digested with 1 unit of XbaI
- 15 in NEBuffer N2 and then purified by phenol/chloroform extraction and ethanol precipitation. One microgram of genomic digest was incubated with 0.05 μ g of digested Alter-1 and 20 U of T4 ligase in 30 μ l of ligase buffer (50 mM Tris-HCl (pH 7.8), 10 mM $MgCl_2$, 10 mM DTT and 1 mM ATP) at 15°C for 12 h. The ligation reaction was transformed into the DH5 α strain of *E. coli* and transformants were
- 20 plated on LB plates containing ampicillin and screened for the *dnaX* insert using the radiolabelled PCR probe as follows:
1. The colonies tested were lifted onto MSI filters, approximately 100 colonies to each filter.
 2. The filters, removed from the LB/Tc plates, were placed side up on a sheet of
 - 25 Whatman 3 MM paper soaked with 0.5 M NaOH for 5 min.
 3. The filters were transferred to a sheet of paper soaked with 1 M Tris-HCl (pH 7.5) for 5 min.
 4. The filters were placed on a sheet of paper soaked in 0.5 M Tris-HCl (pH 7.5), 1.25 M NaCl for 5 min.
 - 30 5. After drying by air, the filters were heated in the oven 80° C for 15 min. and then were analyzed by Southern hybridization.
- Plasmid DNA was prepared from 20 positive colonies; of these 6 contained the expected 4 kb insert when digested with XbaI. Sequencing of the insert was

performed by the Sanger method using the Vent polymerase sequencing kit according to the manufacturers instructions (New England Biolabs).

Identification of the *dnaX* gene

5 The *dnaX* genes of the gram negative *E. coli* and the gram positive *B. subtilis* share more than 50% identity in amino acid sequence within the N-terminal 180 residues containing the ATP-binding domain (Fig. 2). Two highly conserved regions (shown in bold in Fig. 2) were used to design oligonucleotide primers for application of the polymerase chain reaction to *T.th.* genomic DNA. The expected
10 PCR product, including the restriction sites (i.e. before cutting) is 345 nucleotides. Use of these primers with genomic *T.th.* DNA resulted in a product of the expected size. The PCR product was then radiolabelled and used to probe genomic DNA in a Southern analysis (Fig. 3). Genomic DNA was digested with several different restriction endonucleases, electrophoresed in a native agarose gel and then probed
15 with the PCR fragment. The Southern analysis showed an XbaI fragment of approximately 4 kb, more than sufficient length to encode the *dnaX* gene. Other restriction nucleases produced fragments that were significantly longer, or produced two or more fragments indicating presence of a site within the coding sequence of *dnaX*.
20 To obtain full length *dnaX*, genomic DNA was digested with XbaI and ligated into XbaI digested Alter-1 vector. Ligated DNA was transformed into DH5 alpha cells, and colonies were screened with the labeled PCR probe. Plasmid DNA was prepared from 20 positive colonies and analyzed for the appropriate sized insert using XbaI. Six of the twenty clones contained the expected 4 kb XbaI fragment as
25 an insert, the sequence of which is shown in Figs. 4A and 4B.

The frameshift site

 The *dnaX* gene of *E. coli* produces two proteins, the γ and τ subunits, by a -1 frameshift (Tsuchihashi and Kornberg, 1990; Flower and McHenry, 1990;
30 Blinkowa and Walker, 1990). The full length product yields τ , and the frameshift results in addition of one amino acid before encountering a stop codon to produce γ . The -1 frameshift site in the *E. coli dnaX* gene contains the sequence, A AAA AAG, which follows the X XXY YYZ rule found in retroviral genes (Jacks et al., 1988).

This "slippery sequence" preserves the initial two residues of the tRNAs in the aminoacyl and peptidyl sites both before and after the frameshift. Mutagenesis of the *E. coli dnaX* frameshifting site has shown that the first three residues can be nucleotides other than A, but that A's in the second set of three nucleotides is
5 important to frameshifting (Tsuchihashi and Brown, 1992).

Immediately downstream of the stop codon is a potential stem-loop structure which enhances frameshifting, presumably by causing the ribosome to pause. Further, the AAG codon lacks a cognate tRNA in *E. coli* and thus the G residue may facilitate the pause, and has been shown to aid the vigorous frameshifting
10 observed in the *E. coli dnaX* gene (Tsuchihashi and Brown, 1992). A fourth component of frameshifting in the *E. coli dnaX* gene is presence of an upstream Shine-Dalgarno sequence which is thought to pair with the 16S rRNA to increase the frequency of frameshifting still further (Larsen et al., 1994).

Examination of the *T.th. dnaX* sequence reveals a single site that
15 fulfills the X XXY YYZ rule in which positions 4-7 are A residues. The site is unique from that in *E. coli* as all seven residues are A, and the heptanucleotide sequence is flanked by another A residue on each side (i.e. A9). Surprisingly, the stop codon immediately downstream of this site is in the -2 frame, although there is a stop codon in the -1 frame 28 nucleotides downstream of the -2 stop codon. Indeed, a -2
20 frameshift would fulfill the requirement that the first two nucleotides of each codon in the peptidyl and aminoacyl sites be conserved during either a -1 or a -2 frameshift. As with the case of *E. coli dnaX*, there are secondary structure step loop structures immediately downstream. Finally, there is a Shine-Dalgarno sequence immediately adjacent to the frameshift site, as well as another Shine-Dalgarno sequence 22
25 nucleotides upstream of the frameshift site.

Assuming the first stop codon is utilized (i.e. -2 frameshift), the predicted size of the γ subunit in *T.th.* is 454 amino acids for a mass of 49.8 kDa, over 2 kDa larger than the 431 residue γ subunit (47.5 kDa) of *E. coli*. This would result in 2 residues after the -2 frameshift (i.e. after the GluLysLys, the residues LysAla would
30 be added) to be compared to the result of the -1 frameshift in *E. coli* which also results in 2 residues (LysGlu). In the event that a -1 frameshift were utilized in the *T.th. dnaX* gene, then an additional 12 residues would be added following the frameshift for a molecular mass of 50.8 kDa (i.e. after the GluLysLys, the residues

LysProAspProLysAlaProProGlyProThrSer would be added at aa 453-464 of SEQ. ID. No. 4). As explained later, this nucleotide sequence was found to promote both -1 and -2 frameshifting in *E. coli* (Fig. 8). But first, we examined *T.th.* cells by Western analysis for the presence of two subunits homologous to *E. coli* γ and τ .

5

EXAMPLE 2

Frameshifting analysis of the *T.th. dnaX* gene

Frameshifting was analyzed by inserting the frameshift site into lacZ in the three different reading frames, followed by plating on *X-gal* and scoring for blue or white colony formation (Weiss et al., 1987). The frameshifting region within *T.th. dnaX* was subcloned into the EcoRI/BamHI sites of pUC19. These sites are within the polylinker inside of the β -galactosidase gene. Three constructs were produced such that the insert was either in frame with the downstream coding sequence of β -galactosidase, or were out of frame (either -1 or -2). An additional three constructs were designed by mutating the frameshift sequence and then placing this insert into the three reading frames of the β -galactosidase gene. These six plasmids were constructed as described below.

The upstream primer for the shifty sequences was 5'-gcg cgg atc cgg
agg gag aaa aaa gcc tca gcc ca-3' (SEQ. ID. No. 10). The BamHI site for cloning into pUC is underlined. Also, the stop codon, tga, has been mutated to tca (also underlined). The upstream primer for the mutant shifty sequence was: 5'-gcg cgg atc cgg agg gag aga aga aaa gcc tca gcc ca-3' (SEQ. ID. No. 11). The mutant sequence contains two substitutions of a G for an A residue in the polyA stretch (underlined).
Three downstream primers were utilized with each upstream primer to create two sets of three inserts in the 0 frame, -1 frame and -2 frame. The sequence of these primers, and the length of insert (after cutting with EcoRI and BanHI and inserting into pUC19) are as follows: 5'-gaa tta atc tcg cgc ttc ggg agg tgg g-3' (0 frameshift, total 58 nucleotide insert) (SEQ. ID. No. 12); 5'-gcg cga att cgc gct tcg gga ggt ggg-3' (-1 frame, 54mer insert) (SEQ. ID. No. 13); and 5'-gcg cga att cgg gcg ctt cag gag gtg gg-3' (-2 frame, 56mer insert) (SEQ. ID. No. 14). The downstream primers have an EcoRI site (underlined); the EcoRI site of the 0 frame insert was blunt ended to produce the greater length insert (converting the EcoRI site to an aattaatt sequence). Also, the tcg sequence, which produces the tga stop codon (underlined) was mutated

09716964.112100

to tca in the -2 downstream primer so that readthrough would be allowed after the frameshift occurred.

In summary, a region surrounding the frameshift site and ending at least 5 nucleotides past the -1 frameshift stop codon was inserted into the β -galactosidase gene of pUC19 in the three different reading frames (stop codons were mutated to prevent stoppage following a frameshift). These three plasmids were introduced into *E. coli* and plated with *X-gal*. The results, in Fig. 8, show that blue colonies were observed after 24 h incubation with all three plasmids and therefore both -1 and -2 frameshifting had occurred.

To further these results, two γ residues were introduced into the polyA tract which should disrupt the ability of this sequence to direct frameshifts. The mutated slippery sequence was inserted into pUC19 followed by transformation into *E. coli* and plating on *X-gal*. The results showed that both -1 and -2 frameshifting was prevented, further supporting the fact that frameshifting requires the polyA tract as expected (Fig. 8).

EXAMPLE 3

Expression vector for *T.th.* γ and τ

The *dnaX* gene was cloned into the pET16 expression vector in the steps shown in Fig. 9. First, the bulk of the gene was cloned into pET16 by removing the PmlI/XbaI fragment from pAlterdnaX, and placing it into SmaI/XbaI digested Puc19 to yield Puc19dnaXCterm. The N-terminal sequence of the *dnaX* gene was then reconstructed to position an NdeI site at the N-terminus. This was performed by amplifying the 5' region encoding the N-terminal section of γ/τ using an upstream primer containing an NdeI site that hybridizes to the *dnaX* gene at the initiating gtg codon (i.e. to encode Met where the Met is created by the PCR primer, and the Val is the initiating gtg start codon of *dnaX*). The primer sequence for this 5' end was: 5'-gtggtgcatatg gtg agc gcc ctc tac cgc c-3' (SEQ. ID. No. 15) (where the NdeI site is underlined, and the coding sequence of *dnaX* follows). The downstream primer hybridizes past the PmlI site at nucleotide positions 987 - 1004 downstream of the initiating gtg (primer sequence: 5'-gtggtggtcgac cca gga ggg cca cct cca g-3' (SEQ. ID. No. 16) where the initial 12 nucleotides contain a SalGI restriction site, followed

by the sequence from the region downstream the stop codon). The 1.1 kb nucleotide PCR product was digested with PmlI/NdeI and the PmlI/NdeI fragment was ligated into NdeI/PmlI digested Puc19dnaXCterm to form Puc19dnaX. The Puc19dnaX plasmid was then digested with NdeI and SalI and the 1.9 kb fragment containing the *dnaX* gene was purified using the Sephaglas BandPrep Kit (Pharmacia-LKB). pET16b was digested with NdeI and XhoI. Then the full length *dnaX* gene was ligated into the digested pET16b to form pET*dnaX*.

EXAMPLE 4

10

Expression of *T.th.* γ and τ

As discussed in the previous example, the *dnaX* gene was engineered into the T7 based IPTG inducible pET16 vector such that the initiation codon was placed precisely following the Met residue N-terminal leader sequence (Fig. 9). This should produce a protein containing the entire sequence of γ and τ , along with a 21 residue leader containing 10 contiguous His residues (tagged- τ = 60.6 kDa; tagged- γ = 52.4 kDa for -2 frameshift). The pET*dnaX* plasmid was introduced into BL21(DE3)pLysS cells harboring the gene encoding T7 RNA polymerase under control of the lac repressor. Log phase cells were induced with IPTG and analyzed before and after induction in an SDS polyacrylamide gel (Fig. 10, lanes 1 and 2). The result shows that upon induction, two new proteins are expressed with the approximate sizes expected of the *T.th.* γ and τ subunits (larger than *E. coli* γ , and smaller than *E. coli* τ). The two proteins are produced in nearly equal amounts, similar to the case of the *E. coli* γ and τ subunits. Western analysis using antibodies against the *E. coli* γ and τ subunits cross-reacted with the induced proteins further supporting their identity as *T.th.* γ and τ (data not shown, but repeated with the pure subunits shown in Fig. 10, lane 6).

EXAMPLE 5

30

Purification of *T.th.* γ and τ

The His-tagged *T.th.* γ and τ proteins were purified from 6 L of induced *E. coli* cells containing the pET*dnaX* plasmid. Cells were lysed, clarified

from cell debris by centrifugation and the supernatant was applied to a HiTrap chelate affinity column. Elution of the chelate affinity column yielded approximately 35 mg of protein in which the two predominant bands migrated in a region consistent with the molecular weight predicted from the *dnaX* gene (Fig. 10, lane 3), and produced a positive signal by Western analysis using polyclonal antibody directed against the *E. coli* γ and τ subunits (lane 4). The γ and τ subunits are present in nearly equal amounts consistent with the nearly equal expression of these proteins in *E. coli* cells harboring the pET*dnaX* plasmid.

The γ and τ subunits were further purified by gel filtration on a Superose 12 column (Fig. 10, lane 4; Fig. 11). Recovery of *T.th.* γ and τ subunits through gel filtration was 81%. The *E. coli* γ and τ subunits, when separated from one another, elute during gel filtration as tetramers. A mixture of *E. coli* γ/τ results in a mixed tetramer of $\gamma_2\tau_2$ along with γ_4 and τ_4 tetramers (Onrust et al., 1995). The mixture of *T.th.* γ/τ elutes ahead of the 150 kDa marker, and thus is consistent with the expected mass of a $\gamma_2\tau_2$ tetramer (225 kDa) and γ_4 and τ_4 tetramers.

As described earlier, the *dnaX* frameshifting sequence could produce either a -1 or -2 framehift to yield a His-tagged γ subunit of mass either 53.3 kDa or 52.4 kDa, respectively. The difference in these two possible products is too close to determine from migration in SDS gels. It also remains possible that two γ products are present and do not resolve under the conditions used. The exact protocol for this purification is described below.

Six liters of BL21(DE3)pLysSpET*dnaX* cells were grown in LB media containing 50 μ g/ml ampicillin and 25 μ g/ml chloramphenicol at 37°C to an O.D. of 0.8 and then IPTG was added to a concentration of 2 mM. After a further 2 h at 37°C, cells were harvested by centrifugation and stored at -70°C. The following steps were performed at 4°C. Cells (15 g wet weight) were thawed and resuspended in 45 ml 1X binding buffer (5 mM imidazole, 0.5 M NaCl, 20 mM Tris HCl (final pH 7.5)) using a dounce homogenizer to complete cell lysis and 450 ml of 5% polyamine P (Sigma) was added. Cell debris was removed by centrifugation at 18,000 rpm for 30 min. in a Sorvall SS24 rotor at 4°C. The supernatant (Fraction I, 40 ml, 376 mg protein) was applied to a 5 ml HiTrap Chelating Separose column (Pharmacia-LKB). The column was washed with 25 ml of binding buffer, then with 30 ml of binding buffer containing 60 mM imidazole, and then eluted with 30 ml of 0.5 M imidazole, 0.5 M

NaCl, 20 mM Tris-HCl (pH 7.5). Fractions of 1 ml were collected and analyzed on an 8% Coomassie Blue stained SDS polyacrylamide gel. Fractions containing subunits migrating at the *T.th* γ and τ positions, and exhibiting cross reactivity with antibody to *E. coli* γ and τ in a Western analysis, were pooled and dialyzed against
5 buffer A (20 mM Tris-HCl (pH 7.5), 0.1 mM EDTA, 5 mM DTT and 10% glycerol) containing 0.5 M NaCl (Fraction II, 36 mg in 7 ml). Fraction II was diluted 2-fold with buffer A and passed through a 2 ml ATP agarose column equilibrated in buffer A containing 0.2 M NaCl to remove any *E. coli* γ complex contaminant. Then 0.18 mg (300 μ l) Fraction II was gel filtered on a 24 ml Superose 12 column
10 (Pharmacia-LKB) in buffer A containing 0.5 M NaCl. After the first 216 drops, fractions of 200 μ l were collected (Fraction III) and analyzed by Western analysis (by procedures similar to those described in Example 6), by ATPase assays and by Coomassie Blue staining of an 8% Coomassie Blue stained SDS polyacrylamide gel. The Coomassie stained gels and Western analysis of recombinant *T.th.* gamma and
15 tau for these purification steps are summarized in Fig. 10.

EXAMPLE 6

Western Analysis of *T.th.* cells for presence of γ and τ subunits

20 Polyclonal antibody to *E. coli* γ/τ - *E. coli* γ subunit was prepared as described (Studwell-Vaughan and O'Donnell, 1991). Pure γ subunit (100 μ g) was brought up in Freund's adjuvant and injected subcutaneously into a New Zealand Rabbit (Poccono Rabbit Farms). After two weeks, a booster consisting of 50 μ g γ in Freund's adjuvant was administered, followed after two weeks by a third injection (50
25 μ g).

The homology between the amino terminal regions of *T.th.* and *E. coli* γ/τ subunits suggested that there may be some epitopes in common between them. Hence, polyclonal antibody directed against the *E. coli* γ/τ subunits was raised in rabbits for use in probing *T.th.* cells by Western analysis. Fig. 7 shows the results of a
30 Western analysis of whole *T.th.* cells lysed in SDS. The results show that in *T.th.* cells, the antibody is rather specific for two high molecular proteins which migrate in the vicinity of the molecular masses of *E. coli* γ and τ subunits.

Procedure for Western Analysis

Samples were analyzed in duplicate 10 % SDS polyacrylamide gels by the Western method (Towbin et al. 1979). One gel was Coomassie stained to evaluate the pattern of proteins present, and the other gel was then electroblotted onto a nitrocellulose membrane (Schleicher and Schuell). For molecular size markers, the kaliedoscope molecular weight markers (Bio-Rad) were used to verify by visualization that transfer of proteins onto the blotted membrane had occurred. The gel used in electroblotting was also stained after electroblotting to confirm that efficient transfer of protein had occurred. Membranes were blocked using 5% non-fat milk, washed with 0.05% Tween in TBS (TBS-T) and then incubated for over 1 h with a 1/5000 dilution of rabbit polyclonal antibody directed against *E. coli* γ and τ in 1 % gelatin in TBS-T at room temperature. Membranes were washed using TBS-T buffer and then antibody was detected on X-ray film (Kodak) by using the ECL kit from (Amersham) and the manufactures recommended procedures.

Samples included: 1) a mixture of *E. coli* γ (15 ng) and τ (15 ng) subunits; 2) *T.th.* whole cells (100 μ l) suspended in cracking buffer; and 3) purified *T.th.* γ and τ fraction II (0.6 μ g as a mixture).

EXAMPLE 7

Characterization of the ATPase Activity of γ/τ

The *E. coli* τ subunit is a DNA dependent ATPase (Lee and Walker, 1987; Tsuchihashi and Kornberg, 1989). The γ subunit binds ATP but does not hydrolyze it even in the presence of DNA unless other subunits of the DNA polymerase III holoenzyme are also present (Onrust et al., 1991). Next we examined the *T.th.* γ/τ subunits for DNA dependent ATPase activity. The γ/τ preparation was, in fact, a DNA stimulated ATPase (Fig. 11, top panel). The specific activity of the *T.th.* γ/τ was 11.5 mol ATP hydrolyzed/mol γ/τ (as monomer and assuming an equal mixture of the two). Furthermore, analysis of the gel filtration column fractions shows that the ATPase activity coelutes with the *T.th.* γ/τ subunits, supporting evidence that the weak ATPase activity is intrinsic to the γ/τ subunits (Fig. 11). The specific activity of the γ/τ preparation before gel filtration was the same as after gel filtration (within 10%), further indicating that the DNA stimulated ATPase is an

inherent activity of the γ/τ subunits. Presumably, only the τ subunit contains ATPase activity, as in the case of *E. coli*. Assuming only *T.th.* τ contains ATPase activity, its specific activity is twice the observed rate (after factoring out the weight of γ). This rate is still only one-fifth that of *E. coli* τ .

5 The *T.th.* γ/τ ATPase activity is lower at 37°C than at 65°C (middle panel), consistent with the expected behavior of protein activity from a thermophilic source. However, there is no apparent increase in activity in proceeding from 50°C to 65°C (the rapid breakdown of ATP above 65°C precluded measurement of ATPase activity at temperatures above 65°C). In contrast, the *E. coli* τ subunit lost most of its
10 ATPase activity upon elevating the temperature to 50°C (middle panel). These reactions contain no stabilizers such as a nonionic detergent or gelatin, nor did they include substrates such as ATP, DNA or magnesium.

 Last, the relative stability of *T.th.* γ/τ and *E. coli* γ/τ to addition of NaCl (Fig. 12, bottom panel) was examined. Whereas the *E. coli* τ subunit rapidly
15 lost activity at even 0.2 M NaCl, the *T.th.* γ/τ retained full activity in 1.0 M NaCl and was still 80 % active in 1.5 M NaCl. The detailed procedure for the ATPase activity assay is described below.

ATPase assays

20 ATPase assays were performed in 20 μ l of 20 mM Tris-HCl (pH 7.5), 8 mM $MgCl_2$ containing 0.72 μ g of M13mp18 ssDNA (where indicated), 100 mM [γ - ^{32}P]-ATP (specific activity of 2000-4000 cpm/pmol), and the indicated protein. Some reactions contained additional NaCl where indicated. Reactions were incubated at the temperatures indicated in the figure legends for 30 min. and then were
25 quenched with an equal volume of 25 mM EDTA (final). The aliquots were analyzed by spotting them (1 μ l each) onto thin layer chromatography (TLC) sheets coated with Cel-300 polyethyleneimine (Brinkmann Instruments Co.). TLC sheets were developed in 0.5 M lithium chloride, 1 M formic acid. An autoradiogram of the TLC chromatogram was used to visualize Pi at the solvent front and ATP near the origin
30 which were then cut from the TLC sheet and quantitated by liquid scintillation. The extent of ATP hydrolyzed was used to calculate the mol of Pi released per mol of protein per min. One mol of *E. coli* τ was calculated assuming a mass of 71 kDa per monomer. The *T.th.* γ and τ preparation was treated as an equal mixture and thus one

mole of protein as monomer was the average of the predicted masses of the γ and τ subunits (54 kDa).

EXAMPLE 8

5

Homolog of *T.th.* γ/τ to *dnaX* gene products of other organism

The XbaI insert encoded an open reading frame, starting with a GTG codon, of 529 amino acids in length (58.0 kDa), closer to the predicted length of the *B. subtilis* τ subunit (563 amino acids, 62.7 kDa mass)(Alonso et al., 1990) than the *E. coli* τ subunit (71.1 kDa)(Yin et al., 1986). The *dnaX* gene encoding the γ/τ subunits of *E. coli* DNA polymerase III holoenzyme is homologous to the *holB* gene encoding the δ' subunit of the γ complex clamp loader, and this homology extends to all 5 subunits of the eukaryotic RFC clamp loader as well as the bacteriophage gene protein 44 of the gp44/62 clamp loading complex (O'Donnell et al., 1993). These gene products show greatest homology over the N-terminal 166 amino acid residues (of *E. coli dnaX*); the C-terminal regions are more divergent. Fig. 4 shows an alignment of the amino acid sequence of the N-terminal regions of the *T.th. dnaX* gene product to those of several other bacteria. The consensus GXXGXGKT (SEQ. ID. No. 17) motif for nucleotide binding is conserved in all these protein products. Further, the *E. coli* δ' crystal structure reveals one atom of zinc coordinated to four Cys residues (Guenther, 1996). These four Cys residues are conserved in the *E. coli dnaX* gene, and the γ and τ subunits encoded by *E. coli dnaX* bind one atom of zinc. These Cys residues are also conserved in *T.th. dnaX* (shown in Fig. 4). Overall, the level of amino acid identity relative to *E. coli dnaX* in the N-terminal 165 residues of *T.th. dnaX* is 53 %. The *T.th. dnaX* gene is just as homologous to the *B. subtilis dnaX* (53 % identity) gene relative to *E. coli dnaX*. After this region of homology, the C-terminal region of *T.th. dnaX* shares 26% and 20% identity to *E. coli* and *B. subtilis dnaX*, respectively. A proline rich region, downstream of the conserved region, is also present in *T.th. dnaX* (residues 346-375), but not in the *B. subtilis dnaX* (see Figs. 3A and 3B). The overall identity between *E. coli dnaX* and *T.th. dnaX* over the entire gene is 34%. Identity of *T.th. dnaX* to *B. subtilis dnaX* over the entire gene is 28%.

Comparison of *dnaX* genes from *T.th.* and *E. coli*

The above identifies a homologue of the *dnaX* gene of *E. coli* in *Thermus thermophilus*. Like the *E. coli* gene, *T.th. dnaX* encodes two related proteins through use of a highly efficient translational frameshift. The *T.th.* γ/τ subunits are
5 tetramers, or mixed tetramers, similar to the γ and τ subunits of *E. coli*. Further, the γ/τ subunit is a DNA stimulated ATPase like its *E. coli* counterpart. As expected for proteins from a thermophile, the *T.th.* γ/τ ATPase activity is thermostabile and resistant to added salt.

In *E. coli*, γ is a component of the clamp loader, and the τ subunit
10 serves the function of holding the clamp loading apparatus together with two DNA polymerases for coordinated replication of duplex DNA. The presence of γ in *T.th.* suggests it has a clamp loading apparatus and thus a clamp as well. The presence of the τ subunit of *T.th.* implies that *T.th.* contains a replicative polymerase with a structure similar to that of *E. coli* DNA polymerase III holoenzyme.

A significant difference between *E. coli* and *T.th. dnaX* genes is in the
15 translational frameshift sequence. In *E. coli*, the heptamer frameshift site contains six A residues followed by a G residue in the context A AAA AAG. This sequence satisfies the X XXY YYZ rule for -1 frameshifting. The frameshift is made more efficient by the absence of the AAG tRNA for Lys which presumably leads to stalling
20 of the ribosome at the frameshift site and increases the efficiency of frameshifting (Tsuchihashi and Brown, 1992). Two additional aids to frameshifting include a downstream hairpin and an upstream Shine-Dalgarno sequence (Tsuchihashi and Kornberg, 1990; Larsen et al., 1994). The -1 frameshift leads to incorporation of one unique residue at the C-terminus of *E. coli* γ before encounter with a stop codon.

In *T.th.*, the *dnaX* frameshifting heptamer is A AAA AAA, and it is
25 flanked by two other A residues, one on each side. There is also a downstream region of secondary structure. The nearest downstream stop codon is positioned such that gamma would contain only one unique amino acid, as in *E. coli*. However, the *T.th.* stop codon is in the -2 reading frame thus requires a -2 frameshift. No precedent
30 exists in nature for -2 frameshifting, although -2 frameshifting has been shown to occur in test cases (Weiss et al., 1987). *In vivo* analysis of the *T.th.* frameshift sequence shows that this natural sequence promotes both -1 and -2 frameshifting in *E. coli*. Whereas the -2 frameshift results in only one unique C-terminal residue, a -1

frameshift would result in an extension of 12 C-terminal residues. At present, the results do not discriminate which path occurs in *T.th.*, a -1 or -2 frameshift, or a combination of the two.

There are two Shine-Dalgarno sequences just upstream of the
5 frameshift site in *T.th. dnaX*. In two cases of frameshifting in *E. coli*, an upstream Shine-Dalgarno sequence has been shown to stimulate frameshifting (reviewed in Weiss et al., 1897). In release factor 2 (RF2), the Shine-Dalgarno is 3 nucleotides upstream of the shift site, and it stimulates a +1 frameshift event. In the case of *E. coli dnaX*, a Shine-Dalgarno sequence 10 nucleotides upstream of the shift sequence
10 stimulates the -1 frameshift. One of the *T.th. dnaX* Shine-Dalgarno sequences is immediately adjacent to the frameshift sequence with no extra space, the other is 22 residues upstream of the frameshift site. Which of these Shine-Dalgarno sequences plays a role in *T.th. dnaX* frameshifting, if any, will require future study.

In *E. coli*, efficient separation of the two polypeptides, γ and τ , is
15 achieved by mutation of the frameshift site such that only one polypeptide is produced from the gene (Tsuchihashi and Kornberg, 1990). Substitution of G-to-A in two positions of the heptamer of *T.th. dnaX* eliminates frameshifting and thus should be a source to obtain τ subunit free of γ . To produce pure γ subunit free of τ , the frameshifting site and sequence immediately downstream of it can be substituted for
20 an in-frame sequence with a stop codon.

Examination of the *B. subtilis dnaX* gene shows no frameshift sequence that satisfies the X XXY YYZ rule. Hence, it would appear that *dnaX* does not make two proteins in this gram positive organism.

Rapid thermal motions associated with high temperature may make
25 coordination of complicated processes more difficult. It seems possible that organizing the components of the replication apparatus may become yet more important at higher temperature. Hence, production of a τ subunit that could be used to crosslink two polymerases and a clamp loader into one organized particle may be most useful at elevated temperature.

30 As stated above, the following examples describe the continued isolation and purification of the substantial entirety of the Polymerase III from the extreme thermophile *Thermus thermophilus*. It is to be understood that the following exposition is reflective of the protocol and characteristics, both morphological and

functional, of the Polymerase III-type enzymes that are the focus of the present invention, and that the invention is hereby illustrated and comprehends the entire class of enzymes of thermophilic origin.

5

EXAMPLE 9

Purification of the *Thermus thermophilus* DNA polymerase III

All steps in the purification assay were performed at 4°C. The following assay was used in the purification of DNA polymerase from *T.th.* cell
10 extracts. Assays contained 2.5 mg activated calf thymus DNA (Sigma Chemical Company) in a final volume of 25 ml of 20 mM Tris-Cl (pH 7.5), 8 mM MgCl₂, 5 mM DTT, 0.5 mM EDTA, 40 mg/ml BSA, 4% glycerol, 0.5 mM ATP, 3 mM each dCTP, dGTP, dATP, and 20 mM [α -³²P]dTTP. An aliquot of the fraction to be
15 assayed was added to the assay mixture on ice followed by incubation at 60°C for 5 min. DNA synthesis was quantitated using DE81 paper followed by washing off unincorporated nucleotide. Incorporated nucleotide was determined by scintillation counting of the filters.

Thermus thermophilus cell extracts were prepared by suspending 35 grams of cell paste in 200 ml of 50 mM TRIS-HCl, pH=7.5, 30 mM spermidine, 100
20 mM NaCl, 0.5 mM EDTA, 5 mM DTT, 5% glycerol, followed by disruption by passage through a French pressure cell (15,000 PSI). Cell debris was removed by centrifugation (12,000 RPM, 60 min). DNA polymerase III in the clarified supernatant was precipitated by treatment with ammonium sulphate (0.226 gm/liter) and recovered by centrifugation. This fraction was then backwashed with the same
25 buffer (but lacking spermidine) containing 0.20 gm/l ammonium sulfate. The pellet was then resuspended in buffer A and dialyzed overnight against 2 liters of buffer A; a precipitate which formed during dialysis was removed by centrifugation (17,000 RPM, 20 min).

The clarified dialysis supernatant, containing approximately 336 mg of
30 protein, was applied onto a 60 ml heparin agarose column equilibrated in buffer A which was washed with the same buffer until A280 reached baseline. The column was developed with a 500 ml linear gradient of buffer A from 0 to 500 mM NaCl. More tightly adhered proteins were washed off the column by treatment with buffer A

09716964-112100

(20 mM Tris HCl, pH = 7.5, 0.1 mM EDTA, 5mM DTT, and 10% glycerol) and 1M NaCl. Some DNA polymerase activity flowed through the column. Two peaks (HEP.P1 and HEP.P2) of DNA polymerase activity eluted from the heparin agarose column containing 20 mg and 2 mg of total protein respectively (Fig. 13A). These
5 were kept separate throughout the remainder of the purification protocol.

The Pol III resided in HEP.P1 as indicated by the following criteria:

- 1) Western analysis using antibody directed against the α subunit of *E. coli* Pol III indicated presence of Pol III in HEP.P1; 2) Only the HEP.P1 fraction was capable of
10 extending a single primer around an M13mp18 7.2 kb ssDNA circle (explained later in Example 16), such long primer extension being a characteristic of Pol III type enzymes; and 3) Only the HEP.P1 provided DNA polymerase activity that was retained on an ATP-agarose affinity column, which is indicative of a Pol III-type DNA polymerase since the γ and τ subunits are ATP interactive proteins.

The first peak of the heparin agarose column (HEP.P1: 20 mg in 127.5
15 ml) was dialyzed against buffer A and applied onto a 2ml N6-linkage ATP agarose column pre-equilibrated in the same buffer. Bound protein was eluted by a slow (0.05 ml/min) wash with buffer A + 2M NaCl and collected into 200 μ l fractions. Chromatography of peak HEP.P1 yielded a flow-through (HEP.P1-ATP-FT) and a bound fraction (HEP.P1-ATP-Bound) (Fig. 13B). Binding of peak HEP.P2 to the
20 ATP column could not be detected, though DNA polymerase activity was recovered in the flow-through.

The HEP.P1-ATP-Bound fractions from the ATP agarose chromatographic step were further purified by anion exchange over monoQ. The HEP.P1-ATP-Bound fractions were diluted with buffer A to approximately the
25 conductivity of buffer A plus 25 mM NaCl and applied to a 1ml monoQ column equilibrated in Buffer A. DNA polymerase activity eluted in the flow-through and in two resolved chromatographic peaks (MONOQ peak1 and peak2) (Fig. 13C). Peak 2 was by far the major source of DNA polymerase activity. Western analysis using rabbit antibody directed against the *E. coli* α subunit confirmed presence of the α
30 subunit in the second peak (see the Western analysis in Fig. 14B). Antibody against the *E. coli* τ subunit also confirmed the presence of the τ subunit in the second peak. Some reaction against α and τ was also present in the minor peak (first peak). The Coomassie Blue SDS polyacrylamide gel of the MonoQ fractions (Fig. 14A) showed

a band that co-migrated with *E. coli* α and was in the same position as the antibody reactive material (antibody against *E. coli* α). Also present are bands corresponding to τ , γ , δ , and δ' . These subunits, along with β , are all that is necessary for rapid and processive synthesis and primer extension over a long (> 7 kb) stretch of ssDNA in the case of *E. coli* DNA Polymerase III holoenzyme.

The Pol III-type enzyme purified from *T.th.* may be a Pol III*-like enzyme that contains the DNA polymerase and clamp loader subunits (i.e., like the Pol III* of *E. coli*). The evidence for this is: 1) the presence of *dnaX* and *dnaE* gene products in the same column fractions as indicated by Western analysis (see above); 2) the ability of this enzyme to extend a primer around a 7.2 kb circular ssDNA upon adding only β (see Example 16); 3) stimulation of Pol III by adding β on linear DNA, indicating β subunit is not present in saturating amounts (see Example 15); and 4) the presence of τ in *T.th.* which may glue the polymerase and clamp loader into a Pol III* as in *E. coli*; and 5) the comigration of α with subunits τ , γ , δ and δ' of the clamp loader in the column fractions of the last chromatographic step (MonoQ, Fig. 14A).

Micro-sequencing of *T. th* DNA Polymerase III α subunit

The α subunit from the purified *T.th* DNA polymerase III (HEP.P1.ATP-Bound.MONOQ peak2) was blotted onto PVDF membrane and was cut out of the SDS-PAGE gel and submitted to the Protein-Nucleic Acid Facility at Rockefeller University for N-terminal sequencing and proteolytic digestion, purification and microsequencing of the resultant peptides. Analysis of the α candidate band (Mw 130kD) yielded four peptides, two of which (TTH1, TTH2) showed sequence similarity to α subunits from various bacterial sources (see Fig. 15).

EXAMPLE 10

Identification of the *Thermus thermophilus dnaE* gene encoding the α subunit of DNA polymerase III replication enzyme

Cloning of the *dnaE* gene was started with the sequence of the TTH1 peptide from the purified α subunit (FFIEIQNHGLSEQK) (SEQ. ID. No. 61). The fragment was aligned to a region at approximately 180 amino acids downstream of the N-termini of several other known α subunits as shown in Fig. 15. The upstream

5 (SEQ. ID. No. 32) consists of a BamHI site within the first 9 nucleotides (underlined) and the following sequence coding for the peptide FFIEIQNH (SEQ. ID. No. 62).

10

- 15

20

25

30

1. 4 cycles of: 95.5°C – 30 sec., 45°C – 30 sec., 75°C – 8 min.
2. 6 cycles of: 95.5°C – 30 sec., 50°C – 30 sec., 75°C – 6 min.
3. 30 cycles of: 95.5°C – 30 sec., 55°C – 30 sec., 75°C – 5 min.

5 A 1.2kb PCR fragment was obtained and cloned into pUC19:BamHI. The fragment was bracketted by the downstream primer on both sides and contained the region overlapping in 56 bp with the fragment previously cloned.

To obtain yet more *dnaE* sequence, the following primers were used.

The upstream 39mer

(3'-GTGTGGATCCTCGTCCCCCTCATGCGCGACCAGGAAGGG-5') (SEQ. ID.

10 Nos. 35 and 114) consists of a BamHI site within the first 10 nucleotides (underlined) and the sequence from the end of the fragment previously obtained. The downstream 27mer (5'-GTGTGGATCCTTCTTCTTSCCATSGC-3') (SEQ. ID. No. 36) consists of a BamHI site within the first 10 nucleotides (underlined), and the sequence coding for the peptide AMGKKK (SEQ. ID. No. 64) (at position approximately 800 residues from the N terminus) on the complementary strand. The AMGKKK (SEQ. ID. No. 64) sequence was chosen for primer design as it is highly conserved among the known gram-negative α subunits. The amplification reactions contained 10 ng *T.th.* genomic DNA, 0.5 mM of each primer, in a volume of 100 μ l of Taq polymerase reaction mixture containing 10 μ l PCR Buffer, 0.5 mM of each dNTP and 2.5 mM MgCl₂. Amplification was performed using the following cycling scheme:

1. 3 cycles of: 95.5°C – 30 sec., 45°C – 30 sec., 72°C – 8 min.
2. 6 cycles of: 94.5°C – 30 sec., 55°C – 30 sec., 72°C – 6 min.
3. 32 cycles of: 94.5°C – 30 sec., 50°C – 30 sec., 72°C – 5 min.

25 A 2.3kb PCR fragment was obtained instead of the expected 0.6 kb fragment. BamHI digestion of the PCR product resulted in three fragments of 1.1 kb, 0.7kb and 0.5kb. The 1.1 kb fragment was cloned into pUC19:BamHI. It turned out to be the one adjacent to the fragment previously obtained and contained the *dnaE* sequence right up to the region coding for the AMGKKK (SEQ. ID. No. 64) peptide, but was disrupted by an intron just upstream of this region. The sequence that follows this was amplified from the 2.3kb original PCR product using the same conditions and cycling scheme as for the 2.3kb fragment. The downstream primer was the same as in the previous step. The upstream 27mer

(3'-GTGTGGATCCGTGGTGACCTTAGCCAC-5') (SEQ. ID. Nos. 37 and 115)

consisted of a BamHI site within the first 9 nucleotides (underlined) and the sequence from the end of the 1.1kb fragment previously described.

The expected 1.2kb PCR fragment was obtained and cloned into pUC19:SmaI. This fragment coded for the rest of the intein and the end of it was used to obtain the next sequence of *dnaE* downstream of this region. The upstream 30mer (3'-TTCGTGTCCGAGGACCTTGTGGTCCACAAC-5') (SEQ. ID. Nos. 38 and 116) was a sequence from the end of the intron. The downstream 23mer (5'-CCAGAATCGTCTGCTGGTCGTAG-3') (SEQ. ID. No. 39) was the sequence from the end of the *dnaE* gene of *D.rad.* (coding on the complementary strand for the region slightly homologous in the distantly related α subunits and possibly highly homologous between *T.th.* and *D.rad.* α subunits). The amplification reactions contained 10 ng *T.th.* genomic DNA, 0.5 mM of each primer, in a volume of 100 μ l of Vent polymerase reaction mixture containing 10 μ l ThermoPol Buffer, 0.5 mM of each dNTP and 0.1 mM MgSO₄. Amplification was performed using the following cycling scheme:

1. 3 cycles of: 95.5°C – 30 sec., 55°C – 30 sec., 75°C – 8 min.

2. 32 cycles of: 94.5°C – 30 sec., 50°C – 30 sec., 75°C – 5 min.

A 2.5kb PCR fragment was obtained and cloned into pUC19:SmaI. This fragment contained the *dnaE* sequence coding for the 300 amino acids next to the AMGKKK (SEQ. ID. No. 64) region disrupted by yet a second intein inside another sequence that is conserved among the known α subunits (FNKSHSAAY) (SEQ. ID. No. 65).

To obtain the rest of the *dnaE* gene the upstream 19mer (5'-AGCACCTGGAGGAGCTTC-3') (SEQ. ID. No. 40) from the end of the known *dnaE* sequence was used. The downstream primer was: 5'-CATGTCGTACTGGGTGTAC-3' (SEQ. ID. No. 41). The amplification reactions contained 10 ng *T.th.* genomic DNA, 0.5 mM of each primer, in a volume of 100 μ l of Vent polymerase reaction mixture containing 10 μ l ThermoPol Buffer, 0.5 mM of each dNTP and 0.1 mM MgSO₄. Amplification was performed using the following cycling scheme:

1. 3 cycles of: 95.5°C – 30 sec., 55°C – 30 sec., 75°C – 8 min.

2. 32 cycles of: 94.5°C – 30 sec., 50°C – 30 sec., 75°C – 5 min.

A 1.0kb fragment bracketed by this upstream primer was obtained. It contained the 3' end of the *dnaE* gene.

EXAMPLE 11

Cloning and Expression of the *Thermus thermophilus* *dnaQ* gene encoding the ϵ subunit of DNA polymerase III replication enzyme

Cloning of *dnaQ*

The *dnaQ* gene of *E. coli* and the corresponding region of PolC of *B. subtilis*, evolutionary divergent organisms, share approximately 30% identity.

- 10 Comparison of the predicted amino acid sequences for DnaQ (ϵ) of *E. coli* and PolC of *B. subtilis* revealed two highly conserved regions (Fig. 17). Within each of these regions, a nine amino acid sequence was used to design two oligonucleotide primers for use in the polymerase chain reaction.

- The regions highly conservative among Pol III exonucleases were
15 chosen to design the degenerate primers for the amplification of a *T.th. dnaQ* internal fragment (see Fig. 17). DNA oligonucleotides for amplification of *T.th.* genomic DNA were as follows. The upstream 27mer (5'-GTSGTSNNSGACNNSGAGACSACSGGG-3' (SEQ. ID. No. 42)) encodes the following sequence (VVXDXETTG) (SEQ. ID. No. 66). The downstream 27mer
20 (5'-GAASCCSNNGTCGAASNNGGCGTTGTG-3') (SEQ. ID. No. 43) encodes the sequence HNAXFDXGF (SEQ. ID. No. 67) on the complementary strand. The amplification reactions contained 10 ng *T.th.* genomic DNA, 0.5 mM of each primer, in a volume of 100 μ l of Vent polymerase reaction mixture containing 10 μ l ThermoPol Buffer, 0.5 mM of each dNTP and 0.5 mM MgSO₄. Amplification was
25 performed using the following cycling scheme:

1. 5 cycles of: 95.5°C – 30 sec., 40°C – 30 sec., 72°C – 2 min.
2. 5 cycles of: 95.5°C – 30 sec., 45°C – 30 sec., 72°C – 2 min.
3. 30 cycles of: 95.5°C – 30 sec., 50°C – 30 sec., 72°C – 30 min.

- Products were visualized in a 1.5 % native agarose gel. A fragment of the expected
30 size of 270 bp was cloned into the SmaI site of pUC19 and sequenced with the CircumVent Thermal Cycle DNA sequencing kit according to the manufacturer's instructions (New England Biolabs).

To obtain further sequence of the *dnaQ* gene, genomic DNA was digested with either *mho*I, *Bam*HI, *Kpn*I or *Nco*I. These restriction enzymes were chosen because they cut *T.th.* genomic DNA frequently. Approximately 0.1 µg of DNA for each digest was ligated by T4 DNA ligase in 50 µl of ligation buffer (50 mM Tris-HCl (pH 7.8), 10 mM MgCl₂, 10 mM dithiothreitol, 1 mM ATP, 25 mg/ml bovine serum albumin) overnight at 20°C. The ligation mixtures were used for circular PCR.

DNA oligonucleotides for amplification of *T.th.* genomic DNA were the following. The upstream 27mer (5'-CGGGGATCCACCTCAATCACCTCGTGG-3') (SEQ. ID. No. 44) consists of a *Bam*HI site within the first 9 nucleotides (underlined) and the sequence complementary to 42-61bp region of the previously cloned *dnaQ* fragment. The downstream 30mer (5'-CGGGGATCCGCCACCTTGCGGCTCCGGGTG-3') (SEQ. ID. No. 45) consists of a *Bam*HI site within the first 9 nucleotides (underlined) and the sequence corresponding to 240-261 bp region of the *dnaQ* fragment (see Fig. 17).

The amplification reactions contained 1 ng *T.th.* genomic DNA (that had been cut with *Nco*I and religated into circular DNA for circular PCR), 0.4 mM of each primer, in a volume of 100 µl of Vent polymerase reaction mixture containing 10 µl ThermoPol Buffer, 0.5 mM of each dNTP, 0.5 mM MgSO₄, and 10% DMSO.

Circular amplification was performed using the following cycling scheme:

1. 5 cycles of: 95.5°C – 30 sec., 50°C – 30 sec., 72°C – 8 min.
2. 35 cycles of: 95.5°C – 30 sec., 55°C – 30 sec., 72°C – 6 min.
3. 72°C – 10 min.

A 1.5 kb fragment was obtained and cloned into the *Bam*HI site of the pUC19 vector. Partial sequencing of the fragment revealed that it contained the *dnaQ* regions adjacent to sequences corresponding to the PCR primers and hence contained the sequences both upstream and downstream of the previously cloned *dnaQ* fragment. One of *Nco*I sites turned out to be approximately 300 bp downstream of the end of the first cloned *dnaQ* sequence and hence did not include the 3' end of *dnaQ*. To obtain the 3' end, another inverse PCR reaction was performed. Since an *Apa*I restriction site was recognized within this newly sequenced *dnaQ* fragment, the circular PCR procedure was performed using as template an *Apa*I digest of *T.th.* genomic DNA that was ligated (circularized) under the same conditions as described above.

DNA oligonucleotides for amplification of the *Apal*/religated *T.th.* genomic DNA were as follows. The upstream 31mer (5'-GCGCTCTAGACGAGTTCCCAAAGCGTGCGGT-3') (SEQ. ID. No. 46) consists of a *mbaI* site within the first 10 nucleotides (underlined) and the sequence complementary to the region downstream of the *Apal* restriction site in the newly sequenced *dnaQ* fragment. The downstream 25 mer (5'-CGCGTCTAGATCACCTGTATCCAGA-3') (SEQ. ID. No. 47) consists of a *XbaI* site within the first 10 nucleotides (underlined) and the sequence corresponding to another region downstream of the *Apal* restriction site in the newly sequenced *dnaQ* fragment. The 1.7 kb PCR fragment was cloned into the *XbaI* site of the pUC19 vector and partially sequenced. The sequence of *dnaQ*, and the protein sequence of the ϵ subunit encoded by it, is shown in Fig. 18.

The *dnaQ* gene is encoded by an open reading frame of 209 (or 190 depending on which Val is used as the initiating residue) amino acids in length (23598.5 kDa - or 21383.8 kDa for shorter version), similar to the length of the *E. coli* ϵ subunit (243 amino acids, 27099.1 kDa mass) (see Fig. 17).

The entire amino acid sequence of the ϵ subunit predicted from the *T.th. dnaQ* gene aligns with the predicted amino acid sequence of the *dnaQ* genes of other organisms with only a few gaps and insertions (the first two amino acids, and four positions downstream) (Fig. 17). The consensus motifs VVXDXETTG (SEQ. ID. Nos. 66 and 68), HNAXFDXGF (SEQ. ID. No. 67), and HRALYD (SEQ. ID. No. 70), characteristic for exonucleases, are conserved. Overall, the level of amino acid identity relative to most of the known ϵ subunits, or corresponding proofreading exonuclease domains of gram positive PolC genes is approximately 30%. Upstream of start 1 (Fig. 17) there were stop codons in all three reading frames.

Expression of *dnaQ*

The *dnaQ* gene was cloned gene into the pET24-a expression vector in two steps. First, the PCR fragment encoding the N-terminal part of the gene was cloned into the pUC19 plasmid, containing the *Apal* inverse PCR fragment into *NdeI*/*Apal* sites. DNA oligonucleotides for amplification of *T.th.* genomic DNA were as follows. The upstream 33mer (5'-GCGGCGCATATGGTGGTGGTCCTGGACCTGGAG-3') (SEQ. ID. No. 48)

consists of an NdeI site within the first 12 nucleotides (underlined) and the beginning of the *dnaQ* gene. The downstream 25 mer (5'-CGCGTCTAGATCACCTGTATCCAGA-3') (SEQ. ID. No. 49), already used for ApaI circular PCR, consists of an XbaI site within the first 10 nucleotides (underlined) and the sequence corresponding to the region downstream of the ApaI restriction site. The 2.2 kb NdeI/SalI fragment was then cloned into the NdeI/XhoI sites of the pET16 vector to produce pET24-a:*dnaQ*. The ϵ subunit was expressed in the BL21/LysS strain transformed by the pET24-a:*dnaQ* plasmid.

10

EXAMPLE 12

The *Thermus thermophilus* *dnaN* gene encoding the β subunit of DNA polymerase III replication enzyme

15

Strategy of cloning *dnaN* by use of *dnaA*

20

DnaN proteins are highly divergent in bacteria making it difficult to clone them by homology. The level of identity between DnaN representatives from *E. coli* and *B. subtilis* is as low as 18%. These 18% of identical amino acid residues are dispersed through the proteins rather than clustering together in conservative regions, further complicating use of homology to design PCR primers. However, one feature of *dnaN* genes among widely different bacteria is their location in the chromosome. They appear to be near the origin, and immediately adjacent to the *dnaA* gene. The *dnaA* genes show good homology among different bacteria and, thus, *dnaA* was first cloned in order to obtain a DNA probe that is likely near *dnaN*.

25

Identification of *dnaA* and *dnaN*

30

The *dnaA* genes of *E. coli* and *B. subtilis* share 58% identity at the amino acid sequence level within the ATP-binding domain (or among the representatives of gram-positive and gram-negative bacteria, evolutionary divergent organisms). Comparison of the predicted amino acid sequences encoded by *dnaA* of *E. coli* and *B. subtilis* revealed two highly conserved regions (Fig. 19). Within each of these regions, a seven amino acid sequence was used to design two oligonucleotide primers for use in the polymerase chain reaction. The DNA oligonucleotides for amplification of *T.th.* genomic DNA were as follows. The upstream 20mer

(5'-GTSCTSGTSAAGACSCACTT-3') (SEQ. ID. No. 50) encodes the following sequence: VLVKTHL (SEQ. ID. No. 69). The downstream 21mer (5'-SAGSAGSGCGTTGAASGTGTG-3', where S is G or C) (SEQ. ID. No. 51) encodes the sequence: HTFNALL (SEQ. ID. No. 71), on the complementary strand.

5 The amplification reactions contained 10 ng *T.th.* genomic DNA, 0.5 mM of each primer, in a volume of 100 µl of Vent polymerase reaction mixture containing 10 µl ThermoPol Buffer, 0.5 mM of each dNTP and 0.5 mM MgSO₄. Amplification was performed using the following cycling scheme:

1. 5 cycles of: 95.5°C – 30 sec., 45°C – 30 sec., 75°C – 2 min.
- 10 2. 5 cycles of: 95.5°C – 30 sec., 50°C – 30 sec., 75°C – 2 min.
3. 30 cycles of: 95.5°C – 30 sec., 52°C – 30 sec., 75°C – 30 min.

Products were visualized in a 1.5% native agarose gel. A fragment of the expected size of 300 bp was cloned into the SmaI site of pUC19 and sequenced with the CircumVent Thermal Cycle DNA sequencing kit (New England Biolabs).

15 To obtain a larger section of the *T.th.* *dnaA* gene, genomic DNA was digested with either HaeII, HindIII, KasI, KpnI, MluI, NcoI, NgoMI, NheI, NsiI, PaeR7I, PstI, SacI, Sall, SpeI, SphI, StuI, or XhoI, followed by Southern analysis in a native agarose gel. The filter was probed with the 300 bp PCR product radiolabeled by random priming. Four different restriction digests showed a single fragment of
20 reasonable size for further cloning. These were, KasI, NgoMI, and StuI, all of which produced fragments of about 3 kb, and NcoI that produced a 2kb fragment. Also, a KpnI digest resulted in two fragments of about 1.5 kb and 10 kb.

Genomic DNA digests using either NgoMI and StuI were used to obtain the *dnaA* gene by inverse PCR (also referred to as circular PCR). In this
25 procedure, 0.1 µg of DNA from each digest was treated separately with T4 DNA ligase in 50 µl of ligation buffer (50 mM Tris-HCl (pH 7.8), 10 mM MgCl₂, 10 mM dithiothreitol, 1 mM ATP, 25 mg/ml bovine serum albumin) overnight at 20°C. This results in circularizing the genomic DNA fragments. The ligation mixtures were used as substrate in inverse PCR.

30 DNA oligonucleotides for amplification of recircularized *T.th.* genomic DNA were as follows. The upstream 22mer was (5'-CTCGTTGGTGAAAGTTTCCGTG-3') (SEQ. ID. No. 52), and the downstream 24mer was (5'-CGTCCAGTTCATCGCCGGAAGGA-3') (SEQ. ID. No. 53). The

amplification reactions contained 5 ng *T.th.* genomic DNA, 0.5 μ M of each primer, in a volume of 100 μ l of Taq polymerase reaction mixture containing 10 μ l PCR Buffer, 0.5 mM of each dNTP and 2.5 mM $MgCl_2$. Amplification was performed using the following cycling scheme:

- 5 1. 5 cycles of: 95.0°C - 30 sec., 55°C - 30 sec., 72°C – 10 min.
2. 35 cycles of: 95.5°C - 30 sec., 50°C - 30 sec., 72°C – 8 min.

The PCR fragments of the expected length for NgoMI and StuI treated and then ligated chromosomal DNA were digested with either BamHI or Sau3a and cloned into pUC19:BamHI and pUC19:(BamHI+SmaI) and sequenced with CircumVent Thermal
10 Cycle DNA sequencing kit. The 1.6kb (BamHI+BamH) fragment from the NgoMI PCR product contained a sequence coding for the N-terminal part of *dnaN*, followed by the gene for enolase. The 1kb (Sau3a+Sau3a) fragment from the same PCR product included the start of *dnaN* gene and sequence characteristic of the origin of replication (i.e., 9mer DnaA-binding site sequences). The 0.6kb (BamHI+BamHI)
15 fragment from the StuI PCR reaction contained starts for *dnaA* and *gidA* genes in inverse orientation to each other. The 0.4 kb (Sau3a+Sau3a) fragment from the same PCR product contained the 3' end of the *dnaA* gene and DNA sequence characteristic for the origin of replication.

This sequence information provided the beginning and end of both the
20 *dnaA* and the *dnaN* genes. Hence, these genes were easily cloned from this information. Further, the *dnaN* gene was readily cloned and expressed in a pET24-a vector. These steps are described below.

Cloning and sequence of the *dnaA* gene

25 The *dnaA* gene was cloned for sequencing in two parts: from the potential start of the gene up to its middle and from the middle up to the end. For the N-terminal part, the upstream 27mer
(5'-TCTGGCAACACGTTCTGGAGCACATCC-3') (SEQ. ID. No. 54) was 20 bp downstream of the potential start codon of the gene. The downstream 23mer
30 (5'-TGCTGGCGTTCATCTTCAGGATG-3') (SEQ. ID. No. 55) was approximately from the middle of the *dnaA* gene. For the C-terminal part, the upstream 23mer
(5'-CATCCTGAAGATGAACGCCAGCA-3') (SEQ. ID. No. 56) was complementary to the previous primer. The downstream 25mer

(5'-AGGTTATCCACAGGGGTCATGTGCA-3') (SEQ. ID. No. 57) was 20 bp upstream the potential stop codon for the *dnaA* gene. The amplification reactions contained 10 ng *T.th.* genomic DNA, 0.5 μ M of each primer, in a volume of 100 μ l of Vent polymerase reaction mixture containing 10 μ l ThermoPol Buffer, 0.5 mM of each dNTP and 0.5 mM MgSO₄. Amplification was performed using the following cycling scheme:

1. 5 cycles of: 95.5°C - 30 sec., 55°C - 30 sec., 75°C - 3 min.

2. 30 cycles of: 95.5°C - 30 sec., 50°C - 30 sec., 75°C - 2 min.

Products were visualized in a 1.0% native agarose gel. Fragments of the expected sizes of 750 bp and 650 bp were produced, and were sequenced using CircumVent Thermal Cycle DNA sequencing method (New England Biolabs). The nucleotide and amino acid sequences of *dnaA* and its protein product are shown in Fig. 20. The DnaA protein is homologous to the DnaA proteins of several other bacteria as shown in Fig. 19.

Cloning and expression of *dnaN*

The full length *dnaN* gene was obtained by PCR from *T.th.* total DNA. DNA oligonucleotides for amplification of *T.th. dnaN* were the following: the upstream 29mer (5'-GTGTGTCATATGAACATAACGGTCCCAA-3') (SEQ. ID. No. 58) consists of an NdeI site within first 11 nucleotides (underlined), followed by the sequence for the start of the *dnaN* gene; the downstream 29mer (5'-GCGCGAATTCTCCCTTGTGGAAGGCTTAG-3') (SEQ. ID. No. 59) consists of an EcoRI site within the first 10 nucleotides (underlined), followed by the sequence complementary to a section just downstream of the *dnaN* stop codon. The amplification reactions contained 10 ng *T.th.* genomic DNA, 0.5 μ M of each primer, in a volume of 100 μ l of Vent polymerase reaction mixture containing 10 μ l Thermopol Buffer, 0.5 mM of each dNTP and 0.2 mM MgSO₄. Amplification was performed using the following cycling scheme:

1. 5 cycles of: 95.0°C - 30 sec., 55°C - 30 sec., 75°C - 5 min.

2. 35 cycles of: 95.5°C - 30 sec., 50°C - 30 sec., 75°C - 4 min.

The nucleotide and amino acid sequences of *dnaN* and the β subunit, respectively, are shown in Fig. 21. The *T.th.* β subunit shows limited homology to the β subunit sequences of several other bacteria over its entire length (Fig. 22).

The approximately 1 kb *dnaN* gene was cloned into the pET24-a expression vector using the NdeI and EcoRI restriction sites both in the *dnaN* containing PCR product and in pET24-a (Fig. 23). Expression of *T.th.* β subunit was obtained under the following conditions: a fresh colony of BL21(DE3) *E.coli* strain was transformed by the pET24-a:*dnaN* plasmid, and then was grown in LB broth containing 50 mg/ml kanamycin at 37°C until the cell density reached 0.4 OD₆₀₀. The cell culture was then induced for *dnaN* expression upon addition of 2 mM IPTG. Cells were harvested after 4 additional hours of growth under 37°C. The induction of the *T.th.* β subunit is shown in Fig. 24.

Two liters of BL21(DE3)pET*dnaN* cells were grown in LB media containing 50 mg/ml ampicillin at 37°C to an O.D. of 0.8 and then IPTG was added to a concentration of 2 mM. After a further 2 h at 37°C, cells were harvested by centrifugation and stored at -70°C. The following steps were performed at 4°C. Cells were thawed and resuspended in 40 ml of 5 mM Tris-HCl (pH 8.0), 1% sucrose, 1M NaCl, 5 mM DTT, and 30 mM spermidine. Cells were lysed using a French Pressure cell at 20,000 psi. The lysate was allowed to sit at 4°C for 30 min. and then cell debris was removed by centrifugation (Sorvall SS-34 rotor, 45 min. 18,000 rpm). The supernatant was incubated at 65°C for 20 minutes with occasional stirring. The resulting protein precipitate was removed by centrifugation as described above. The supernatant was dialyzed against 4 liters of buffer A containing 50 mM NaCl overnight. The dialyzed supernatant was clarified by centrifugation (35 ml, 150 mg total) and then loaded onto an 8 ml MonoQ column equilibrated in buffer A containing 50 mM NaCl. The column was washed with 5 column volumes of the same buffer and then eluted with a 120 ml gradient of buffer A plus 50 mM NaCl to buffer A plus 500 mM NaCl. Fractions of 2 ml were collected. Over 50 mg of *T.th.* β was recovered in fractions 5-21.

Identification and cloning of *T. thermophilus* *holaA*

TPKGKDLVRHLENRAKRLGLRLPGGVAQYLA-SLEGDLEALERELEKLALLSP
10 -PLTLEKVEKVVALRPPLTGFDLVRSVLEKDPKEALLRLGRLKEEGEELRLL
GALSWQFALLARAFFLLREMPRPKEEDLARLEAHPYAAKKALL-EAARRLTE
EALKEALDALMEAEKRAKG-GKDPWLALAAVLRLAR-PAGQPRVD

20	AGACTCGAGG	CCCTGGAGCG	GGAGCTGGAG	AAGCTTGCCC	TCCTCTCCCC	ACCCCTCACC	60
	CTGGAGAAGG	TGGAGAAGGT	GGTGGCCCTG	AGGCCCCCCC	TCACGGGCTT	TGACCTGGTG	120
	CGCTCCGTCC	TGGAGAAGGA	CCCCAAGGAG	GCCCTCTCTG	GCCTCAGGCG	CCTCAGGGAG	180
	GAGGGGGAGG	AGCCCCCTCAG	GCTCCTCGGG	GCCCTCTCCT	GGCAGTTCGC	CCTCCTCGCC	240
	CGGGCCTTCT	TCCTCCTCCG	GGAAAACCCC	AGGCCCAAGG	AGGAGGACCT	CGCCCGCCTC	300
25	GAGGCCCCACC	CCTACGCCGC	CAAGAAGGCC	A			331

30 RLEALERELEKLALLSPPLTLEKVEKVVALRPPLTGFDLVRVLEKDPKEALL
RLRRLREEGEEPLRLLGALSWQFALLARAFFLLRENPRPKEEDLARLEAHPYA
AKKA

The DNA sequence obtained by PCR (SEQ. ID. No. 188) was used to
35 design internal primers for inverted PCR. The upstream 31mer (5'-

GTGGTGTCTAGACATCATAACGGTTCTGGCA-3') (SEQ. ID. NO. 190) introduced an XbaI site for cloning *holA* into a pGEX vector. The downstream 27mer (5'-GAGGGCCACCACCTTCTCCACCTTCTC-3') (SEQ. ID. No. 191) encodes *holA* sequence EKVEKVVAL (aa residues 159-167 of SEQ. ID. No. 158) on the complementary strand. The amplification reactions contained 50ng *T.th.* genomic DNA and 0.1 μ M of each primer in a volume of 100 μ l of Vent polymerase reaction mixture containing 10 μ l ThermoPol Buffer, 2.5 mM of each dNTP, 2 mM MgSO₄, and 10 μ l of formamide. Amplification was performed using the following cycling scheme:

1. 5 cycles of: 95°C - 30 sec., 65°C - 20 sec., 75°C - 5 min.
2. 5 cycles of: 95°C - 20 sec., 58°C - 10 sec., 75°C - 5 min.
3. 35 cycles of: 95°C - 20 sec., 50°C - 5 sec., 75°C - 4 min.

Products were visualized in a 1.0% native agarose gel. A fragment of 1.5 Kb was gel purified and partially sequenced.

- 15 A different set of primers were used to obtain the 3'-end of *T.th. holA*, including an upstream 25mer (5'-CTCCGTCCTGGAGAAGGACCCCAAG-3') (SEQ. ID. No. 192) which encoded the amino acid sequence SVLEKDPK from *T.th. holA* (aa residues 179-186 of SEQ. ID. No. 158), and a downstream 29mer (5'-CGCGAATTCAACGCCTCCTCAAGACST-3' where S = C or G) (SEQ. ID. No. 193) was not related to the *holA* sequence. The amplification reactions contained 50ng *T.th.* genomic DNA and 0.1 μ M of each primer in a volume of 100 μ l of Vent polymerase reaction mixture containing 10 μ l ThermoPol Buffer, 2.5 mM of each dNTP, and 1-2 mM MgSO₄, and 10 μ l of formamide. Amplification was performed using the following cycling scheme:

- 25 1. 5 cycles of: 95°C - 30 sec., 65°C - 20 sec., 75°C - 5 min.
2. 5 cycles of: 95°C - 20 sec., 55°C - 10 sec., 75°C - 5 min.
3. 35 cycles of: 95°C - 20 sec., 50°C - 5 sec., 75°C - 4 min.

Products were visualized in a 1.0% native agarose gel. A fragment of 1.2 Kb was gel purified and partially sequenced to obtain the remainder of the *T.th. holA* gene.

- 30 The *T.th. holA* gene was cloned into the NdeI/EcoRI sites in the pET24 vector using a pair of primers. The upstream 31mer (5'-GACACTTAACATATGGTCATCGCCTTCACCG-3') (SEQ. ID. No. 194) contains a NdeI site within the first 15 nucleotides (underlined) and has a sequence

corresponding to 5' region of *T.th. holA*. The downstream 38 mer (5'-GTGTGTGAATTCGGGTCAACGGGCGAGGCGGAGGACCG-3') (SEQ. ID. No. 195) contains a EcoRI site within the first 12 nucleotides (underlined) and has a sequence complementary to the 3' end of *holA* gene.

5

EXAMPLE 14

Identification of *T.th. holB* encoding δ' subunit

10

To clone the ends of *T.th. holB* gene, it was assumed that the order of genes in *Thermus thermophilis* could be the same as in related *Deinococcus radiodurance*. Multiple alignment of the upstream neighbor (probable phosphoesterase, DNA repair Rad24c related protein) revealed a conservative region close to the C-terminus of the protein sequence:

15

<i>Deinococcus radiodurance</i>	VILNPGSVGQ	(SEQ. ID. No. 196)
<i>Methanococcus janaschii</i>	YLINPGSVGQ	(SEQ. ID. No. 197)
<i>Thermotoga maritima</i>	LVLNPGSAGR	(SEQ. ID. No. 198)

20

The *D.rad.* sequence was used to design an upstream 28mer primer (5'-CTGGTGAACCCGGGCTCCGTGGGCCAGC-3') (SEQ. ID. No. 199) that encodes the amino acid sequence LLVNPGSVGQ (SEQ. ID. No. 200) and a downstream 27mer (5'-CTCGAGGAGCTTGAGGAGGGTGTGGC-3') (SEQ. ID. No. 201) encodes the sequence ANTLLKLLLE (SEQ. ID. No. 202) on the complementary strand. The amplification reactions contained 50ng *T.th.* genomic DNA and 0.1 μ M of each primer in a volume of 100 μ l of Deep Vent polymerase reaction mixture containing 10 μ l ThermoPol Buffer, 2.5 mM of each dNTP, 1.5 mM MgSO₄, and 10 μ l formamide. Amplification was performed using the following cycling scheme:

30

1. 5 cycles of: 95°C - 30 sec., 68°C - 20 sec., 75°C - 3 min.
2. 5 cycles of: 95°C - 20 sec., 63°C - 20 sec., 75°C - 3 min.
3. 35 cycles of: 95°C - 20 sec., 55°C - 10 sec., 75°C - 3 min.

Product was visualized in a 1.0% native agarose gel as a single band of 0.7 Kb. The fragment was purified and partially sequenced.

5 Multiple alignment of the gene downstream of *D.rad.* identified the following conservative region:

Deinococcus radiodurans GFGG**VQLHAAHGYLL**SQFLSPRHNVRDEYGG
(SEQ. ID. No. 203)

10 *Caenorhabditis elegans* GFDGI**QLHGAHGYLL**SQFTSPTTNKRVDKYGG
(SEQ. ID. No. 204)

Pseudomonas aeruginosa GFSGV**EIHAAHGYLL**SQFLSPLSNRRSDAWGG
(SEQ. ID. No. 205)

15 *Archaeoglobus fulgidus* GFDAV**QLHAAHGYLL**SEFISPHVNRRKDEYGG
(SEQ. ID. No. 206)

The fragment in bold was used to design primers, specifically the downstream primer, for cloning of the 3' region of the *T.th. holB* gene. The upstream 30mer (5'-CATCCTGGACTCGGCCACCTCCTCACCGA-3') (SEQ. ID. No. 207) encodes the amino acid sequence ILDSAHLT (SEQ. ID. No. 208). The downstream 33mer (5'- GAGGAGGTAGCCGTGGGCCGCGTGGAGCTCCAC-3') (SEQ. ID. No. 209) encodes the sequence VELHAAHGYLL (SEQ. ID. No. 210) on the complementary strand. The amplification reactions contained 50ng *T.th.* genomic DNA and 0.1 µM of each primer in a volume of 100µl of Deep Vent polymerase reaction mixture containing 10 µl ThermoPol Buffer, 2.5 mM of each dNTP, 2 mM MgSO₄, and 10 µl DMSO. Amplification was performed using the following cycling scheme:

1. 5 cycles of: 95°C - 30 sec., 70°C - 20 sec., 75°C - 4 min.
2. 5 cycles of: 95°C - 20 sec., 66°C - 20 sec., 75°C - 4 min.
- 30 3. 30 cycles of: 95°C - 20 sec., 60°C - 10 sec., 77°C - 4 min.

Products were visualized in a 1.0% native agarose gel as a single band of 1.1 kb. The Kb fragment was gel purified and sequenced to provide the remainder of the *holB* gene encoding *T.th.* δ'.

For protein expression, the *T.th. holB* gene was cloned into the pET24 vector at the Nde:EcoR sites using a pair of primers. The upstream 32mer (5'-GGCTTTCCCATATGGCTCTACACCCGGCTCAC-3') (SEQ. ID. No. 211) contains a NdeI site within the first 15 nucleotides (underlined) and the sequence
5 corresponding to the 5' region of *T.th. holB*. The downstream 29 mer (5'-GCGTGGATCCACGGTCATGTCTCTAAGTC-3') (SEQ. ID. No. 212) contains a BamHI site within the first 10 nucleotides (underlined) and a sequence complementary to the 3' end of the *holB* gene.

10

EXAMPLE 15

Alternate synthetic path in absence of clamp loader activity

As discussed earlier, the Pol III-type enzyme of the present invention is
15 capable of application and use in a variety of contexts, including a method wherein the clamp loader component that is traditionally involved in the initiation of enzyme activity, is not required. The clamp loader generally functions to increase the efficiency of ring assembly onto circular primed DNA, because both the ring and the DNA are circles and one must be broken transiently for them to become interlocked
20 rings. In such a reaction, the clamp loader increases the efficiency of opening the ring.

The procedure described below illustrates the instance where the clamp loader need not be present. For example, the β clamp can be assembled onto DNA in the absence of the clamp loader. Particularly, the bulk of primed templates in PCR
25 reactions are linear ssDNA fragments that are primed at the ends. On linear primed DNA, the ring need not open at all. Instead, the ring can simply thread onto the end of the linear primed template (Bauer and Burgers, 1988; Tan et al, 1986; O'Day et al., 1992; Burgers and Yoder, 1993). Hence, on linear primed templates, such as those generated in PCR, the beta clamp can simply slide over the DNA end. After the ring
30 slides onto the end, the DNA polymerase can associate with the ring for enhanced DNA synthesis.

Such "end assembly" is common among Pol III-type enzymes and has been demonstrated in yeast and human systems. Rings assembling onto linear DNA for use by their respective DNA polymerases are shown in the following example

09716964-112100

demonstrated in the *E. coli* bacterial system, in the human system, and in the *T.th.* system.

The bulk of the primed templates in PCR reactions are linear ssDNA fragments that are primed at their ends. However, these end primed linear fragments
5 are not generated until after the first step of PCR has already been performed. In the very first step, PCR primers generally anneal at internal sites in a heat denatured ssDNA template. Primed linear templates are then generated in subsequent steps enabling use of this alternate path. For this first step, the clamp may be assembled onto an internal site in the absence of the clamp loader using special conditions that
10 allow clamp assembly in the absence of a clamp loader.

For example, a set of conditions that lead to assembly of the clamp onto circular DNA (i.e., internal primed sites) have been described in the protocol for the use of the bacteriophage T4 ring shaped clamp (gene 45 protein) without the clamp loader (Reddy et al., 1993). In this case, polyethylene glycol leads to
15 "macromolecular crowding" such that the clamp and DNA are pushed together in close proximity, leading to the ring self assembling onto internal primed sites on circular DNA. Other possible conditions that may lead to assembly of rings onto internal sites include use of a high concentration of beta such that use of heat or denaturant to break the dimeric ring into two half rings (crescents) followed by
20 lowering the heat (or dilution or removal of denaturant) leading to rings assembling around the DNA.

The ring shaped sliding clamps of *E. coli* and human slide over the end of linear DNA to activate their respective DNA polymerase in the absence of the clamp loader. This clamp loader independent assay is performed in the bacterial
25 system in Fig. 25A. For this assay, the linear template is polydA primed with oligodT. The polydA is of average length 4500 nucleotides and was purchased from SuperTecs. OligodT35 was synthesized by Oligos etc. The template was prepared using 145µl of 5.2 mM (as nucleotide) polydA and 22 µl of 1.75 mM (as nucleotide) oligodT. The mixture was incubated in a final volume of 2100 µl T.E. buffer (ratio as
30 nucleotide was 21:1 polydA to oligodT). The mixture was heated to boiling in a 1 ml Eppendorf tube, then removed and allowed to cool to room temperature. Assays were performed in a final volume of 25 µl 20 mM Tris-Cl (pH 7.5), 8 mM MgCl₂, 5 mM DTT, 0.5 mM EDTA, 40 mg/ml BSA, 4% glycerol, containing 20 µM [α -³²P]dTTP,

0.1 μ g polydA-oligodT, 25 ng Pol III and, where present, 5 μ g of β subunit. Proteins were added to the reaction on ice, then shifted to 37°C for 5 min. DNA synthesis was quantitated using DE81 paper as described (Rowen and Kornberg, 1978).

In the linear template assay, no ATP or dATP is provided and therefore, a clamp loader, even if present, is not active. Thus, the clamp (e.g., β) can only stimulate the DNA polymerase provided the clamp threads onto the DNA (see diagram in Fig. 25). Hence, threading of the clamp is shown by a stimulation of the DNA polymerase. In lane 1 of Fig. 25A, the DNA polymerase is incubated with the linear DNA in the absence of the clamp, and lane 2 shows the result of adding the clamp. The results show that the clamp is able to thread onto the DNA ends and stimulate the DNA polymerase in the absence of ATP and thus, in the absence of clamp loading as well.

This clamp loader independent assay is performed in the human system in Fig. 25B. The assay reaction (25 μ l) contains 50 mM Tris-HCl (pH=7.8), 8 mM MgCl₂, 1 mM DTT, 1 mM creatine phosphate, 40 μ g/ml bovine serum albumin, 0.55 μ g human SSB, 100 ng PCNA (where present), 7 units DNA polymerase delta (1 unit incorporates 1 pmol dTMP in 60 min.), 40 mM [α -³²P]dTTP and 0.1 μ g polydA-oligodT. Proteins were added to the reaction on ice, then shifted to 37°C for 60 min. DNA synthesis was quantitated using DE81 paper as described (Rowen and Kornberg, 1978). In lane 3, (Fig. 25) the DNA polymerase δ is incubated with the linear DNA in the absence of the clamp, and lane 4 shows the result of adding the PCNA clamp. The results demonstrate that the clamp is able to thread onto the DNA ends and stimulate the DNA polymerase in the absence of ATP and thus, the absence of clamp loading.

This clamp loader independent assay is performed in the *T.th.* system in Fig. 25C. The assay reaction is exactly as described above for use of the *E. coli* Pol III and beta system except the temperature is 60°C and here the Pol III is HEP.P1 *T.th.* Pol III (0.5 μ l, providing 0.1 units where one unit is equal to 1 pmol of dTTP incorporated in 1 minute under these conditions and in the absence of beta), and the beta subunit is 7 μ g *T.th.* β (from the MonoQ column). Proteins were added to the reaction on ice, then shifted to 37°C for 60 min. DNA synthesis was quantitated using DE81 paper as described (Rowen and Kornberg, 1978). In lane 3 (Fig. 25C), the *T.Th.* Pol III is incubated with the linear DNA in the absence of the clamp, and

lane 4 shows the result of adding the *T.th.* β clamp. The results demonstrate that the clamp is able to thread onto the DNA ends and stimulate the DNA polymerase in the absence of clamp loader activity.

5

EXAMPLE 16

Use of *T.th.* Pol III in long chain primer extension

A characteristic of Pol III-type enzymes is their ability to extend a single primer for several kilobases around a long (e.g. 7 kb) circular single stranded DNA genome of a bacteriophage. This reaction uses the circular β clamp protein. For the circular β to be assembled onto a circular DNA genome, the circular β must be opened, positioned around the DNA, and then closed. This assembly of the circular beta around DNA requires the action of the clamp loader, which uses ATP to open and close the ring around DNA. In this example, the 7.2 kb circular single strand DNA genome of bacteriophage M13mp18 was used as a template. This template was primed with a single DNA 57mer oligonucleotide and the Pol III enzyme was tested for conversion of this template to a double strand circular form (RFII). The reaction was supplemented with recombinant *T.th.* β produced in *E. coli*. This assay is summarized in the scheme at the top of Fig. 26. M13mp18 ssDNA was phenol extracted from phage purified as described (Turner and O'Donnell, 1995). M13mp18 ssDNA was primed with a 57mer DNA oligomer synthesized by Oligos etc. The replication assays contained 73 ng singly primed M13mp18 ssDNA and 100 ng *T.th.* β subunit in a 25 μ l reaction containing 20 mM Tris-HCl (pH 7.5), 8 mM MgCl₂, 40 μ g/ml BSA, 0.1 mM EDTA, 4% glycerol, 0.5 mM ATP, 60 μ M each of dCTP, dGTP, dATP and 20 μ M α -³²P-TTP (specific activity 2,000-4,000 cpm/pmol). Either *T.th.* Pol III from the Heparin, peak 1 (HEP.P1; 5 μ l, 0.21 units where 1 unit equals 1 pmol nucleotide incorporated in 1 min.) or a non-Pol III from the Heparin peak 2 (HEP.P2; 5 μ l, 2.6 units) were added to the reaction. Reactions were shifted to 60° C for 5 min., and then DNA synthesis was quenched upon adding 25 μ l of 1% SDS, 40 mM EDTA. One half of the reaction was analyzed in a 0.8% native agarose gel, and the other half was quantitated using DE81 paper as described (Studwell and O'Donnell, 1990).

09716964-112100

The results of the assay are shown in Fig. 26. Lane 1 is the result obtained using the *T.th.* Pol III (HEP.P1) which was capable of extending the primer around the ssDNA circle to form RFII. Lane 2 shows the result of using the non-Pol III (HEP.P2) which was not capable of this extension and produced only incomplete DNA products (the result shown included 0.8 µg *E. coli* SSB which did not increase the chain length of the product). In the absence of SSB, the same product was observed, although the band contained more counts. The greater amount of total synthesis observed in lane 2 is due to the build up of immature products in a small region of the gel. The presence of immature products in lane 1 is likely due to a contaminating polymerase in the preparation that can not convert the single primer to the full length RFII form. Alternatively, the presence of incomplete products in lane 1 (Pol III type enzyme) is due to secondary structure in the DNA which causes the Pol III to pause. In this case it may be presumed that performing the reaction at higher temperature could remove the secondary structure barrier. Alternatively, SSB could be added to the assay (although *T.th.* SSB would be needed, because addition of *E. coli* SSB was tried and did not alter the quality of the product profile). Generally, SSB is needed to remove secondary structure elements from ssDNA at 37°C for complete extension of primers by mesophilic Pol III-type enzymes.

The assay described above was performed at 60°C. The *T.th.* Pol III HEP.P1 gained activity as the temperature was increased from 37°C to 60°C, as expected for an enzyme from a thermophilic source. The *E. coli* Pol III lost activity at 60°C compared to 37°C, as expected for an enzyme from a mesophilic source.

EXAMPLE 17

25

Materials used in Examples 18-29

Radioactive nucleotide were from Dupont NEN; unlabeled nucleotides were from Pharmacia Upjohn. DNA oligonucleotides were synthesized by Gibco BRL. M13mp18 ssDNA was purified from phage that was isolated by two successive bandings in cesium chloride gradients. M13mp18 ssDNA was primed with a 30-mer (map position 6817-6846) as described. The pET protein expression vectors and BL21 (DE3) protein expression strain of *E. coli* were purchased from Novagen. DNA modification enzymes were from New England Biolabs. *Aquifex aeolicus* genomic

DNA was a gift of Dr. Robert Huber and Dr. Karl Stetter (Regensburg University, Germany). Protein concentrations were determined by absorbance at 280nm using extinction coefficients calculated from their known Trp and Tyr content using the equation $\epsilon_{280} = \text{Trp}_m (5690 \text{ M}^{-1} \text{ cm}^{-1}) + \text{Tyr}_n (1280 \text{ M}^{-1} \text{ cm}^{-1})$.

5

EXAMPLE 18

Purification of α Encoded by *dnaE*

The *Aquifex aeolicus dnaE* gene was previously identified (Deckert et al., 1998). The *dnaE* was obtained by searching the *Aquifex aeolicus* genome with the amino acid sequence of *T.th* α subunit (encoded by *dnaE*). The *dnaE* gene was amplified from *Aquifex aeolicus* genomic DNA by PCR using the following primers: the upstream 37mer (5'-GTGTGTCATATGAGTAAG GATTTCGTCCACCTTCACC-3') (SEQ. ID. No. 157) contains an NdeI site (underlined); the downstream 34mer (5'-GTGTGTGGATCCGGGGACTACTCGGAAGTAAGGG-3') (SEQ. ID. No. 158) contains a BamHI site (underlined). The PCR product was digested with NdeI and BamHI, purified, and ligated into the pET24 NdeI and BamHI sites to produce pETAadnaE.

The pETAadnaE plasmid was transformed into the BL21 (DE3) strain of *E. coli*. Cells were grown in 50L of LB containing 100 μ g/ml of kanamycin, 5mM MgSO₄ at 37°C to OD₆₀₀ = 2.0, induced with 2mM IPTG for 20h at 20°C, then collected by centrifugation. Cells were resuspended in 400ml 50mM Tris-HCl (pH 7.5), 10% sucrose, 1M NaCl, 30mM spermidine, 5mM DTT and 2mM EDTA. The following procedures were performed at 4°C. Cells were lysed by passing them twice through a French Press (15,000 psi) followed by centrifugation at 13,000 rpm for 90 min at 4°C. In this protein preparation, as well as each of those that follow, the induced *Aquifex aeolicus* protein was easily discernible as a large band in an SDS polyacrylamide gel stained with Coomassie Blue. Hence, column fractions were assayed for the presence of the *Aquifex aeolicus* protein by SDS PAGE analysis, which forms the basis for pooling column fractions.

The clarified cell lysate was heated to 65°C for 30 min and the precipitate was removed by centrifugation at 13,000 rpm in a GSA rotor for 1h. The

09716964.12100

09716964.112100
supernatant (1.4gm, 280ml) was dialyzed against buffer A (20mM Tris-HCl (pH 7.5)), 10% glycerol, 0.5 mM EDTA, 5mM DTT) overnight, then diluted to 320ml with buffer A to a conductivity equal to 100mM NaCl. The dialysate was applied to a 150ml Fast Flow Q (FFQ) Sepharose column (Pharmacia) equilibrated in buffer A, and eluted with a 1.5L linear gradient of 0-500mM NaCl in buffer A. Eighty fractions were collected. Fractions 38-58 (1g, 390ml) were pooled, dialyzed versus buffer A overnight, and applied to a 250ml Heparin Agarose column (Bio-Rad) equilibrated with buffer A. Protein was eluted with a 1L linear 0-5mM NaCl gradient in buffer A. One hundred fractions were collected. Fractions 69-79 (320 mg in 200 ml) were pooled and dialyzed against buffer A containing 100 mM NaCl. The α preparation was aliquoted and stored frozen at -80°C (see Fig. 27).

EXAMPLE 19

15 Purification of δ Encoded by *holA*

The *Aquifex aeolicus holA* gene was not previously identified by the genome sequencing group at Diversa (Deckert et al., 1998). *Aquifex aeolicus holA* was identified by searching the *Aquifex aeolicus* genome with the amino acid sequence of the *T.th.* δ subunit (encoded by *holA*). The *Aquifex aeolicus holA* was amplified by PCR using the following primers: the upstream 36mer (5'-GTGTGTCATATGGAAACCACAATATTCCAGTTCCAG-3') (SEQ. ID. No. 159) contains an NdeI site (underlined); the downstream 39mer (5'-GTGTGTGGATCCTTATCCACCATGAGAAGTATTTTTCAC-3') (SEQ. ID. No. 160) contains a BamHI site (underlined). The PCR product was digested with NdeI and BamHI, purified, and ligated into the pET24 NdeI and BamHI sites to produce pETAaholA.

The pETAaholA plasmid was transformed into *E. coli* strain BL21 (DE3). Cells were grown in 50L of LB media containing 100 μ g/ml kanamycin. Cells were grown at 37°C to OD₆₀₀ = 2.0, induced for 20h upon addition of 2mM IPTG, then collected by centrifugation. Cells from 25L of culture were lysed as described in Example 18.

The cell lysate was heated to 65°C for 30 min and the precipitate was removed by centrifugation. The supernatant (650mg, 240ml) was dialyzed against

buffer A, adjusted to a conductivity equal to 160mM NaCl by addition of 40ml of
buffer A, and applied to a 220ml Heparin Agarose column equilibrated in buffer A
containing 100mM NaCl. The column was eluted with 1.0L linear gradient of 150-
700 mM NaCl in buffer A. One hundred and four fractions were collected. Fractions
5 45-56 were pooled (250mg, 210 ml), diluted with 230ml buffer A to a conductivity
equal to 230mM NaCl, then loaded onto a 100ml FFQ Sepharose column equilibrated
in buffer A containing 150mM NaCl. The column was eluted with 200ml linear
gradient of 150-750mM NaCl in buffer A; seventy-three fractions were collected.
Fractions 16-38 were pooled (95mg, 40ml), aliquoted, and stored at -80°C (see Fig.
10 27).

EXAMPLE 20

Purification of δ' Encoded by *holB*

15 The *Aquifex aeolicus holB* gene was previously identified by the
genome sequencing facility at Diversa (Deckert et al., 1998). The *Aquifex aeolicus*
holB sequence was obtained by searching the *Aquifex aeolicus* genome with the
sequence of the *T.th.* δ' (encoded by *holB*). The *Aquifex aeolicus holB* gene was
amplified by PCR using the following primers: the upstream 39mer (5'-
20 GTGTGTCATATGGAAAAAGTTTTTTTGGAAA AAAGTCCAG-3') (SEQ. ID.
No. 161) contains an NdeI site (underlined); the downstream 35mer (5'-
GTGTGTGGATCCTTAATCCGCCTGAACGGCTAACG-3') (SEQ. ID. No. 162)
contains a BamHI site (underlined). The PCR product was digested with NdeI and
BamHI, purified, and ligated into the pET24 NdeI and BamHI site to produce
25 pETAaholB.

The pETAaholB plasmid was transformed into *E. coli* strain BL21
(DE3). Cells were grown at 37°C in 50L media containing 100µg/ml kanamycin to
OD₆₀₀ 2.0, then induced for 3h upon addition of 0.2mM IPTG. Cells were collected
by centrifugation and were lysed using lysozyme by the heat lysis procedure (Wickner
30 and Kornberg, 1974). The cell lysate was heated to 65°C for 30 min and precipitate
was removed by centrifugation. The supernatant (2.4g, 400ml) was dialyzed versus
buffer A, then applied to a 220ml FFQ Sepharose column equilibrated in buffer A.
Protein was eluted with a 1L linear gradient of 0-500mM NaCl in buffer A; eighty

fractions were collected. Fractions 23-30 were pooled and diluted 2-fold with buffer A to a conductivity equal to 100mM NaCl, then loaded onto a 200ml Heparin Agarose column equilibrated in buffer A. Protein was eluted with a 1L linear gradient of 0-1.0M NaCl in buffer A; eighty-four fractions were collected. Fractions 46-66
5 were pooled (1.3g, 395ml), dialyzed versus buffer A containing 100mM NaCl, then aliquoted and stored frozen at -80°C (see Fig. 27)

EXAMPLE 21

10 Purification of τ Encoded by *dnaX*

The *Aquifex aeolicus dnaX* gene was previously identified (Deckert et al., 1998). The *dnaX* gene sequence was obtained by searching the *Aquifex aeolicus* genome with the sequence of *T.th.* τ subunit (encoded by *dnaX*). The *Aquifex aeolicus dnaX* was amplified by PCR using the following primers: the upstream
15 41mer (5'-GTGTGTCATATGAACTACGTTCCCTTCGCGAGAAAGTACAG-3') (SEQ. ID. No. 163) contains an NdeI site (underlined); the downstream 36mer (5'-GTGTGTGGATCCTTAAACAGCCTCGTCCCGCTGGA-3') (SEQ. ID. No. 164) contains a BamHI site (underlined). The PCR product was digested with NdeI and BamHI, purified, and ligated into the pET24 NdeI and BamHI sites to produce
20 pETAadnaX.

The pETAadnaX plasmid was transformed into *E. coli* strain BL21 (DE3). Cells were grown in 50L LB containing 100 μ g/ml kanamycin at 37°C to OD₆₀₀ = 0.6, then induced for 20h at 20°C upon addition of IPTG to 0.2mM. Cells were collected by centrifugation and lysed as described in Example 18. The clarified
25 cell lysate was heated to 65°C for 30 min and the protein precipitate was removed by centrifugation. The supernatant (1.1g in 340ml) was treated with 0.228g/ml ammonium sulfate followed by centrifugation. The τ subunit remained in the pellet which was dissolved in buffer B (20mM Hepes (pH 7.5), 0.5mM EDTA, 2mM DTT, 10% glycerol) and dialyzed versus buffer B to a conductivity equal to 87mM NaCl.
30 The dialysate (1073mg, 570ml) was applied to a 200ml FFQ Sepharose column equilibrated in buffer A. The column was eluted with a 1.5L linear gradient of 0-500mM NaCl in buffer A; eighty fractions were collected. Fractions 28-37 were pooled (289mg, 138ml), dialyzed against buffer A to a conductivity equal to 82mM

NaCl, then loaded onto a 150ml column of Heparin Agarose equilibrated in buffer A. The column was eluted with a 900ml linear gradient of 0-500mM NaCl in buffer A; thirty-two fractions were collected. Fractions 15-18 (187mg, 110ml) were dialyzed versus buffer A, then aliquoted and stored at -80°C (see Fig. 27).

5

EXAMPLE 22

Purification of β Encoded by *dnaN*

The *Aquifex aeolicus dnaN* gene was previously identified (Deckert et al., 1998). The *dnaN* sequence was obtained by searching the *Aquifex aeolicus* genome with the sequence of *T.th.* β subunit (encoded by *dnaN*). The *Aquifex aeolicus dnaN* gene was amplified by PCR using the following primers: the upstream 33mer (5'-GTGTGTCATATG CGCGTTAAGGTGGACAGGGAG-3') (SEQ. ID. No. 165) contains an NdeI site (underlined); the downstream 36mer (5'-

15 TGTGTCTCGAG TCATGGCTACACCCTCATCGGCAT-3') (SEQ. ID. No. 166) contains a XhoI site (underlined). The PCR product was digested with NdeI and BamHI, purified, and ligated into the pET24 NdeI and BamHI sites to produce pETAadnaN.

The pETAadnaN plasmid was transformed into *E. coli* strain BL21 (DE3). Cells were grown in 1L LB containing 100mg/ml kanamycin at 37°C to OD₆₀₀ = 1.0, then induced for 6h upon addition of 2mM IPTG. Cells were collected (7g) and lysed as described in Example 18. The cell lysate was heated to 65°C for 30 min and the protein precipitate was removed by centrifugation. The supernatant (39mg, 45ml) was applied to a 10ml DEAE Sephacel column (Pharmacia)

25 equilibrated in buffer A. The column was eluted with a 100ml linear gradient of 0-500mM NaCl in buffer A; seventy-five fractions were collected. Fractions 45-57 were pooled (18.7mg), dialyzed versus buffer A, and applied to a 30ml Heparin Agarose column equilibrated in buffer A. The column was eluted with a 300ml linear gradient of 0-500mM NaCl in buffer A; sixty-five fractions were collected. Fractions 27-33

30 were pooled (11mg, 28ml) and stored at -80°C (see Fig. 27).

09716964.1.2100

EXAMPLE 23

Purification of SSB Encoded by *ssb*

The *Aquifex aeolicus ssb* gene was previously identified (Deckert et al., 1998g). The *ssb* gene sequence was obtained by searching the *Aquifex aeolicus* genome with the sequence of *T.th.* SSB (encoded by *ssb*). The *Aquifex aeolicus ssb* gene was amplified by PCR using the following primers: the upstream 47mer (5'-GTGTGTCATATGCTCAA TAAGGTTTTTATAATAGGAAGACTTACGGG-3') (SEQ. ID. No. 167) contains an NdeI site (underlined); the downstream 39mer (5'-GTGTGGATCCTTA AAAAGGTATTTTCGTCCTCTTCATCGG-3') (SEQ. ID. No. 168) contains a BamHI site (underlined). The PCR product was digested with NdeI and BamHI, purified, and ligated into the pET16 NdeI and BamHI sites to produce pETAassb.

The pETAassb plasmid was transformed into *E. coli* strain BL21 (DE3). Cells were grown in 6L of LB media containing 200µg/ml ampicillin. Cells were grown at 37°C to OD₆₀₀= 0.6, then induced at 15°C overnight in the presence of 2mM IPTG and collected by centrifugation. Cells were lysed as described above in Example 18, except cells were resuspended in buffer C (20mM Tris-HCl (pH 7.9), 500mM NaCl).

The cell lysate was heated to 65°C for 30 min, then the precipitate was removed by centrifugation. The supernatant (1.4g, 190ml) was applied to 25ml Chelating Sepharose column (Pharmacia-Biotech) charged with 50mM Nickel Sulfate and then equilibrated in buffer C containing 5mM Imidazole. The column was eluted with a 300ml linear gradient of 5-100mM Imidazole in buffer C. Fractions of 4ml were collected. Fractions 81-92 were pooled (~240mg in 48ml) and dialyzed overnight against 2L of buffer B containing 200mM NaCl. The dialysate was diluted to a conductivity equal to 92mM NaCl using buffer A and then loaded onto an 8ml MonoQ column equilibrated in buffer A containing 100mM NaCl. The column was eluted with a 120ml linear gradient of 100-500mM Imidazole in buffer A. Seventy-four fractions were collected. Fractions 57-70 were pooled (100mg, 25ml), aliquoted, and stored at -80°C (see Fig. 27).

EXAMPLE 24

MonoQ Preparation of $\tau\delta\delta'$

The δ subunit (0.29mg) purified in Example 19 and δ' subunit
5 (0.31mg) purified in Example 20 were mixed in a volume of 2.8ml of buffer A at
15°C. After 30min, the τ subunit (0.5mg in 1.4ml), purified in Example 21, was
added and the reaction was incubated a further 1h at 15°C. The reaction was applied
to a 1ml MonoQ column equilibrated in buffer A. The $\tau\delta\delta'$ complex elutes later than
either τ , δ or δ' alone. Protein was eluted with a 32ml linear gradient of 100-500mM
10 NaCl in buffer A; eighty fractions were collected. Analysis of the MonoQ fractions in
a SDS polyacrylamide gel shows a peak of $\tau\delta\delta'$ complex that elutes in fractions of
32-38 (see Fig. 28). The peak fractions 850 μ g were stored at -80°C. This procedure
can easily be scaled up. For example, a much larger amount of $\tau\delta\delta'$ was constituted
by following a similar protocol and using a 8ml MonoQ column, which yielded 9.6mg
15 of $\tau\delta\delta'$.

EXAMPLE 25

Constitution of $\alpha\tau\delta\delta'$ Complex

20 The reaction mixture contained 1.2 mg α subunit (9nmol; 133,207 da)
purified in Example 18, 0.41mg τ subunit (7.5 nmol; 54,332 da) purified in
Example 21, 0.41 mg δ subunit (10 nmol; 40,693 da) purified in Example 19, and 0.2
mg δ' subunit (9nmol; 29,000 da) purified in Example 20 in 1.1ml buffer A. The α
and τ subunit solutions were premixed in 871 μ l for 2h at 15°C before adding δ and δ'
25 subunit solution, then the complete mixture was allowed to incubate an additional
12 h at 15°C. The reaction may not require an order of addition, or these extended
incubation times. The reaction mixture was concentrated to 200 μ l using a Centricon
30 at 4°C, then applied to an FPLC Superose 6 HR 10/30 column (25ml) at 4°C
developed with a continuous flow of buffer A containing 100mM NaCl. After the
30 first 216 drops (6.6ml), fractions of 7 drops each were collected. Fractions were
analyzed on a SDS polyacrylamide gel stained with Coomassie Blue (Fig. 29). The
analysis was repeated using the α subunit alone (Fig. 29). The results show that the

peak fractions of α shift to a considerably earlier position when τ , δ and δ' are present and α comigrates with τ , δ , and δ' , when compared to the elution position of α alone, indicating that α assembles with τ , δ and δ' into a $\alpha\tau\delta\delta'$ complex.

5

EXAMPLE 26

$\alpha\tau\delta\delta'$ Functions with the β Clamp

Replication reactions were performed using circular M13mp18 ssDNA primed with a synthetic DNA 90 mer oligonucleotide. Reactions contained 8.6 μ g primed M13mp18 ssDNA, 9.4 μ g SSB purified in Example 23, 1.0 μ g $\alpha\tau\delta\delta'$ prepared in Example 25, and 2.0 μ g β subunit purified in Example 22 (when present), in 230 μ l of 20mM Tris-HCl (pH 7.5), 5mM DTT, 4% glycerol, 8mM MgCl₂, 0.5mM ATP, 60 μ M each dATP and dGTP (buffer composition is for a final volume of 250 μ l). Reactions were mixed on ice, then aliquoted into separate tubes containing 25 μ l each. For each timed reaction, the mixture was brought to 65°C for 2 min before initiating syntheses upon addition of 2 μ l of dCTP and α^{32} P-dTTP (final concentrations, 60 and 40 μ M, respectively). Aliquots were quenched at the times indicated in Fig. 30 upon adding 4 μ l of 0.25M EDTA, 1% SDS. Quenched reactions were then analyzed in a 0.8% alkaline agarose gel. The results, illustrated in Fig. 30, demonstrate that efficient synthesis requires addition of the β subunit. Comparison with size standards in the same gel indicates an average speed of ~125 nucleotides; the leading edge of the product smear indicates a maximum speed of 375 nucleotides/s.

15
20

EXAMPLE 27

25

Purification of *T.th.* α subunit

To obtain *T.th.* α subunit, 8 L of *E. coli* BL21(DE3) cells harboring pETtthalpha were grown to O.D. = 0.3 and induced upon adding IPTG. Cells were collected by centrifugation and resuspended in 200 ml 50mM Tris-HCl (pH 7.5), 10% sucrose, 1M NaCl, 30mM spermidine, 5mM DTT and 2mM EDTA. The following procedures were performed at 4°C. Cells were lysed by passing them three times through a French Press (20,000 psi) followed by incubation at 4°C for 30 min and then centrifugation at 18,000 rpm in an SS-34 rotor for 45 min at 4°C. Induced

30

protein was less than 1% total cell protein but was discernible as a band that migrated in the appropriate position for its predicted molecular weight in an SDS polyacrylamide gel stained with Coomassie Blue. Hence, column fractions were assayed for the presence of the protein by SDS PAGE analysis, which forms the basis for pooling column fractions.

The clarified cell lysate was heated to 65°C for 30 min and the precipitate was removed by centrifugation. The supernatant (1.4gm, 280ml) was dialyzed against buffer A (20mM Tris-HCl (pH 7.5), 10% glycerol, 0.5 mM EDTA, 5mM DTT) overnight, then diluted to 320ml with buffer A to a conductivity equal to 100mM NaCl. The dialysate (approximately 150 mg) was applied to a 60ml DEAE Fast Flow Q (FFQ) Sepharose column (Pharmacia) equilibrated in buffer A, and eluted with a 600 ml linear gradient of 0-500mM NaCl in buffer A. Fractions of 8 ml each were collected. The *Tth. α* subunit could be seen as a major band in several fractions, especially in fractions 26-30. In these peak fractions the *Tth. α* subunit was approximately 20-30 percent pure.

EXAMPLE 28

Purification of *Tth. ε* subunit

The *dnaQ* gene was cloned into the pET16 expression plasmid using the Val within the context "VGLWEW..." and transformed into *E. coli* (BL21(DE3)). This pET plasmid places an N-terminal leader containing six histidines onto the expressed protein to facilitate purification via use of chelate affinity chromatography. Twelve liters of cells were grown to an OD of 0.7 and induced with IPTG. Induced cells were collected by centrifugation and resuspended in 150 ml of buffer C (20mM Tris-HCl (pH 7.9), 500mM NaCl). Cells were lysed by passing them two times through a French Press (20,000 psi) followed by incubation at 4°C for 30 min and then centrifugation at 13,800 rpm in an SLA-1500 rotor for 45 min at 4°C. Induced protein appeared greater than 5% total cell protein and was easily discernible as a band that migrated in the appropriate position for its predicted molecular weight in an SDS polyacrylamide gel stained with Coomassie Blue. Hence, column fractions were assayed for the presence of the protein by SDS PAGE analysis, which forms the basis for pooling column fractions.

Upon analyzing the precipitate from the cell lysis, and the supernatant, it was determined that the epsilon subunit was insoluble and appeared in the precipitate. Therefore the cell pellet was resuspended in 100 ml of binding buffer containing 6M freshly deionized urea. This resuspension was then placed in centrifuge bottles and spun at 13,800 rpm for 45 min in the SLA-1500 rotor. The epsilon was in the supernatant and was applied to a 25 ml Chelating Sepharose column (Pharmacia-Biotech) charged with 50 mM Nickel Sulfate and then equilibrated in buffer C containing 5mM Imidazole. The column was washed with two column volumes of buffer C, then washed with 5 column volumes of beffer C containing 80 mM Imidazole (final). Then the *Tth* epsilon was eluted with a 250 ml linear gradient of 60-1000 mM Imidazole in buffer C. Fractions of 4ml were collected. Fractions 15-24 were pooled (~131 mg) and dialyzed overnight against 2L of buffer A containing 6M urea, but no NaCl or glycerol. The dialysate was then loaded onto an 8ml MonoQ column equilibrated in buffer A containing 6M urea. The column was eluted with a 120ml linear gradient of 0-500 mM NaCl in buffer A containing urea. Sixty five fractions were collected. The epsilon is approximately 80-90 percent pure at this stage. Fractions 13-17 were stored at -80°C. The epsilon is in urea but is at a concentration of 5-10 mg/ml, and thus can be used with other proteins by diluting it such that the final urea concentration is less than 0.5 M. This level of urea does not generally denature protein, and should allow epsilon to renature for catalytic activity.

EXAMPLE 29

25 Temperature optimum of *Aquifex* and *Thermus* α subunit DNA polymerases

The temperature optimum of the alpha subunits of the *Aquifex* and *Thermus* replicases was tested in the calf thymus DNA replication assay. In this experiment, the reactions were assembled on ice in 25 μ l containing 2.5 μ g calf thymus activated DNA, and either 0.88 μ g *Aquifex* α , or 0.6 μ g of the *Thermus* α DEAE pool of peak fractions (obtained from Examples 18 and 28, respectively) in 20 mM Tris-HCl (pH 8.8), 8 mM MgCl₂, 10 mM KCl, 10 mM (NH₄)SO₄, 2 mM MgSO₄, 0.1% Triton X-100, 60 μ M each dATP, dCTP, dGTP, and 20 μ M α^{32} P-dTTP. Reactons were shifted to either 30, 40, 50, 60, 70, 80, or 90°C, then stopped

after 5 minutes and spotted onto DE81 filters to quantitate DNA synthesis. The results, illustrated in Figs. 31-32, show that these enzymes increase in activity as the temperature is raised. The *Thermus* α has a broad peak of activity from 70-80°C (Fig. 31), while the *Aquifex* α is maximal at 80°C (Fig. 32). The *Aquifex* α retains considerable activity at 90°C, whereas the *Thermus* α is nearly inactive at 90°C, a result that is consistent with the higher temperature at which the *Aquifex aeolicus* may live relative to the *Thermus* bacterium.

EXAMPLE 30

10

Temperature optimum of *Aquifex* $\alpha\tau\delta\delta'/\beta$

Aquifex α , β , $\tau\delta\delta'$, SSB and $\alpha\tau\delta\delta'$ were tested for stability at different temperatures by incubating the protein in a solution, followed by performing a replication assay of the protein. Incubation was performed in 0.4 ml tubes under mineral oil. The 5 μ l reaction mixture contained: buffer B (20 mM Tris-HCl (pH 7.5), 5 mM DTT, 5 mM EDTA), and either: 0.352 μ g of α (Fig. 33A), 0.2 μ g of β (Fig. 33B), 0.125 μ g τ complex (Fig. 33C), 0.32 μ g SSB and 0.042 μ g primed M13mp18 ssDNA (Fig. 33D), 0.82 μ g Pol III* (Fig. 33E). Reactions were incubated for 2 min. at either 70, 80, 85, or 90°C in the presence of either 0.1% Triton X-100 (filled diamonds); 0.05% Tween-20 and 0.01% NP-40 (filled circles); 4 mM CaCl₂ (filled triangles); 40% Glycerol (inverted filled triangles); 0.01% Triton X-100, 0.05% Tween-20, 0.01% NP-40, 4 mM CaCl₂ (half-filled square); 40% Glycerol, 0.1% Triton X-100 (open diamonds); 40% Glycerol, 0.05% Tween-20, 0.01% NP-40 (open circles); 40% Glycerol, 4 mM CaCl₂ (open triangles); 40% Glycerol, 0.01% Triton X-100, 0.05% Tween-20, 0.01% NP-40, 4 mM CaCl₂ (half-filled diamonds). After heating, reactions were shifted to ice and 20 μ l of replication assay buffer was added followed by incubation for 1.5 min at 70°C; 15 μ l was then spotted onto a DE81 filter and DNA synthesis was quantitated. The replication assay buffer contained: 60 mM Tris-HCl (pH 9.1 at 25°C), 8mM MgCl₂, 18 mM (NH₄)₂SO₄, 2 mM ATP, 60 μ M each of dATP, dCTP, dGTP, and 20 μ M [α -³²P] TTP (specific activity 10,000 cpm/pmol), and 0.264 μ g primed M13mp18 ssDNA. To assay for β , 0.1 ng $\alpha\tau\delta\delta'$ was added to the reaction. To assay $\tau\delta\delta'$, 0.9 ng β and 0.17 ng α were added to the reaction. To

09716564-112100

assay for SSB, 0.17 ng *E. coli* β and 0.1 ng *E. coli* $\alpha\tau\delta\delta'$ were added to the reaction followed by incubation for 1.5 min at 37°C. To assay for $\alpha\tau\delta\delta'$, 0.9 ng β was added to the reaction. To assay α , the calf thymus DNA replication assay was performed in the buffer as described above but 2.5 μ g activated calf thymus DNA was used instead of primed M13mp18 ssDNA, no other replication proteins were added, and incubation was for 8 min at 70°C.

REFERENCES

The following is a list of documents related to the above disclosure and particularly to the experimental procedures and discussions. The documents should be considered as incorporated by reference in their entirety.

Alonso, J.C., Shirahige, K., and Ogasawara, N. (1990) Molecular cloning, genetic characterization and DNA sequence analysis of the recM region of *Bacillus subtilis*. *Nuc. Acids Res.* **18**:6771-6777.

Altschul et al., (1997) Gapped BLAST and PSI-BLAST: A new generation of protein database search programs. *Nucl. Acids Res.* **25**:3389-3402.

Ausubel, R.M., ed., *Current Protocols in Molecular Biology*, Vol. I-III (1994).

Bambara, R.A., and Jessee, C.B. (1991) Properties of DNA polymerases δ and ϵ , and their role in eukaryotic DNA replication. *Biochimica et Biophysica Acta* **1088**:11-24.

Barnes, M.H., Tarantino, Jr., P.M., Spaccioappoli, P., Brown, N.C., Yu, H., and Dybvig, K. (1994) DNA polymerase III of *Mycoplasma pulmonis*: isolation of characterization of the enzyme and its structural gene, polC. *Molec Microbiol.* **13**:843-854.

Barnes, W.M., (1994) PCR amplification of up to 35-kb DNA with high fidelity and high yield from lambda bacteriophage templates. *Proc. Natl. Acad. Sci. USA* **91**:2216-2220.

Bauer, G.A., and Burgers, P.M. (1988) Protein-protein interactions of yeast DNA polymerase III with mammalian and yeast proliferating cell nuclear antigen (PCNA)/cyclin. *Biochim. Biophys. Acta* **951**(2-3):274-9.

- Blinkowa, A.L., and Walker, J.R., (1990) Programmed ribosomal frameshifting generates the Escherichia coli DNA polymerase III gamma subunit from within the tau subunit reading frame. *Nucl. Acids Res.* **18**(7):1725-1729.
- Bonner, C.A., Stukenberg, P.T., Rajagopalan, M., Eritja, R., O'Donnell, M., McEntee, K., Echols, H., and Goodman, M.F. (1992) Processive DNA synthesis by DNA polymerase II mediated by DNA polymerase III accessory proteins. *J. Biol. Chem.*, **267**:11431-11438.
- Braithwaite, D.K., and Ito, J. (1993) Compilation, alignment, and phylogenetic relationships of DNA polymerases. *Nucl. Acids Res.* **21**(4):787-802.
- 10 Brock, T.D., and Freeze, H., (1969) *Thermus aquaticus* gen. n. and sp. n., a nonsporulating extreme thermophile. *J. Bacteriol.* **98**(1):289-297.
- Burgers, P.M., Yoder, B.L. (1993) ATP-independent loading of the proliferating cell nuclear antigen requires DNA ends. *J. Biol. Chem.* **268**(27):19923-19926.
- Caetano-Anollés et al., (1991) DNA amplification fingerprinting using very short arbitrary oligonucleotide primers. *Bio/Technology* **9**:553-557.
- 15 Carter, J.R., Franden, M.A., Aebersold, R., and McHenry, C.S. (1993) Identification, isolation, and characterization of the structural gene encoding the δ' subunit of *E. coli* DNA polymerase III holoenzyme. *J. Bacteriol.* **175**:3812-3822.
- Celis, J.E., ed., *Cell Biology: A Laboratory Handbook*, Vol. I-III (1994).
- 20 Chen, M., Pan, Z.-Q., and Hurwitz, J. (1992) Studies of the cloned 37-kDa subunit of activator 1 (replication factor C) of HeLa cells. *Proc. Natl. Acad. Sci. USA* **89**(12):5211-5215.
- Cheng, S., Fockler, C., Barnes, W.M., and Higuchi, R., (1994) Effective amplification of long targets from cloned inserts and human genomic DNA. *Proc. Natl. Acad. Sci. USA* **91**:5695-5699.
- 25 Coligan, J.E., ed., *Current Protocols in Immunology*, Vol. I-III (1994).
- Cullman, G., Fien, K., Kobayashi, R., and Stillman, B. (1995) Characterization of the five replication factor C genes of *Saccharomyces cerevesiae*. *Mol. and Cell. Biol.* **15**:4661-4671.

- Davis, L.G., *Basic Methods In Molecular Biology*, Elsevier Edit., New York (1986).
- Decket et al., (1998) The complete genome of the hyperthermophilic bacterium *Aquifex aeolicus*. *Nature* **392**:353-358.
- Dulbecco, R., et al. (1959) Plaque production by the polyoma virus. *Virology* **8**:396-397.
- 5 Edge, M.D., et al., (1981) Total synthesis of a human leukocyte interferon gene. *Nature* **292**:756.
- Flower, A.M. and McHenry, C.S. (1990) The γ subunit of DNA polymerase III holoenzyme of *Escherichia coli* is produced by ribosomal frameshifting. *Proc. Natl. Acad. Sci. USA* **87**:3713-3717.
- 10 Freshney, R.I., ed., *Animal Cell Culture* (1986).
- Gait, M.J., ed., *Oligonucleotide Synthesis* (1984).
- Glover, ed., *DNA Cloning: A Practical Approach*, Vol. I & II, MRL Press, Ltd., Oxford, U.K. (1985).
- Guenther, B.D. (1996) Structural studies on the DNA replication apparatus: X-ray crystal structure of the δ' subunit of *Escherichia coli* DNA Pol III. Ph.D. Thesis, Rockefeller University.
- 15 Guibus, J.M., Kelman, Z., Hurwitz, J., O'Donnell, M., and Kuriyan, J. (1996) Structure of the C-terminal region of p21waf1/cip1 complexed with human PCNA. *Cell* **87**:297-306.
- 20 Hames, B.D., and Higgins, S.J., eds., *Nucleic Acid Hybridization* (1985).
- Hames, B.D., and Higgins, S.J., eds., *Transcription and Translation* (1984).
- Hammerling et al., *Monoclonal Antibodies and T-cell Hybridomas* (1981).
- Harlow and Lane, eds., *Antibodies - A Laboratory Manual*, Cold Spring Harbor, New York (1988).
- 25 Heath, D.D., Iwama, G.K., and Devlin, R.H., (1993) PCR primed with VNTR core sequences yields species specific patterns and hypervariable probes. *Nucl. Acids Res.* **21**(24):5782-5785.

- Hughes, Jr., A.J., Bryan, S.K., Chen, H., Moses, R.E., and McHenry, C.S. (1991) Escherichia coli DNA polymerase II is stimulated by DNA polymerase III holoenzyme auxiliary subunits. *J. Biol. Chem.* **266**:4568-4573.
- IRL Press, Publ., *Immobilized Cells and Enzymes* (1986).
- 5 Ito, J., and Braithwaite, D., (1991) Compilation and alignment of DNA polymerase sequences. *Nucl. Acids Res.* **19**(15):4045-4057 (1991).
- Jacks, T., Madhami, H.D., Masiarz, F.R., and Varmus, H.E. (1988) Signals for ribosomal frameshifting in the Rous sarcoma virus *gag-pol* region. *Cell* **55**:447-458.
- Jay, E., et al., (1984) Chemical synthesis of a biologically active gene for human
10 immune interferon-gamma. Prospect for site-specific mutagenesis and structure-function studies. *J. Biol. Chem.* **259**:6311-6317.
- Kelman Z., and O'Donnell, M. (1995) DNA Polymerase III holoenzyme: Structure and function of a chromosomal replicating machine, *Annu. Rev. Biochem.*, **64**:171-200.
- 15 Kelman, Z., and O'Donnell, M. (1994) DNA replication: enzymology and mechanisms. *Current Opinions in Genetics and Development* **4**:185-195.
- Kennett et al., *Monoclonal Antibodies* (1980).
- Kong, X. -P., Onrust, R., O'Donnell, M., and Kuriyan, J. (1992). Three dimensional structure of the β subunit of *Escherichia coli* DNA polymerase III holoenzyme: a
20 sliding DNA clamp. *Cell* **69**:425-437.
- Kornberg, A., and Baker, T. (1992). *DNA Replication*, second edition. (New York:W. H. Freeman and Company), pp. 165-194.
- Krishna, T.S., Kong, X.-P., Gary, S. , Burgers, P.M., and Kuriyan, J. (1994) Crystal structure of the eukaryotic DNA polymerase processivity factor PCNA. *Cell* **79**(7):
25 1233-1243.
- Kuriyan, J. and O'Donnell, M. (1993) Sliding clamps of DNA polymerases. *J. Mol. Biol.* **234**:915-925.

- Larsen, B., Wills, N.M., Gesteland, R.F., and Atkins, J.F. (1994) rRNA-mRNA base pairing stimulates a programmed -1 ribosomal frameshift. *J. Bact.* **176**: 6842-6851.
- Lin, J.J., and Kuo, J. (1995) *Focus* **17**(2):66-70.
- Linn, S. (1991) How many pols does it take to replicate nuclear DNA? *Cell* **66**:185-187.
- Lee, S.H. and Walker, J.R. (1987) Escherichia coli dnaX product, the τ subunit of DNA polymerase III, is a multifunctional protein with single-stranded DNA-dependent ATPase activity. *Proc. Natl. Acad. Sci. USA* **84**:2713-2717.
- Low, R. L., Rashbaum, S.A., and Cozzarelli, N.R. (1976) Purification and characterization of DNA polymerase III from Bacillus subtilis. *J. Biol. Chem.*, **251**:1311-1325.
- Maki, H., Maki, S., and Kornberg, A. (1988) DNA polymerase III holoenzyme of *Escherichia coli* IV. The holoenzyme is an asymmetric dimer with twin active sites. *J. Biol. Chem.* **263**:6570-6578.
- Maniatis, T., Fritsch, E.F., and Sambrook, J. (1992) *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
- McHenry, C.S. (1991) DNA Polymerase III Holoenzyme. *J. Biol. Chem.*, **266**:19127-19130.
- McHenry, C.S., and Crow, W. (1979) DNA polymerase III of *Escherichia coli*: Purification and identification of Subunits. *J. Biol. Chem.* **254**(5):1748-1753.
- McHenry, C.S. (1982) Purification and characterization of DNA polymerase III'. Identification of τ as a subunit of the DNA polymerase III holoenzyme. *J. Biol. Chem.* **257**:2657-2663.
- Moarefi, I., Jeruzalmi, D., Turner, J., O'Donnell, M., and Kuriyan, J. (2000) Crystal structure of the DNA polymerase processivity factor of T4 bacteriophage. *J. Molec. Biol.*, **296**:1215-1223.
- Naktinis, V., Turner, J., and O'Donnell, M. (1996) A molecular switch in a replication machine defined by an internal competition for protein rings. *Cell* **84**(1):137-145.

09716964-112100

- Naktinis, V., Onrust, R., Fang, L., O'Donnell, M. (1995) Assembly of a chromosomal replication machine: two DNA polymerases, a clamp loader, and sliding clamps in one holoenzyme particle. II. Intermediate complex between the clamp loader and its clamp. *J. Biol. Chem.* **270**:13358-13365.
- 5 Nambair, K.P., et al., (1984) Total synthesis and cloning of a gene coding for the ribonuclease S protein. *Science* **223**:1299-1300.
- Nelson, K.E., et al., (1999) Evidence for lateral gene transfer between *Archaea* and bacteria from genome sequence of *Thermotoga maritima*. *Nature* **399**:323-329.
- Noren, C.J., et al., (1989) A general method for site-specific incorporation of
10 unnatural amino acids into proteins. *Science* **244**:182-188.
- O'Day, C.L., Burgers, P.M., Taylor, J.S. (1992) PCNA-induced DNA synthesis past *cis*-syn and *trans*-syn-I thymine dimers by calf thymus DNA polymerase delta *in vitro*. *Nucl. Acids Res.* **20**(20):5403-6.
- O'Donnell, M., (1987) Accessory proteins bind a primed template and Mediate Rapid
15 Cycling of DNA polymerase III Holoenzyme from *E. coli*. *J. Biol. Chem.* **262**:16558-16565.
- O'Donnell, M., Kuriyan, J., Kong, X-P., Stukenberg, P.T. and Onrust, R. (1992) The sliding clamp of DNA polymerase III holoenzyme encircles DNA. *Molec. Biol. Cell* **3**:953-957.
- 20 O'Donnell, M., Onrust, R., Dean, F.B., Chen, M., and Hurwitz, J. (1993) Homology in accessory proteins of replicative polymerases-*E. coli* to humans. *Nucl. Acids Res.* **21**:1-3.
- Onrust, R., Finkelstein, J., Turner, J., Naktinis, V., and O'Donnell, M. (1995)
Assembly of a chromosomal replication machine: two DNA polymerases, a clamp
25 loader and sliding clamps in one holoenzyme particle. III) Interface between two polymerases and the clamp loader. *J. Biol. Chem.* **270**:13366-13377.
- Onrust, R. and O'Donnell, M. (1993) DNA polymerase III accessory proteins. I) holoA and holoB encoding δ and δ' . *J. Biol. Chem.* **268**:11758-11765.

09716964-12100

- Onrust, R., Stukenberg, P.T., and O'Donnell, M. (1991) Analysis of the ATPase subassembly which initiates processive DNA synthesis by DNA polymerase III holoenzyme. *J. Biol. Chem.* **266**:21681-21686.
- Oshima, T., and Imahori, K. (1974) Description of *Thermus thermophilus* (Yoshida and Oshima) comb-nov, a nonsporulating bacterium from a Japanese spa. *Int. J. Syst. Bacteriol.* 24(1):102-112.
- Pacitti, D.F., Barnes, M.H., Li, D.H., and Brown, N.C. (1995) Characterization and overexpression of the gene encoding *Staphylococcus aureus* DNA polymerase III. *Gene*, **1165**:51-56.
- 10 Perbal, B., *A Practical Guide to Molecular Cloning* (1984).
- Perrino, F.W., and Loeb, L.A. (1990) Hydrolysis of 3'-terminal mispairs *in vitro* by the 3' 5' exonuclease of DNA polymerase δ permits subsequent extension by DNA polymerase α . *Biochem.* **29**:5226-5231.
- Reddy et al., (1993) Assembly of a functional replication complex without ATP hydrolysis: a direct interaction of bacteriophage T4 gp45 with T4 DNA polymerase. *Proc. Natl. Acad. Sci. USA* **90**(8):3211-3215.
- 15 Rowen, L., and Kornberg, A. (1978) Primase, the DnaG protein of *Escherichia coli*. An enzyme which starts DNA chains. *J. Biol. Chem.* **253**:758-764.
- Ruttimann, C., Cotoras, M., Zaldivar, J., and Vicuna, R. (1985) DNA polymerases from the extremely thermophilic bacterium *Thermus thermophilus* HB-8. *European J. Biochem.* **149**:41-46.
- 20 Sambrook et al., *Molecular Cloning: A Laboratory Manual* (1989).
- Sanger, F., and Coulson, A.R., (1975) A rapid method for determining sequences in DNA by primed synthesis with DNA polymerase. *J. Mol. Biol.* **94**:441-448.
- 25 Sanger, F., et al., (1977) DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA* 74:5463-5467
- Schreier, M., et al., *Hybridoma Techniques* (1980).

09716964-112100

- Studwell-Vaughan, P.S. and O'Donnell, M. (1991) Constitution of the twin polymerase of DNA polymerase III holoenzyme. *J. Biol. Chem.* **266**:19833-19841.
- Studwell-Vaughan, P.S. and O'Donnell, M. (1990) Processive replication is contingent on the exonuclease subunit of DNA polymerase III holoenzyme. *J. Biol. Chem.* **265**(2):1171-1178.
- Stukenberg, P.T., Studwell-Vaughan, P.S., and O'Donnell, M. (1991) Mechanism of the sliding β -clamp of DNA polymerase III holoenzyme. *J. Biol. Chem.* **266**:11328-11334.
- Sugino, A. (1995) Yeast DNA polymerases and their role at the replication fork. Elsevier Science Ltd., 319-323.
- Tabor, S. and Richardson, C.C. (1995) A single residue in DNA polymerases of the *Escherichia coli* DNA polymerase I family is critical for distinguishing between deoxy- and dideoxyribonucleotides. *Proc. Natl. Acad. Sci. USA*, **92**(14):6339-6343.
- Tan, C.K., Castillo, C., So, A.G., Downey, K.M. (1986) An auxiliary protein for DNA polymerase-delta from fetal calf thymus. *J. Biol. Chem.* **261**(26):12310-6.
- Towbin, H., Staehelin, T., Gordon, J. (1979) Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proc. Natl. Acad. Sci. USA* **76**(9):4350-4354.
- Tsuchihashi, Z., and Kornberg, A. (1989) ATP interactions of the tau and gamma subunits of DNA polymerase III holoenzyme of *Escherichia coli*. *J. Biol. Chem.* **264**:17790-95.
- Tsuchihashi, Z. and Kornberg, A. (1990) Translational frameshifting generates the γ subunit of DNA polymerase III holoenzyme. *Proc. Natl. Acad. Sci. USA* **87**:2516-2520.
- Tsuchihashi, Z., and Brown, P.O. (1992) Sequence requirements for efficient translational frameshifting in the *Escherichia coli* dnaX gene and the role of an unstable interaction between tRNA^{Lys} and an AAG lysine codon. *Genes and Dev.* **6**:511-519.

00716964-12100

- Turner, J., and O'Donnell, M. (1995) Cycling of Escherichia coli DNA polymerase III from one sliding clamp to another: model for lagging strand. *Methods Enzymol.* **262**:442-449.
- Vos, P., et al., (1995) AFLP: a new technique for DNA fingerprinting. *Nucl. Acids Res.* **23**(21):4407-4414.
- Weiss, R.B., Dunn, D.M., Atkins, J.F., and Gesteland, R.F., (1987) Slippery runs, shifty stops, backward steps, and forward hops: -2, -1, +2, +5 and +6 ribosomal frameshifting. in Cold Spring Harbor Symposia on Quantitative Biology **52**: 687-693.
- Welsh, J., and McClelland, M., (1990) Fingerprinting genomes using PCR with arbitrary primers. *Nucl. Acids Res.* **18**(24):7213-7218.
- Wickner, W., and Kornberg, A., (1974) A holoenzyme form of DNA Polymerase III. Isolation and Properties. *J. Biol. Chem.* **249**(19):6244-6249.
- Williams, J.G., et al., (1990) DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucl. Acids Res.* **18**(22):6531-6535.
- Yin, K-C., Blinkowa, A., and Walker, J.R. (1986) Nucleotide sequence of the Escherichia replication gene *dnaZX*. *Nuc. Acids. Res.* **14**:6541-6549.
- Young, M.C., Reddy, M.K., and von Hippel, P.H. (1992) Structure and function of the bacteriophage T4 DNA polymerase holoenzyme. *Biochem.*, **31**:8675-8690.
- Yuzhakov, A., Turner, J. and O'Donnell, M. (1996) Replisome assembly reveals the basis for asymmetric function in leading and lagging strand replication. *Cell* **86**:877-886.
- U.S. Patent No. 5,668,004 to O'Donnell.
- U.S. Patent No. 5,583,026 to O'Donnell.
- U.S. Patent No. 5,545,552 to Mathur.
- U.S. Patent No. 5,498,523 to Tabor et al.
- U.S. Patent No. 5,455,166 to Walker.
- U.S. Patent No. 5,409,818 to Davey et al.
- U.S. Patent No. 5,374,553 to Gelfand et al.
- U.S. Patent No. 5,352,778 to Comb et al.

EP 534,858 to Vos et al.

This invention may be embodied in other forms or carried out in other ways without departing from the spirit or essential characteristics thereof. The present disclosure is
25 therefore to be considered as in all respects illustrative and not restrictive, the scope of
the invention being indicated by the appended claims, and all changes which come
within the meaning and range of equivalency are intended to be embraced therein.

WHAT IS CLAIMED:

1. An isolated DNA molecule from a thermophilic bacterium, the isolated DNA molecule encoding a DNA polymerase III-type enzyme subunit.

5

2. The isolated DNA molecule according to claim 1, wherein the enzyme subunit is selected from the group consisting of alpha, beta, tau, gamma, epsilon, delta, delta prime, and SSB subunits.

10

3. The isolated DNA molecule according to claim 2, wherein the enzyme subunit is a delta subunit.

4. The isolated DNA molecule according to claim 3, wherein the thermophilic bacterium is *Aquifex aeolicus*.

15

5. The isolated DNA molecule according to claim 4, wherein the delta subunit comprises an amino acid sequence of SEQ. ID. No. 124.

20

6. The isolated DNA molecule according to claim 4, wherein the DNA molecule comprises a nucleotide sequence of SEQ. ID. No. 123 or hybridizes to a DNA molecule comprising the nucleotide sequence of SEQ. ID. No. 123 under stringent conditions.

25

7. The isolated DNA molecule according to claim 3, wherein the thermophilic bacterium is *Thermus thermophilus*.

8. The isolated DNA molecule according to claim 7, wherein the delta subunit comprises an amino acid sequence of SEQ. ID. No. 158.

30

9. The isolated DNA molecule according to claim 7, wherein the DNA molecule comprises a nucleotide sequence of SEQ. ID. No. 157 or hybridizes to a DNA molecule comprising the nucleotide sequence of SEQ. ID. No. 157 under stringent conditions.

10. The isolated DNA molecule according to claim 3, wherein the thermophilic bacterium is *Thermatoga maritima*.

11. The isolated DNA molecule according to claim 10, wherein the delta subunit comprises an amino acid sequence of SEQ. ID. No. 146.

12. The isolated DNA molecule according to claim 10, wherein the DNA molecule comprises a nucleotide sequence of SEQ. ID. No. 145 or hybridizes to a DNA molecule comprising the nucleotide sequence of SEQ. ID. No. 145 under stringent conditions.

13. The isolated Dna molecule according to claim 3, wherein the thermophilic bacterium is *Bacillus stearothermophilus*.

14. The isolated DNA molecule according to claim 13, wherein the delta subunit comprises an amino acid sequence of SEQ. ID. No. 178.

15. The isolated DNA molecule according to claim 13, wherein the DNA molecule comprises a nucleotide sequence of SEQ. ID. No. 177 or hybridizes to a DNA molecule comprising the nucleotide sequence of SEQ. ID. No. 177 under stringent conditions.

16. The isolated DNA molecule according to claim 2, wherein the replication enzyme subunit is a delta prime subunit.

17. The isolated DNA molecule according to claim 16, wherein the thermophilic bacterium is *Aquifex aeolicus*.

18. The isolated DNA molecule according to claim 17, wherein the delta prime subunit comprises an amino acid sequence of SEQ. ID. No. 126.

19. The isolated DNA molecule according to claim 17, wherein the DNA molecule comprises a nucleotide sequence of SEQ. ID. No. 125 or hybridizes to

a DNA molecule comprising the nucleotide sequence of SEQ. ID. No. 125 under stringent conditions.

5 20. The isolated DNA molecule according to claim 16, wherein the thermophilic bacterium is *Thermus thermophilus*.

 21. The isolated DNA molecule according to claim 20, wherein the delta prime subunit comprises an amino acid sequence of SEQ. ID. No. 156.

10 22. The isolated DNA molecule according to claim 20, wherein the DNA molecule comprises a nucleotide sequence of SEQ. ID. No. 155 or hybridizes to a DNA molecule comprising the nucleotide sequence of SEQ. ID. No. 155 under stringent conditions.

15 23. The isolated DNA molecule according to claim 16, wherein the thermophilic bacterium is *Thermatoga maritima*.

 24. The isolated DNA molecule according to claim 23, wherein the delta prime subunit comprises an amino acid sequence of SEQ. ID. No. 146.

20 25. The isolated DNA molecule according to claim 23, wherein the DNA molecule comprises a nucleotide sequence of SEQ. ID. No. 147 or hybridizes to a DNA molecule comprising the nucleotide sequence of SEQ. ID. No. 147 under stringent conditions.

25 26. The isolated DNA molecule according to claim 16, wherein the thermophilic bacterium is *Bacillus stearothermophilus*.

30 27. The isolated DNA molecule according to claim 26, wherein the delta prime subunit comprises an amino acid sequence of SEQ. ID. No. 180.

 28. The isolated DNA molecule according to claim 26, wherein the DNA molecule comprises a nucleotide sequence of SEQ. ID. No. 179 or hybridizes to

0074969460

a DNA molecule comprising the nucleotide sequence of SEQ. ID. No. 179 under stringent conditions.

5 29. An isolated replication enzyme subunit of a thermophilic bacterium which is encoded by the isolated DNA molecule of claim 1.

 30. The isolated replication enzyme subunit according to claim 29, wherein the replication enzyme subunit is selected from the group of consisting alpha, beta, tau, gamma, epsilon, delta, delta prime, and SSB subunits.

10 31. The isolated replication enzyme subunit according to claim 30, wherein the replication enzyme subunit is a delta subunit.

 32. The isolated replication enzyme subunit according to claim 31, wherein the thermophilic bacterium is *Aquifex aeolicus*.

 33. The isolated replication enzyme subunit according to claim 32, wherein the delta subunit comprises an amino acid sequence of SEQ. ID. No. 124.

20 34. The isolated replication enzyme subunit according to claim 31, wherein the thermophilic bacterium is *Thermus thermophilus*.

 35. The isolated replication enzyme subunit according to claim 34, wherein the delta subunit comprises an amino acid sequence of SEQ. ID. No. 158.

25 36. The isolated replication enzyme subunit according to claim 31, wherein the thermophilic bacterium is *Thermotoga maritima*.

 37. The isolated replication enzyme subunit according to claim 36, wherein the delta subunit comprises an amino acid sequence of SEQ. ID. No. 146.

30 38. The isolated replication enzyme subunit according to claim 31, wherein the thermophilic bacterium is *Bacillus stearothermophilus*.

39. The isolated replication enzyme subunits according to claim 38,
wherein the delta subunit comprises an amino acid sequence of SEQ. ID. No. 178.

5 40. The isolated replication enzyme subunit according to claim 30,
wherein the replication enzyme subunit is a delta prime subunit.

41. The isolated replication enzyme subunit according to claim 40,
wherein the thermophilic bacterium is *Aquifex aeolicus*.

10 42. The isolated replication enzyme subunit according to claim 41,
wherein the delta prime subunit comprises an amino acid sequence of SEQ. ID. No.
126.

15 43. The isolated replication enzyme subunit according to claim 40,
wherein the thermophilic bacterium is *Thermus thermophilus*.

20 44. The isolated replication enzyme subunit according to claim 43,
wherein the delta prime subunit comprises an amino acid sequence of SEQ. ID. No.
156.

45. The isolated replication enzyme subunit according to claim 40,
wherein the thermophilic bacterium is *Thermotoga maritima*.

25 46. The isolated replication enzyme subunit according to claim 45,
wherein the delta prime subunit comprises an amino acid sequence of SEQ. ID. No.
148.

30 47. The isolated replication enzyme subunit according to claim 40,
wherein the thermophilic bacterium is *Bacillus stearothermophilus*.

48. The isolated replication enzyme subunit according to claim 47,
wherein the delta prime subunit comprises an amino acid sequence of SEQ. ID. No.
180.

09716964.12100

49. An expression system comprising an expression vector into which is inserted a heterologous DNA molecule according to claim 1.

50. The expression system according to claim 40, wherein the heterologous DNA molecule is in sense orientation and correct reading frame.

51. A host cell comprising a heterologous DNA molecule according to claim 1.

52. A method of producing a recombinant thermostable DNA polymerase III-type enzyme, or subunit thereof, from a thermophilic bacterium, said method comprising:

transforming a host cell with at least one heterologous DNA molecule according to claim 1 under conditions suitable for expression of the DNA polymerase III-type enzyme, or subunit thereof, and

isolating the DNA polymerase III-type enzyme, or subunit thereof.

53. The method according to claim 52, wherein the enzyme subunit is selected from the group consisting of alpha, beta, tau, gamma, epsilon, delta, delta prime, and SSB subunits.

54. The method according to claim 53, wherein the enzyme subunit is a delta or delta prime subunit.

55. The method according to claim 54, wherein the thermophilic bacteria is *Thermus thermophilus*, *Aquifex aeolicus*, *Thermotoga maritima*, or *Bacillus stearothermophilus*.

56. The method according to claim 52, wherein said transforming is carried out by transforming the host cell with a plurality of heterologous DNA molecules according to claim 1 under conditions suitable for expression of the DNA polymerase III-type enzyme, or a plurality of subunits thereof, and said isolating is carried out by isolating the DNA polymerase III-type enzyme, or the plurality of subunits thereof.

57. An isolated clamp loader of a DNA polymerase III-type enzyme comprising either a heterologously expressed delta subunit, a heterologously expressed delta prime subunit, or both, derived from a thermophilic eubacteria.

5

58. The isolated clamp loader according to claim 57, wherein the thermophilic bacteria is a *Thermus* species, a *Thermotoga* species, an *Aquifex* species, or a *Bacillus* species.

10

59. The isolated clamp loader according to claim 58, wherein the thermophilic bacteria is *Thermus thermophilus*.

60. The isolated clamp loader according to claim 59, wherein the delta subunit comprises an amino acid sequence of SEQ. ID. No. 158.

15

61. The isolated clamp loader according to claim 59, wherein the delta prime subunit comprises an amino acid sequence of SEQ. ID. No. 156.

20

62. The isolated clamp loader according to claim 58, wherein the thermophilic bacteria is *Thermotoga maritima*.

63. The isolated clamp loader according to claim 62, wherein the delta subunit comprises an amino acid sequence of SEQ. ID. No. 146.

25

64. The isolated clamp loader according to claim 62, wherein the delta prime subunit comprises an amino acid sequence of SEQ. ID. No. 148.

30

65. The isolated clamp loader according to claim 58, wherein the thermophilic bacteria is *Aquifex aeolicus*.

66. The isolated clamp loader according to claim 65, wherein the delta subunit comprises an amino acid sequence of SEQ. ID. No. 124.

0974594-12100

67. The isolated clamp loader according to claim 65, wherein the delta prime subunit comprises an amino acid sequence of SEQ. ID. No. 126.

5 68. The isolated clamp loader according to claim 58, wherein the thermophilic bacteria is *Bacillus stearothermophilus*.

69. The isolated clamp loader according to claim 68, wherein the delta subunit comprises an amino acid sequence of SEQ. ID. No. 178.

10 70. The isolated clamp loader according to claim 68, wherein the delta prime subunit comprises an amino acid sequence of SEQ. ID. No. 180.

0071694.13100
0071694.13100


ABSTRACT OF THE INVENTION

The present invention relates to an isolated DNA molecule from a thermophilic bacterium which encodes a DNA polymerase III-type enzyme subunit.

5 Also encompassed by the present invention are host cells and expression system including the heterologous DNA molecule of the present invention, as well as isolated replication enzyme subunits encoded by such DNA molecules. Also disclosed is a method of producing a recombinant thermostable DNA polymerase III-type enzyme, or subunit thereof, from a thermophilic bacterium, which is carried out by
10 transforming a host cell with at least one heterologous DNA molecule of the present invention under conditions suitable for expression of the DNA polymerase III-type enzyme, or subunit thereof, and then isolating the DNA polymerase III-type enzyme, or subunit thereof.

001694.1300

FIG.1



ATP binding

E. coli
 MSYQVLARKWRPQTFADVVGQEHVLTALANGLSLGRIHHAYLFSGTRGVGKTSIARLLAK
 B. subtilis
 MSYQALYRVFRPQRFEDVVGQEHITKTLQNALLQKKFSHAYLFSGPRGTGKTSAAKIFAK
 **** * * *** * ***** . * * * . ***** ** ***** **

E. coli
 GLNCETGITATPCGVCDNCREIEQGRFVDLIEIDAASRTKVEDTRDLLDNVQYAPARGRF
 B. subtilis
 AVNCEHAPVDEPCNECAACKGITNGSISDVIEIDAASNNGVDEIRDIDKVKFAPSAVTY
 *** ** * * * * * . * * * * * . * * * * * . * * * * * .

E. coli
 KVYLIDEVHMLSRHSFNALLKTLLEPPPEHVKFLATTDPQKLPVTILSRCLQFHLKALDV
 B. subtilis
 KVYIIDEVHMLSIGAFNALLKTLLEPPPEHCIFILATTEPHKIPLTIISRCQRFDFKRITS
 *** . ***** . ***** . ***** . ***** . ***** . ***** .

FIG. 2




FIG.3

TCCGGGGGTG	GGGTTCCAG	GTAGACCCCG	GCCCCTCGG	TGAGCCCCTT	TACCCAGGCC	60
GCCACCTCCT	CCAGGGGGGC	CAAGCGGTGC	AAGGAGAGGA	ACGTCCGCAC	<u>CAGGCCCTAT</u>	120
ACTAGCCCTT	GTG AGC GCC CTC TAC CGC CGC TTC CGC CCC CTC ACC TTC CAG GAG GTG GTG				S.D.	180
	met ser ala leu tyr arg phe arg pro leu thr phe gln glu val val					(17)
GGG CAG GAG CAC GTG AAG GAG CCC CTC CTC AAG GCC ATC CGG GAG GGG AGG CTC GCC CAG					CAC	240
gly gln glu his val lys glu pro leu leu lys ala ile arg glu gly arg leu ala gln						(37)
GCS TAC CTS TTC TCC GGS AC						
GCC TAC CTC TTC TCC GGG CCC AGG GGC GTG GGC AAG ACC ACC GCG AGG CTC CTC GCC						300
ala tyr leu phe ser gly pro arg gly val gly lys thr thr ala arg leu leu ala						(57)
ATG GCG GTG GGG TGC CAG GGG GAA GAC CCC CCT TGC GGG GTC TGC CCC CAC TGC CAG GCG						360
met ala val gly cys gln gly glu asp pro pro cys gly val cys pro his cys gln ala						(77)
GtG CAG AGG GGC GCC CAC CCG GAC GTG GTG GAC ATT GAC GCC AGC AAC AAC TCC GTG						420
val gln arg gly ala his pro asp val val asp ile asp ala ala ser asn ser val						(97)
GAG GAC GTG CGG GAG CTG AGG GAA AGG ATC CAC CTC GCC CCC CTC TCT GCC CCC AGG AAG						480
glu asp val arg glu leu arg glu arg ile his leu ala pro leu ser ala pro arg lys						(117)
GTC TTC ATC CTG GAC GAG GCC CAC ATG CTC TCC AAA AGC GCC TTC AAC GCC CTC CTC AAG					^C	540
val phe ile leu asp Glu ala his met leu ser lys ser ala phe asn ala leu leu lys						(137)

FIG.4A-1

CCC CTTT CCC CCG CCC CAG GCC CTG ATC GCC ATG ACC GCC CTG GAC GAG GCC ATG 1080
pro leu pro ala pro pro gln ala leu ile ala ala met thr ala leu asp glu ala met (317

GAG CGC CTC GCC CGC CGC TCC GAC GCC TTA AGC CTG GAG GTG GCC CTC CTG GAG GCG GGA	1140
glu arg leu ala arg arg ser asp ala leu ser leu glu val ala leu glu ala gly	(337)
AGG GCC CTG GCC GAG GCC CTA CCC CAG CCC ACG GGC GCT CCT TCC CCA GAG GTC GGC	1200
arg ala leu ala ala glu ala leu pro gln pro thr gly ala pro ser pro glu val gly	(357)
CCC AAG CCG GAA AGC CCC CCG ACC CCG GAA CCC CCA AGG CCC GAG GAG GCG CCC GAC CTG	1260
pro lys pro glu ser pro pro thr pro pro glu pro arg pro glu ala pro asp leu	(377)
CGG GAG CGG TGG CGG GCC TTC CTC GAG GCC CTC AGG CCC ACC CTA CCG GCC TTC GTG CGG	1320
arg glu arg trp arg ala phe leu glu ala leu arg pro thr leu arg ala phe val arg	(397)
GAG GCC CGC CGC GAG GTC CCG GAA GGC CAG CTC TGC CTC GCT TTC CCC GAG GAC AAG GCC	1380
glu ala arg pro glu val arg glu gly gln leu cys leu ala phe pro glu asp lys ala	(417)
TTC CAC TAC CGC AAG GCC TCG GAA CAG AAG GTG AGG CTC CTC CCC CTG GCC CAG GCC CAT	1440
phe his tyr arg lys ala ser glu gln lys val arg leu leu pro leu ala gln ala his	(437)
frameshift site	
TTC GGG GTG GAG GAG GTC GTC CTC GTC GAG GGA GAA AAA AAA AGC CTG AGC CCA AGG	1500
phe gly val glu glu val val leu val leu glu gly glu lys lys ser leu ser pro arg	(457)

FIG.4B-1

CCC CGC CCG GCC CCA CCT CCT GAA GCG CCC GCA CCC CCG GGC CCT CCC GAG GAG GAG GTA	1560
pro arg pro ala pro pro pro pro glu ala pro ala pro pro gly pro pro glu glu val	(477)
GAG GCG GAG GAA GCG GCG GAG GAG GCG CCG GAG GAG GAG GAG GTC CGC CTC	1620
glu ala glu glu ala ala glu glu ala pro glu ala leu arg arg val val arg leu	(497)
CTG GGG GGG CCG GTG CTC TGG GTG CCG CCG AGG ACC CCG GAG GCG CCG GAG GAG GAA	1680
leu gly gly arg val leu trp val arg arg pro arg thr arg glu ala pro glu glu glu	(517)
CCC CTG AGC CAA GAC GAG ATA GGG GGT ACT GGT ATA TAA	1740
pro leu ser gln asp glu ile gly thr gly ile *	(529)
CGACCTCGGA CAAGAGACCG TGGACAACAT CCTCAAGCGC	1820
GGTGCGGGG CTCCAGAAGA TGGTGGCCGA GGGCCGCCCC	1880
GATGACCGCC ACCAAGAAGG CCATGGAGGC GCGGGCCACC	1940
GAACGTCTGC GCCGCCGAGG TCTCCGAGGG CAAGGTGAAC	2000
CGCCACCATG CTGAAGAACT TCATCTA	2027

FIG.4B-2

GGG	CAG	GAG	CAC	GTG	AGC	GCC	CTC	TAC	CGC	CGC	TTC	CGC	CCC	CTC	ACC	TTC	CAG	GAG	GTG	GTG	51
GCC	TAC	CTC	TTC	TCC	GTG	AAG	GAG	CCC	CTC	GCC	AAG	GCC	ATC	CGG	GAG	GGG	AGG	CTC	GCC	CAG	111
ATG	GCG	GTG	GGG	TGC	CAG	GGG	CCC	GAA	GAC	GGG	CCT	TGC	GGG	GTC	ACC	TGC	CAC	TGC	CTC	GCC	171
GtG	CAG	AGG	GGC	GCC	CAC	CCG	CCG	GAC	GTG	GAC	ATT	GAC	GCC	GCC	GCC	GCC	AAC	TGC	CAG	GCG	231
GAG	GAC	GTG	CGG	GAG	CTG	AGG	AGG	GAA	AGG	ATC	CAC	CTC	GCC	CCC	CTC	TCT	GCC	CCC	AGG	AAG	291
GTC	TTC	ATC	CTG	GAC	GAG	GCC	GCC	CAC	ATG	CTC	TCC	AAA	AGC	GCC	TTC	AAC	GCC	CTC	CTC	AAG	351
ACC	CTG	GAG	GAG	CCC	CCC	CCC	CCC	CAC	GTC	CTC	TTC	GTC	TTC	GCC	ACC	ACC	GAG	CCC	GAG	AGG	411
ATG	CCC	CCC	ACC	ATC	CTC	TCC	CGC	ACC	CGC	ATC	CAG	TTC	CGC	TTC	CGC	CGC	CTC	ACG	GAG	AGG	471
GAG	ATC	GCC	TTT	AAG	CTC	CGG	CGC	ATC	CGC	ATC	CTG	GAG	GCC	GTG	CGG	GAG	GCG	GAG	GAG	GAG	531
GCC	CTC	CTC	CTC	CTC	GCC	CGC	CGC	CTG	GCG	CTG	GAC	GGG	GCC	CTT	AGG	GAC	GCG	AGC	CTC	CTG	591
GAG	CGC	TTC	CTC	CTC	CTC	CTG	GAA	GGC	CCC	CGC	ATC	CGG	ACC	CTT	AGG	GAG	GCG	GAA	AGC	CTC	651
TCC	CCC	CCA	GGG	ACC	GGG	GTG	GTG	GCC	GAG	ATC	GCC	GCC	TCC	CTC	CTC	GTG	GAG	CGC	GCC	CTA	711
GAG	GCC	CTG	GGC	CTC	GCC	CGG	CGC	CTC	CGC	CTC	TAC	GGG	GAA	GGG	TAC	GCC	AGG	GGG	AAA	ACG	771
TCG	GGC	CTT	TTG	GAG	GTG	GTG	TTC	CGG	GAA	GGC	CTC	TAC	GCC	GGC	GGG	TTC	GGC	CTC	AGC	CTG	831
CCC	CTT	CCC	GCC	CGC	CGC	CGC	CGC	CGC	CTG	GCC	ATC	GCC	GCC	ATG	ACC	GCC	CTG	GAC	GAG	GCC	891
GAG	CGC	CTC	GCC	CGC	CGC	TCC	TCC	GAC	GCC	CTG	TTA	AGC	CTG	GAG	GTG	GCC	CTC	GAG	GCG	GGA	951
AGG	GCC	CTG	GCC	GCC	GAG	GCC	GCC	CTA	CCC	CTA	CCC	ACG	GGC	GGC	GCT	CCT	TCC	CCA	GAG	GCG	1011
CCC	AAG	CCG	GAA	AGC	CCC	CCG	CCG	ACC	CCG	ACC	CAA	CCC	AGG	CCC	AGG	GAG	GAG	CCC	CCC	CTG	1071
CGG	GAG	CGG	TGG	CGG	GCC	TTC	CTC	CTC	GAG	GCC	CTC	AGG	CCC	AGG	CCC	GAG	GAG	GCG	GAC	CTG	1131
GAG	GCC	CGC	CCG	GAG	GTC	CGG	CGG	GAA	GGC	GAG	CTC	TGC	CTC	CTC	ACC	CTA	CGG	GCC	TTC	GTG	1191
TTC	CAC	TAC	CGC	AAG	GCC	TCG	TCG	GAA	CAG	AGG	AAG	GTG	AGG	CCC	ACC	TTC	CCC	GAG	AAG	GCC	1251
TTC	GGG	GTG	GAG	GAG	GTC	GTC	GTC	CTC	GTG	AGG	GAG	GGA	GAA	AAA	AGC	CTG	GCC	CAG	GCC	CAT	1311
CCC	CGC	CCG	GCC	CCA	CCT	CCT	CCT	GAA	GCG	CCC	GCA	CCC	CCG	GGC	AGC	CTG	AGC	AGC	CCA	AGG	1371
GAG	GCG	GAG	GAA	GCG	GCG	GAG	GAG	GAG	GCC	CCG	CCG	CCC	GCC	TTG	AGG	CGG	GAG	GTG	GTC	CGC	1431
CTG	GGG	GGG	CGG	GTG	CTC	TGG	GTG	CGG	CGG	ACC	CGG	AGG	ACC	CGG	GAG	GCG	GAG	CCG	GAG	GAA	1491
																					1551

CCC CTG AGC CAA GAC GAG ATA GGG GGT ACT GGT ATA TAA (1590)

FIG.4C

Met	ser	ala	leu	tyr	arg	arg	phe	lys	ala	ile	pro	leu	thr	phe	gln	glu	val	val	gln	glu	gln	glu	20
his	val	lys	glu	pro	arg	leu	lys	ala	arg	thr	ile	arg	glu	gly	arg	leu	ala	ala	ala	tyr	leu	40	
phe	ser	gly	pro	arg	glu	val	gly	lys	thr	thr	thr	thr	thr	ala	arg	leu	ala	ala	met	ala	val	60	
gly	cys	gln	gly	glu	asp	pro	pro	cys	gly	val	val	cys	pro	pro	his	cys	gln	ala	val	gln	arg	80	
gly	ala	his	pro	asp	val	val	asp	ile	asp	ala	asp	ala	ala	ser	asn	asn	ser	val	glu	asp	val	100	
arg	glu	leu	arg	glu	arg	ile	his	leu	leu	ala	ser	pro	leu	ser	ala	pro	arg	lys	val	phe	ile	120	
leu	asp	glu	ala	his	met	leu	ser	lys	val	phe	arg	ala	phe	thr	thr	leu	lys	lys	thr	leu	glu	140	
glu	pro	pro	pro	his	val	leu	phe	leu	gln	leu	thr	gln	thr	arg	glu	pro	glu	arg	met	pro	pro	160	
thr	ile	leu	ser	arg	thr	ile	his	phe	ala	val	arg	phe	arg	arg	leu	thr	glu	glu	glu	ile	ala	180	
phe	lys	leu	arg	arg	leu	leu	glu	ala	leu	val	gly	arg	arg	glu	ala	glu	glu	glu	ala	leu	leu	200	
leu	leu	ala	arg	leu	ala	pro	asp	gly	leu	leu	lys	arg	asp	ala	glu	ser	leu	leu	glu	arg	phe	220	
gly	thr	gly	val	ala	glu	pro	thr	arg	ile	glu	lys	glu	val	glu	arg	ala	leu	gly	ser	pro	pro	240	
gly	leu	ala	arg	arg	leu	leu	ala	ala	ala	ser	lys	leu	ala	arg	gly	lys	thr	ala	glu	ala	leu	260	
leu	glu	val	phe	arg	glu	leu	gly	leu	tyr	gly	ala	ala	ala	phe	arg	ser	leu	val	ser	gly	leu	280	
ala	pro	pro	gln	ala	leu	ile	ile	ala	ala	met	thr	ala	ala	leu	leu	ala	gly	thr	pro	leu	pro	300	
ala	arg	arg	ser	asp	ala	leu	ile	leu	leu	thr	gly	val	ala	leu	leu	glu	ala	met	glu	arg	leu	320	
ala	ala	glu	ala	leu	pro	pro	gln	pro	thr	gly	ala	pro	pro	glu	leu	glu	ala	gly	arg	ala	leu	340	
glu	ser	pro	pro	thr	pro	pro	glu	pro	pro	thr	gly	ala	pro	glu	ala	pro	val	gly	pro	lys	pro	360	
trp	arg	ala	phe	leu	glu	glu	ala	leu	arg	pro	arg	pro	thr	leu	arg	phe	val	arg	leu	glu	arg	380	
pro	glu	val	arg	glu	gly	gln	gln	leu	cys	leu	leu	ala	phe	pro	glu	asp	lys	ala	phe	his	tyr	400	
arg	lys	ala	ser	glu	gln	lys	val	arg	arg	leu	leu	leu	pro	leu	ala	gln	ala	his	phe	gly	val	420	
glu	glu	val	val	leu	val	leu	glu	glu	glu	glu	glu	lys	pro	lys	ser	leu	pro	arg	pro	arg	pro	440	
ala	pro	pro	pro	glu	ala	pro	ala	pro	pro	pro	pro	gly	pro	pro	pro	glu	glu	val	glu	ala	glu	460	
glu	ala	ala	glu	glu	ala	pro	pro	glu	glu	ala	ala	leu	arg	arg	val	val	arg	leu	glu	ala	glu	480	
arg	val	leu	trp	val	arg	arg	pro	arg	thr	thr	thr	arg	arg	glu	ala	pro	glu	leu	leu	gly	gly	500	
gln	asp	glu	ile	gly	thr	thr	thr	thr	thr	thr	thr	thr	thr	thr	thr	thr	thr	thr	thr	thr	thr	520	
																						529	

FIG.4D

Met	ser	ala	lys	glu	pro	arg	tyr	arg	arg	pro	leu	thr	phe	gln	glu	val	gly	gln	glu	20	
his	val	lys	glu	pro	arg	leu	lys	ala	arg	ile	arg	glu	gly	arg	leu	ala	ala	tyr	leu	40	
phe	ser	gly	pro	arg	glu	val	gly	lys	thr	thr	thr	thr	ala	arg	leu	ala	met	ala	val	60	
gly	cys	gln	gly	glu	asp	pro	pro	cys	gly	val	cys	pro	his	arg	cys	gln	ala	val	gln	arg	80
gly	ala	his	pro	arg	val	val	asp	ile	asp	ala	pro	ala	ser	asn	asn	ser	val	glu	asp	val	100
arg	glu	leu	arg	glu	arg	ile	his	leu	ala	pro	leu	ser	ala	pro	arg	lys	val	phe	ile	120	
leu	asp	glu	ala	his	met	leu	ser	lys	ser	ala	phe	thr	asn	ala	leu	lys	thr	leu	glu	140	
glu	pro	pro	pro	his	val	leu	phe	val	phe	ala	thr	thr	thr	glu	pro	glu	arg	met	pro	160	
thr	ile	leu	ser	arg	thr	gln	his	phe	arg	phe	arg	arg	arg	leu	thr	glu	glu	ile	ala	180	
phe	lys	leu	arg	arg	ile	leu	glu	ala	val	gly	gly	glu	glu	ala	glu	glu	ala	leu	leu	200	
leu	leu	ala	arg	leu	ala	asp	gly	ala	leu	arg	asp	ala	ala	glu	ser	leu	leu	glu	phe	220	
leu	thr	leu	glu	gly	pro	leu	thr	arg	lys	glu	val	glu	glu	arg	ala	leu	gly	ser	pro	240	
gly	thr	gly	val	ala	arg	leu	ala	ala	ser	leu	ala	arg	arg	gly	lys	thr	ala	glu	ala	260	
gly	leu	ala	arg	arg	arg	leu	gly	glu	tyr	tyr	ala	pro	arg	leu	ser	leu	val	ser	gly	280	
leu	glu	val	phe	gln	phe	gly	leu	ile	ala	ala	phe	gly	leu	ala	ala	gly	thr	pro	leu	300	
ala	pro	pro	pro	ala	ala	leu	ile	ala	met	thr	ala	leu	asp	glu	glu	ala	met	glu	arg	320	
ala	arg	arg	ser	asp	ala	ala	leu	ser	leu	val	ala	leu	leu	leu	glu	ala	gly	arg	ala	340	
ala	ala	glu	ala	leu	pro	gln	pro	pro	thr	gly	ala	pro	ser	pro	glu	val	gly	pro	lys	360	
glu	ser	pro	pro	thr	pro	glu	pro	pro	arg	pro	glu	glu	ala	pro	asp	leu	arg	glu	arg	380	
trp	arg	ala	phe	leu	glu	ala	leu	leu	arg	pro	thr	leu	arg	phe	val	arg	glu	ala	arg	400	
pro	glu	val	arg	glu	gly	gln	leu	leu	cys	leu	ala	phe	pro	glu	asp	lys	ala	phe	his	420	
arg	lys	ala	ser	glu	gln	lys	val	arg	leu	leu	pro	leu	ala	ala	gln	ala	his	phe	gly	440	
glu	glu	val	val	leu	val	leu	glu	gly	glu	lys	lys	lys	pro	pro	asp	pro	lys	ala	pro	460	
gly	pro	thr	ser																	464	

FIG.4E

Met	ser	ala	leu	tyr	arg	arg	phe	arg	pro	leu	thr	phe	gln	glu	val	val	gly	gln	glu	20
his	val	lys	glu	pro	leu	arg	lys	ala	ile	arg	glu	gly	arg	leu	ala	gln	ala	tyr	leu	40
phe	ser	gly	pro	arg	gly	val	gly	lys	thr	thr	thr	ala	arg	leu	ala	ala	met	ala	val	60
gly	cys	gln	gly	glu	asp	pro	pro	cys	gly	val	cys	pro	his	cys	gln	ala	val	gln	arg	80
gly	ala	his	pro	asp	val	val	asp	ile	asp	ala	ala	ser	asn	asn	ser	val	glu	asp	val	100
arg	glu	leu	arg	glu	arg	ile	his	leu	ala	pro	leu	ser	ala	pro	arg	lys	val	phe	ile	120
leu	asp	glu	ala	his	met	leu	ser	lys	ser	ala	phe	asn	ala	leu	leu	lys	thr	leu	glu	140
glu	pro	pro	pro	his	val	leu	phe	val	phe	ala	thr	thr	glu	pro	glu	arg	met	pro	pro	160
thr	ile	leu	ser	arg	thr	gln	his	phe	arg	phe	arg	arg	leu	thr	glu	glu	glu	ile	ala	180
phe	lys	leu	arg	arg	ile	leu	glu	ala	val	gly	arg	glu	ala	glu	glu	glu	ala	leu	leu	200
leu	leu	ala	arg	leu	ala	asp	gly	ala	leu	arg	asp	ala	glu	ser	leu	leu	glu	arg	phe	220
leu	leu	leu	glu	gly	pro	leu	thr	arg	lys	glu	val	glu	arg	ala	leu	gly	ser	pro	pro	240
gly	thr	gly	val	ala	glu	leu	ile	ala	ala	ser	leu	ala	arg	lys	thr	ala	glu	ala	leu	260
gly	leu	ala	arg	arg	leu	tyr	gly	glu	gly	tyr	ala	pro	arg	ser	leu	val	ser	gly	leu	280
leu	glu	val	phe	arg	glu	glu	leu	tyr	ala	ala	phe	gly	leu	ala	gly	thr	pro	leu	pro	300
ala	pro	pro	gln	ala	leu	ile	ala	ala	met	thr	ala	leu	asp	glu	ala	met	glu	arg	leu	320
ala	arg	arg	ser	asp	ala	leu	ser	leu	glu	val	ala	leu	leu	glu	ala	gly	arg	ala	leu	340
ala	ala	glu	ala	leu	pro	gln	pro	thr	gly	ala	pro	ser	pro	glu	val	gly	pro	lys	pro	360
glu	ser	pro	pro	thr	pro	glu	pro	pro	arg	pro	glu	glu	ala	pro	asp	leu	arg	glu	arg	380
trp	arg	ala	phe	leu	glu	ala	leu	leu	arg	pro	thr	leu	arg	phe	val	arg	glu	ala	arg	400
pro	glu	val	arg	glu	gly	gln	leu	leu	cys	leu	phe	pro	glu	asp	lys	ala	phe	his	tyr	420
arg	lys	ala	ser	glu	gln	lys	val	arg	leu	leu	pro	leu	ala	gln	ala	his	phe	gly	val	440
glu	glu	val	val	leu	val	leu	glu	gly	glu	lys	lys	lys	lys	ala						454

FIG.4F

		ATP site	
E.coli	MSYQVLARKWRPQTFADVVGQEHVLTALANGLSLGRHHAYLFSGTRGVGKTSIARLLAK	60	
H.inf.K.....II.....KDN.L.....F..	60	
B.sub.	...A.Y.VF...R.E.....ITKT.Q.A.LQKKFS.....P.T...A.KIF..	60	
C.cres.	DA.T....Y..R..E.LI...AMVRT...AF.T...A..FMLT.V.....TT.....R	113	
M.gen.	-MH..FYQ.Y..IN.KQTL...SIRKI.V.AINRDKLPNG.I...E.T...TF.KII..	59	
T.th.	--VSA.Y.RF..L..QE.....KEP.LKAIRE..LAQ.....P.....TT.....M	58	
	Zn ⁺⁺ finger		
	* * *		
E.coli	GLNCET----GITATPCGVCDNCREIEQGRFVDLIEIDAASRTKVEDTRDLLDNVQYAPA	116	
H.inf.	...VH-----V.....E.E..KA...N.I.....E.....K.V	116	
B.sub.	AV...H----APVDE..NE.AA.KG.TN.SIS.V.....NNG.DEI..IR.K.KF..S	116	
C.cres.	A..Y..DTVK.PSVDLTTEGYH..S.IE..HM.VL.L.....DEM.E...G.R...V	173	
M.gen.	AI..LN----WDQIDV.NS..V.KS.NTNSAI.IV.....KNGIN.I.E.VE..FNH.F	115	
T.th.	AVG.QG-----EDP.....PH.QAVQR.AHP.VVD.....NNS...V.E.RERIHL...L	112	
E.coli	RGRFKVYLIDEVHMLSRHSFNALLKTLLEPPPEHVKFLLATDPQKLPVTILSRCLQFHLK	176	
H.inf.	V.....Y.....	176	
B.sub.	AVTY...I.....IGA.....CI.I...E.H.I.L..I...QR.DF.	176	
C.cres.	EA.Y...I.....TAA.....P.A..IF...EIR.V.....QR.D.R	233	
M.gen.	TFKK...IL..A...TTQ.WGG.....S.PY.L.IFT..EFN.I.L.....QS.FF.	175	
T.th.	SAPR..FIL..A...KSA.....P..L.VF...E.ERM.P.....TQH.RFR	172	

FIG.5A

E.coli	ALDVEQIRHQLEHILNEEHIAHEPRALQQLARAAEGSLRDALSITDQAIASGDQ--VST	234
H.inf.	...ET...SQH.A...TQ.N.PF.DP..VK..K..Q..I..S.....M..R...TN	234
B.sub.	RITSQA.VGRMNK.VDA.QLQV.EGS.EII.S..H.GM.....L....SFSGDI--LKV	234
C.cres.	RVEPDVLVKHFDR.SAK.GARI.MD..A.I.....V..G...L....VQTERGQT.TS	293
M.gen.	KITSDL.LER.ND.AKK.K.KI.KD..IKI.DLSQ.....G...L..LAI.LIVKKL.LL	235
T.th.	R.TE.E.AFK.RR..EAVGREA.EE..L...L.D.A...E..LERFLLLEGP---LTR	229
E.coli	QAVSAMLGTLDDDDQALSVEAMVEANGERVMALINEAAARGIEWEALLVEMGLLHRIAM	294
H.inf.	NV..N...L...NYSVDILY.LHQG...LL.RTLQRV.DAAGD.DK..G.CAEK...Q..L	294
B.sub.	EDALLIT.AVSQLYIGK.AKSLHDK.VSDALETL..LLQQ.KDPK.IED.IFYFRDMLL	294
C.cres.	TV.RD...LA.RS.TIA.Y.HVMAGKTKDALEGFRALWGF.ADPVVMLDV.DHC.AS.V	353
M.gen.	MLKKHLISLIEMQN.L.KQFYQ.I	260
T.th.	KE.ERA...SPPGTGVAEIAASLARGKTAELG.ARRLYGE.YAPRS.VSGL.EVFREGLY	289

FIG.5B




FIG.6





FIG.7

FIG.8A



	Reading frame	Blue	White
Shifty sequence	0	+	
	- 1	+	
	- 2	+	
Mutant sequence	0	++	
	- 1		+
	- 2		+

FIG.8B




FIG.9




FIG.10A FIG.10B FIG.10C




FIG. 11A




FIG. 11B

FIG.12A




FIG.12B




FIG.12C




FIG. 13A

FIG.13C





FIG.14A


load FT 9 10 11 12 13 14 15 16 17 18 19 ^{E. coli}
 _{α γ δ}



↑ ↑
T.th E. coli
subunits subunits

FIG.14B

load FT 9 10 11 12 13 14 15 16 17 18 19



← α

Alignment of TTH1 with alphas subunits of other organisms.

E.coli	DRYFLELIRTGRPDEESYLHAAVELAEARGLPVV	197	(ID#72)
V.chol.	DHFYLELIRTGRADEESYLHFALDVAEQYDLPVV	197	(ID#73)
H.inf.	DHFYALSRTPNEERYIQAAKLAERCDDLPLV	197	(ID#74)
R.prow.	DRFYFEIMRHDLP EEGFIENSYIQIASELSIPIV	195	(ID#75)
H.pyl.	DDFYLEIMRHGILDQRFIDEQVIKMSLETGLKII	213	(ID#76)
S.sp.	DDYYLEIQDHGSVEDRLVNINLVKIAQELDIKIV	202	(ID#77)
M.tub.	DNYFLELMDHGLTIERRVRDGLLEIGRALNIPPL	220	(ID#78)
T.th.	FFIEIQNHGLSEQK		(ID#61)

FIG.15A

Alignment of TTH2 with alphas subunits of other organisms.

E.coli	NKRRAKNGEPPLDIAAIPLDDKKSFDM LQRSETTAVFQLESRGMKD	618	(ID#79)
V.chol.	NPRLKKAGKPPVRIEAIPLDDARSFRNLQDAKTTAVFQLESRGMKKE	618	(ID#80)
H.inf.	NVRMVREGKPRVDIAAIPLDDPESFELLKRSETTAVFQLESRGMKD	618	(ID#81)
R.prow.	CKKLLKEQGIKIDFDDMTFDDDKKTYQMLCKGKGVGFQFESIGMKD	624	(ID#82)
H.pyl.	LKIIKTQHKISVDFLSLMDMDPKVYKTIQSGDTVGFQIES-GMFQ	648	(ID#83)
S.sp.	QERKALQIRARTGSKKLPDDVKKTHKLL EAGDLEGIFQLESQGMKQ	643	(ID#84)
M.tub.	IDNVRANRGIDLDESVP LDDKATYELLGRGDTLGVFQLDGGPMRD	646	(ID#85)
T.th.	RVELDYDALTLDD		(ID#60)

FIG.15B

ATGGGCCGGGAGCTCCGCTTCGCCCACCTCCACCAGCACA	
CCCAGTTCTCCCTCCTGGACGGGGCGGCGAAGCTTTCCGA	
CCTCCTCAAGTGGGTCAAGGAGACGACCCCCGAGGACCCC	120
GCCTTGGCCATGACCGACCACGGCAACCTCTTCGGGGCCG	
TGGAGTTCTACAAGAAGGCCACCGAAATGGGCATCAAGCC	
CATCCTGGGCTACGAGGCCTACGTGGCGGCGGAAAGCCGC	240
TTTGACCGCAAGCGGGGAAAGGGCCTAGACGGGGGCTACT	
TTACCTCACCTCCTCGCCAAGGACTTCACGGGGTACCA	
GAACCTGGTGCGCTGGCGAGCCGGGCTTACCTGGAGGGG	360
TTTTACGAAAAGCCCCGGATTGACCGGGAGATCCTGCGCG	
AGCACGCCGAGGGCCTCATCGCCCTCTCGGGGTGCCTCGG	
GGCGGAGATCCCCAGTTTCATCCTCCAGGACCGTCTGGAC	480
CTGGCCGAGGCCCGGCTCAACGAGTACCTCTCCATCTTCA	
AGGACCGCTTCTTCATCGAGATCCAGAACCACGGCCTCCC	
CGAGCAGAAAAAGGTCAACGAGGTCTCAAGGAGTTCGCC	600
CGAAAGTACGGCCTGGGGATGGTGGCCACCAACGACGGCC	
ATTACGTGAGGAAGGAGGACGCCCCGCGCCACGAGGTCCT	
CCTCGCCATCCAGTCCAAGAGCACCTTGGACGACCCCGGG	720
CGCTGGCGCTTCCCCTGCGACGAGTTCTACGTGAAGACCC	
CCGAGGAGATGCGGGCCATGTTCCCCGAGGAGGAGTGGGG	
GGACGAGCCCTTTGACAACACCGTGGAGATCGCCCGCATG	840
TGCAACGTGGAGCTGCCCATCGGGGACAAGATGGTCTACC	
GAATCCCCCGCTTCCCCCTCCCCGAGGGGCGGACCGAGGC	
CCAGTACCTCATGGAGCTCACCTTCAAGGGGCTCCTCCGC	960
CGCTACCCGGACCGGATCACCGAGGGCTTCTACCGGGAGG	
TCTTCCGCCTTTTGGGGAAGCTTCCCCCCCCACGGGGACGG	
GGAGGCCTTGGCCGAGGCCTTGGCCCAGGTGGAGCGGGAG	1080
GCTTGGGAGAGGCTCATGAAGAGCCTCCCCCTTTGGCCG	
GGGTCAAGGAGTGGACGGCGGAGGCCATTTTCCACCGGGC	
CCTTTACGAGCTTTCCGTGATAGAGCGCATGGGGTTTCCC	1200
GGCTACTTCTCATCGTCCAGGACTACATCAACTGGGCCC	
GGAGAAACGGCGTCTCCGTGGGGCCCCGGCAGGGGGAGCGC	
CGCCGGGAGCCTGGTGGCCTACGCCGTGGGGATCACCAAC	1320
ATTGACCCCCCTCCGCTTCGGCCTCCTCTTTGAGCGCTTCC	
TGAACCCGGAGAGGGTCTCCATGCCCGACATTGACACGGA	
CTTCTCCGACCGGGAGCGGGACCGGTGATCCAGTACGTG	1440
CGGGAGCGCTACGGCGAGGACAAGGTGGCCCAGATCGGCA	
CCCTGGGAAGCCTCGCCTCCAAGGCCGCCCTCAAGGACGT	
GGCCCGGGTCTACGGCATCCCCACAAGAAGGCGGAGGAA	1560
TTGGCCAAGCTCATCCCGGTGCAGTTCGGGAAGCCCAAGC	
CCCTGCAGGAGGCCATCCAGGTGGTGCCGGAGCTTAGGGC	
GGAGATGGAGAAGGACCCCAAGGTGCGGGAGGTCTTCGAG	1680
GTGGCCATGCGCCTGGAGGGCCTGAACCGCCACGCCTCCG	
TCCACGCCGCCGGGGTGGTGATCGCCGCCGAGCCCCCTCAC	
GGACCTCGTCCCCCTCATGCGCGACCAGGAAGGGCGGCCC	1800
GTCACCCAGTACGACATGGGGGCGGTGGAGGCCTTGGGGC	
TTTTGAAGATGGACTTTTTTGGGCCTCCGCACCCTCACCTT	

FIG. 16A

CCTGGACGAGGTCAAGCGCATCGTCAAGGCGTCCCAGGGG	1920
GTGGAGCTGGACTACGATGCCCTCCCCCTGGACGACCCCA	
AGACCTTCGCCCTCCTCTCCCGGGGGGAGACCAAGGGGGT	
CTTCCAGCTGGAGTCGGGGGGGATGACCGCCACGCTCCGC	2040
GGCCTCAAGCCGCGGCGCTTTGAGGACCTGATCGCCATCC	
TCTCCCTCTACCGCCCCGGGCCCATGGAGCACATCCCCAC	
CTACATCCGCCGCCACCACGGGCTGGAGCCCGTGAGCTAC	2160
AGCGAGTTTCCCCACGCCGAGAAGTACCTAAAGCCCATCC	
TGGACGAGACCTACGGCATCCCCGTCTACCAGGAGCAGAT	
CATGCAGATCGCCTCGGCCGTGGCGGGGTACTCCCTGGGC	2280
GAGGCGGACCTCCTGCGGCGGTCCATGGGCAAGAAGAAGG	
TGGAGGAGATGAAGTCCCACCGGGAGCGCTTCGTCCAGGG	
GGCCAAGGAAAGGGGCGTGCCCCGAGGAGGAGGCCAACC GC	2400
CTCTTTGACATGCTGGAGGCCTTCGCCAACTACGGCTTCA	
ACAAATCCCACGCTGCCGCCTACAGCCTCCTCTCCTACCA	
GACCGCCTACGTGAAGGCCCACTACCCCGTGGAGTTCATG	2520
GCCGCCCTCCTCTCCGTGGAGCGGCACGACTCCGACAAGG	
TGGCCGAGTACATCCGCGACGCCCGGGCCATGGGCATAGA	
GGTCCTTCCCCCGGACGTCAACCGCTCCGGGTTTGACTTC	2640
CTGGTCCAGGGCCGGCAGATCCTTTTCGGCCTCTCCGCGG	
TGAAGAACGTGGGCGAGGCGGCGGCGGAGGCCATTCTCCG	
GGAGCGGGAGCGGGGCGGCCCTACCGGAGCCTCGGCGAC	2760
TTCTCAAGCGGCTGGACGAGAAGGTGCTCAACAAGCGGA	
CCCTGGAGTCCCTCATCAAGGCGGGCGCCCTGGACGGCTT	
CGGGGAAAGGGCGCGGCTCCTCGCCTCCCTGGAAGGGCTC	2880
CTCAAGTGGGCGGCCGAGAACC GGGAAGGCCCGCTCGG	
GCATGATGGGCCTCTTCAGCGAAGTGGAGGAGCCGCCTTT	
GGCCGAGGCCGCCCCCTTGACGAGATCACCCGGCTCCGC	3000
TACGAGAAGGAGGCCCTGGGGATCTACGTCTCCGGCCACC	
CCATCTTGCGGTACCCCGGGCTCCGGGAGACGGCCACCTG	
CACCCTGGAGGAGCTTCCCCACCTGGCCCGGGACCTGCCG	3120
CCCCGGTCTAGGGTCCTCCTTGCCGGGATGGTGGAGGAGG	
TGGTGCGCAAGCCCACAAAGAGCGGCGGGATGATGGCCCG	
CTTCGTCTCTCCGACGAGACGGGGGCGCTTGAGGCGGTG	3240
GCATTTCGGCCGGGCCTACGACCAGGTCTCCCGAGGCTCA	
AGGAGGACACCCCCGTGCTCGTCCTCGCCGAGGTGGAGCG	
GGAGGAGGGGGGCGTGCGGGTGCTGGCCCAGGCCGTTTGG	3360
ACCTACGAGGAGCTGGAGCAGGTCCCCCGGGCCCTCGAGG	
TGGAGGTGGAGGCCTCCCTCCTGGACGACCGGGGGGTGGC	
CCACCTGAAAAGCCTCCTGGACGAGCACGCGGGGACCTC	3480
CCCCGTGTACGTCCGGGTCCAGGGCGCCTTCGGCGAGGCC	
TCCTCGCCCTGAGGGAGGTGCGGGTGGGGGAGGAGGCTGT	
AGGCGGCCGCGTGGTTCCGGGCCTACCTCCTGCCCGACCG	3600
GGAGGTCTTCTCCAGGGCGGCCAGGCGGGGGAGGCCAG	
GAGGCGGTGCCCTTCTAGGGGGTGGGCCGTGAGACCTAGC	
GCCATCGTTCTCGCCGGGGCAAGGAGGCCTGGGCCCGAC	3720
CCCTTTTGG	

FIG. 16B

MGRELRF	FAHLHQHTQFSL	LDGAPKLSDLLK	WEETTPEDP	
ALAMTDHGNLFG	AVEFYKKATEMGI	KPILGYEAYVAA	ESR	
FDRKR	GKGLDGGYFHLTLL	AKDFTGYQNLVRL	ASRAYLEG	120
FYEKPRIDREIL	REHAEGLI	ALSGCLGAEIPQF	ILQDRLD	
LAEARLNEYLSI	FKDRFFIEIQNHGL	PEQKKVNEVLKEFA		
RKYGLGMVATND	GHYVRKEDARAHEVLL	AIQSKSTLDDPG		240
ALALPCEEFYVKT	PEEMRAMFP	EEEEVGGRSPLTTP	WRSPH	
VQRGAAIGTRW	STRIPRFPLPEGRTE	AQYLMELTFKGLLR		
RYPDRITEGFY	REVFRLSGKLPPHGD	GEALAEALAQVERE		360
AWERLMKSLPPL	AGVKEWTAEAI	FHRALYELSAIERM	GFP	
GLLPHRPG	LHQLGPEKGVSVGP	GRGAAGSLVAYAVG	ITN	
IDPLRFGLL	FERFLNPERVSMPD	IDTDFSDRERDRVIQYV		480
RERYGEDKVAQ	IGTLGSLASKAALKE	VARVYGI	PRKKAEE	
LAKLIPVQFGKPK	PLQEAIQVVP	ELRAEMEKDPKV	REVL	
VAMRLEGLNRHAS	VHAGRGV	FSEPLTDLVPLCATR	KGGP	600
YTQYDMGAVEAL	GLLKMDFLGLR	TLTFLDEVKRIVKASQ		
VELDYDALPLDD	PKTFALLSRGETK	GVFQLESGGMTATLR		
GLKPRRFEDLIA	ILSLYRPGMEHI	PTYIRRHHGLEPVS	Y	720
SEFPHA	EKYLKPILDETYG	IPVYQEQIMQIASA	VAGYSLG	
EADLLRRSMG	KKKVEEMKSHRER	FVQGAKE	RGVPEEEANR	
LFDMLEAF	FANYGFNKSHAAAY	SLLSYQTAYVKA	HYPVEFM	840
AALLSVERHDS	DKVAEYIRDARAMG	IEVLPPDVNRSG	FDF	
LVQGRQILFGL	SAVKNVGEAAAE	AILRERERGGPYR	SLGD	
FLKRLDEKVLN	KRTLES	LIKAGALDGFGERARLL	ASLEGL	960
LKWAAENREKAR	SGMMGLFSEVEEP	PLAEAAPLDEITRLR		
YEKEALGIYVSG	HPILRYPGLRETAT	CTLEELPHLARDLP		
PRSRVLLAGM	VEEVVRKPTKSGG	MARFVLSDETGA	LEAV	1080
AFGRAYDQVSP	RKEDTPVLVLA	EVEREEGGVRVLAQ	AVW	
TYQELEQVPR	ALEVEVEASLPDD	RGVAHLKSLLDEHAG	TL	
PLYVRVQGA	FGEALLALREVRV	GEEALGALEAAGFP	PAYLL	1200
PNREVSPRLT	GS	GPRGRALSTGLALK	TYP	
IALPG	NEAL			
ARPLL				

FIG. 16C

	Start1	Start2	3'-Exo I
T.th.	VERVVRTLLDGRFLLEEGVGLWEWRYPPFLEGEAVVLDLETTGLAG		LDEVIEVGLLRLEGG---RRLPF
D.rad.		PWPQDVVVFDDLETTGFSPA	SAAIVEIGAVRIVGGQIDETLKF
Bac.sub.	HGIKMIYGMEANLVDDGVPIAYNAAHRLLEEETVVVFDVETTGLSAV		YDTIIELAAVKVKGGE---IIDKF
H.inf.		MINPNRQIVLDTETTGMNQLGHAHYEGHCHIEIGAVELINRR	YTGNNX
E.c.		MSTAITRQIVLDTETTGMNQIGAHSEGHKIIIEIGAVEVNNRR	LTGNNF
H.pyl.	NLEYLKACGLNFIETSENLTILKNLKTPLKDEVFSFIDLETTGSCPI		KHEILEIGAVQVKGGE---IINRF

	3'-Exo II
T.th.	QSLVR-PLPP---AEARSWNLT---GIPREALLEAPSLSEVLEKAYPLRGDATLVIHNAAFDLGFL-RPALEGLG
D.rad.	ETLVR-PTRPDGSMLSI PWQAQRVHGISEMVRRAPAXKDVLPDFFDFVDGSVAVAHNVSFDDGGFM-RAGAERLG
Bac.sub.	EAFAN-PHRP---LSATIIELT---GITDDMLQADPDVVDVIRDFREWIGDDILVAHNASFDDMGFL-NVAYKKLL
H.inf.	HIYIK-PDRP---XDPDAIKVH---GITDEMLADKPEFKEVAQDFLDYINGAELLIHNAFPFDDVGFM-DYEFKRLN
E.c.	HVYLK-DRLV---DPEAFGVH---GIAVDFLLDKPTFAEVAVEFMDYIRGAELVHNAAFDVGFM-DYEFSLLK
H.pyl.	ETLVKVKSVP---DYIAELT---GITYEDTLNAPSAHEALQELRLFLGNSVFAHNAFDDYNF LGRYFVEKHLH

	3'-Exo IIIC
T.th.	-----YRLENPVVDSLRLARRGLPGLRRYGLDALSEVLELPRRT--CHRALEDVERTLAVVHEVYMLT-----SG
D.rad.	-----LSWAPERELCTMQLSRRAPFRERTHNLTVLAERLGLEFAPGGRHSYGDVQVTAQAYLRLLLELLG-----ER
Bac.sub.	E----VEKAKNPVIDTLELGRFLYPEFKNHRNLTLCKKFDIELTQ--HHRATYDTEATAYLLLKMLKDA-----EK
H.inf.	-LNVKTDDICLVDTLQMARQMPGKRN-NLDALCDRLGIDNSKRTLHGALLDAEILADVILMMTGGQTNLFDEEE
E.c.	RDIAKTNTFCVKVTDLSAVARKMFPGKRN-SLDALCARYEIDNSKRTLHGALLDAQILAEVYLAMTGGQTSMAFAME
H.pyl.	-----CPLLNLKLTLDLSKRAILSMRY-SLSFLKELLGFGIEV--SHRAYADALASYKLFELCLNLP--SYIKT

FIG.17

FIG.18A

ATGGTGGAGCGGGTGGTGCGGACCCTTCTGGACGGGAGGT 40
 TCCTCCTGGAGGAGGGGGTGGGGCTTTGGGAGTGGCGCTA
 CCCCTTTCCCCTGGAGGGGGAGGCGGTGGTGGTCCTGGAC 120
 CTGGAGACCACGGGGCTTGCCGGCCTGGACGAGGTGATTG
 AGGTGGGCCTCCTCCGCCTGGAGGGGGGGAGGCGCCTCCC 200
 CTTCCAGAGCCTCGTCCGGCCCCCTCCCGCCCCGCCGAAGCC
 CGTTCGTGGAACCTCACCGGCATCCCCCGGGAGGCCCTGG 280
 AGGAGGCCCCCTCCCTGGAGGAGGTTCTGGAGAAGGCCTA
 CCCCCTCCGCGGCGACGCCACCTTGGTGATCCACAACGCC 360
 GCCTTTGACCTGGGCTTCCTCCGCCCCGGCCTTGGAGGGCC
 TGGGCTACCGCCTGGAAAACCCCGTGGTGGACTCCCTGCG 440
 CTTGGCCAGACGGGGCTTACCAGGCCTTAGGCGCTACGGC
 CTGGACGCCCTCTCCGAGGTCCTGGAGCTTCCCCGAAGGA 520
 CCTGCCACCGGGCCCTCGAGGACGTGGAGCGCACCCCTCGC
 CGTGGTGCACGAGGTATACTATATGCTTACGTCCGGCCGT 600
 CCCCCACGCTTTGGGAACTCGGGAGGTAG

MVERVVRTLDDGRFLLEEGVGLWEWRYPFPLEGEAVVVLD 40
 LETTGLAGLDEVIEVGLLRLEGGRRLPFQSLVRPLPPAEA
 RSWNLTGIPREALEEAPSLEEVLEKAYPLRGDATALVIHNA 120
 AFDLGFLRPALEGLGYRLNPVVDLRLARRGLPGLRRYG
 LDALSEVLELPRRTCHRALEDVERTLAVVHEVYYMLTSGR 200
 PRTLWELGRZ

FIG.18B

Alignment of dnaA genes.

P.mar.	MLEASWEK	VQSSL--KQNLK--	-----PSYE	TWIRPTEFSG--FKN	GELTLIAPNSFSSAW	LKNYSQTIQETAE-	65
Syn.sp.	MVSCENLWQQ	ALAIL--ATQLTK--	-----PAFD	TWIKASVLIS--LGD	GVATIQVENGFVLNH	LQKSYGPLLMEVLT-	67
B.sut.	MENILDLWNQ	ALAQI--EKKLSK--	-----PSFE	TWMKSTKAHS--LQG	DTLTITAPNEFARDW	LESRYLHLIADTIY-	67
M.tub.	MTDDPGSGFTVWNA	VVSELNGDPKVDGDP	SSDANLSAPLTPQQR	AWLNLVQPLT--IVE	GFAILSVPSFVQNE	IERRHLRAPITDALS-	87
T.th.	MSHEAVWQH	VLEHI--RRSITE--	-----VEFH	TWFERIRPLG--IRD	GVLELAVPTSFALDW	IRRHVAGLIQEGPR-	66
E.coli	MSLSLWQQ	CLARL--QDELPA--	-----TEFS	MWIRPLQAE--LSD	NTLALYAPNRFVLDW	VRDKYLNININGLLT-	64
T.mar.	MKER	ILQEI--KTRVNR--	-----KSWE	LWFSFVDVKS--IEG	NKVVSFVGNLFIKEW	LEKKVYSVLKAVK-	61
H.pyl.	MDTNNNIEKE	ILALVKQNPVSL--	-----IEYE	NYFSQLKYNPNASKS	DIAFFYAPNQVLCTT	ITAKYGALLKEILSQ	72
P.mar.	EIFG----	VKANAESSDEHYSSA	P-----	---ITPPLEASPGSV	DSSGSSLRLSK----	-KTLPLLNLRVYVFN	130
Syn.sp.	DLTG----	TDGLEPHS---LIGQ	E-----	---SSLPMEITP----	-----	-KNATALNGKYTFSR	115
B.sut.	ELTG----	EELSIFKV	IPQNQDVEDFMKPQ	VKKAVKEDTSDFPQN	-----	-----MLNPKYTFDT	119
M.tub.	RRIGH-QIQLGVRIA	PPATDEADDTTVPPS	ENPATSPDVTIDND	EIDDSAAARGDNQHS	WPSYFTEPERPHNTDSA	TAGVTSINRRYTFDT	176
T.th.	LLGAQ-APRFELRVV	PGVVQEDIFQPPPS	PPAQAP-----	-----	-----	-----EDTFKT	108
E.coli	SFCGADAPQLRFEVG	TKPVTQTPQAAVTSN	VAAPQAQVATQOPORA	APSTRSGWDNVPAPA	EP-----	-TYRSNVNVKHTFDN	140
T.mar.	VVLG----	NDATFEIT	YEAFEPHSSYSEPLV	KKRAVLLTP-----	-----	-----LNPDTYTFEN	106
H.pyl.	NKVG-MHLAHSVDVR	IEVAPKIQINAQSN	NYKAIKTS-----	-----	-----	-----VKDSYTFEN	118
P.mar.	FVVGPNSRMAHAAAM	AVAESPGREENPLFI	CGGVGLGKTHLMQAI	GHYRLEIDPGAKVSY	VSTETFTNDLIL--A	IRQDRMQAFDRDYR-	217
Syn.sp.	FVVGPTNRMHAHAASL	AVAESPGREENPLFL	CGGVGLGKTHLMQAI	AHYRLEWYPNAKVY	VSTERFTNDLIT--A	IRQDNMEDFRSYR-	202
B.sut.	FVIGSGNRFHAHAASL	AVAEAPAKAYNPLFI	YGGVGLGKTHLMHAI	GHYVIDHNPNAKVY	LSSEKFTNEFIN--S	IRDNAKAVDFRNRYR-	206
M.tub.	FVIGASNRFAHAAAL	AIAEAPARAYNPLFI	WGESGLGKTHLLHAA	GNYAQRLFPGMRVKY	VSTEEFTNDFIN--S	LRDDRKVAFKRSYR-	263
T.th.	SMWGPTTPWPHGGAV	AVAESPGRAYNPLFI	YGGRGLGKTVMLHAV	GPLRAKRFPHPMRLEY	VSTETFTNELINRPS	AR-DRMTFRERYR-	196
E.coli	FVEGKSNQLARAAAR	QVADNPGGAYNPLFL	YGGTGLGKTHLLHAV	GNGIMARKPNNAKVY	MHSERFVQDMVK--A	LQNNALIEEFKRYR-	227
T.mar.	FVVGPGNSFAYHAAL	EVAKHPGR--YNPLFI	YGGVGLGKTHLLQSI	GNVYVQNEPDLRVMY	ITSEKFLNDLVD--S	MKEGKLNEFREKYRK	193
H.pyl.	FVVGSCNNTVYELAK	KVAQSDTPPPYNFVLF	YGGTGLGKTHILNAI	GNHALEK--HKKVVL	VTSEDFLTDFLK--H	LDNKTMDSFKAKEYR-	203

FIG.19A

P.mar.	AADLLVDDIQFIEG	KEYTQEEFFHTFNAL	HDAGSQIVLASDRPP	SQIPRLQERLMSRFS	MGLIADVQAPDLETR	MAILQKKAHERVGL	307
Syn.sp.	SADFLLLIDDIQFIKG	KEYTQEEFFHTFNLS	HEAGQVWVASDRAP	QRIPGLQDRLISRFS	MGLIADIQVPDLETR	MAILQKKAEDRIRL	292
B.sut.	NVDVLLIDDIQIFLAG	KEQTQEEFFHTFNIL	HEESQIVISSDRPP	KEIPTLEDRLRSRFE	WGLITDITPPDLETR	IAILRKKAKAEGLDI	296
M.tub.	DVDVLLVDDIQFIEG	KEGIQEEFFHTFNIL	HNANKQIVISSDRPP	KQLATLEDRLRTRFE	WGLITDVQPPPELETR	IAILRKKAKAEMERLAV	353
T.th.	SVDLLLVDDVQFIAG	KERTQEEFFHTFNAL	YEAHKQIILSSDRPP	KDILTLEARLSRFE	WGLITDNPAPELETR	IAILKMNAS-SGPED	285
E.coli	SVDALLIDDIQFFAN	KERSQEEFFHTFNAL	LEGNQIILTSDRYP	KEINGVEDRLKSRFG	WGLITVAIEPPELETR	VAILMKKADENDIRL	317
T.mar.	KVDILLIDDVQFLIG	KTGVQTELFHTFNEL	HDGKQIVICSDREP	QKLSEFQDRLVSRFQ	MGLVAKLEPPDEETR	KSIARKMLEIEHGEEL	283
H.pyl.	HCDFLLDDAQFLQG	KPKLEEEFFHTFNEL	HANSKQIVLISDRSP	KNIAGLEDRLKSRFE	WGITAKVMPDLETK	LSIVKQKQCQLNQITTL	293
P.mar.	PRDLIQFIAGRFTSN	IRELEGALTRAIATA	SITGLPMTVDISIAPM	LD-----PNGQGVEVT	PKQVLDKVAEVFKVT	PDEMRSASRRR-PVS	392
Syn.sp.	PKEVIEYIASHYTSN	IRELEGALIRAIAYT	SLSNVAMTVENIAFV	LN-----PPVEKVAAA	PETIITIVAQHYQLK	VEELLSNSRRR-EVS	377
B.sut.	PNEVMLYIANQIDSN	IRELEGALIRVVAYS	SLINKDINADLAAEA	LKDIIPSSKPKVIT	IKETQVVGQQFNK	LEDFKAKRTK-SVA	384
M.tub.	PDDVLELIIASSIERN	IRELEGALIRVTATA	SLNKTPIDKALAEIV	LRDLI-ADANTMQIS	AATMAATAEYFDTT	VEELRGPGKTR-ALA	441
T.th.	PEDALEYIARQVTSN	IREWEGALMRASPFA	SLNGVELTRAVAACA	LRHLR-P--RELEAD	PLEIIRKAAGPVRPE	TPGGAHGERRKKEVV	372
E.coli	PGEVAFFIAKRLRSN	VRELEGALNRVIANA	NFTGRAITIDFVREA	LRDLL-A-LQEKLV	IDNIQKTVAEYKIK	VADLLSKRRSR-SVA	404
T.mar.	PEEVNLFVAENVDDN	LRRLRGAIKLLVYK	ETTGKVDLKEAILL	LKDFIKPNRVKAMDP	IDELIEIVAKVTGVP	REEILSNSRNV-KAL	372
H.pyl.	PEEVMEYIAQHISDN	IRQMEGAIKISVNA	NLMNASIDLNLAKTV	LEDL--QKDHAEGLSS	LENILLAVAQSLNLK	SSEIKVSSRQK-NVA	380
P.mar.	QARQVGMVLMRQGTN	LSLPRIGDTFGGKDH	TTVMYAIEQVEKKLS	S-----DPQIA	SQVQKIRDLLQIDSR	RKR-----	461
Syn.sp.	LARQVGMVLMRQHTD	LSLPRIGEAFGGKDH	TTVMYSCDKITQLQQ	K-----DWETS	QTILTSLSHRINLAGQ	APES-----	447
B.sut.	FPRQIAMVLSREMTD	SSLPKIGEEFGGRDH	TTVIHAHEKISKLLA	D-----DEQLQ	QHVKEIKEQLK----	-----	446
M.tub.	QSRQIAMVLCRELTD	LSLPKIGQAFG-RDH	TTVMYAQRKILSEMA	E-----RREVF	DHVKELTTRIRQSK	R-----	507
T.th.	LPRQLAMVILVRELTP	ASLPEIGQLFGGRDH	TTVRVAIQVQELAG	KP-----DREVQ	GLLRTLREACTDPVD	NLWITCG	446
E.coli	RPRQWAMALAKELTN	HSLPEIGDAFGGRDH	TTVLHACRKIEQLRE	E-----SHDIK	EDFSNLRTLSS----	-----	467
T.mar.	TARRIGMVAKNVLYK	SSLRTIAEFN-RSH	PVVVDSVKVKVDSLL	KG-----NKQLK	ALIDEVIGEISRRAL	SG-----	440
H.pyl.	LARKLVVYFARLYTP	NPTLSLAQFLDLKDH	SSISKMYSGVKKMLE	EEKSPFVLSLREEIK	NRLNEINDKKTAFFNS	SE-----	457

FIG.19B

GTGTCGCACGAGGCCGTCTGGCAACACGTTCTGGAGCA⁻CA
 TCCGCCGCAGCATCACCGAGGTGGAGTTCCACACCTGGTT
 TGAAAGGATCCGCCCTTGGGGATCCGGGACGGGGTGCTG 120
 GAGCTCGCCGTGCCACCTCCTTTGCCCTGGACTGGATCC
 GGCGCCACTACGCCGGCCTCATCCAGGAGGGCCCTCGGCT
 CCTCGGGGGCCAGGCGCCCCGGTTTGAGCTCCGGGTGGTG 240
 CCCGGGGTTCGTAGTCCAGGAGGACATCTTCCAGCCCCCGC
 CGAGCCCCCGGCCCAAGCTCAACCCGAAGATACCTTTAA
 AACTTCGTGGTGGGGCCCAACAACCTCCATGGCCCCACGGC 360
 GGCGCCGTGGCCGTGGCCGAGTCCCCCGGCCGGGCCTACA
 ACCCCCTCTTCATCTACGGGGGCCGTGGCCTGGGAAAGAC
 CTACCTGATGCACGCCGTGGGCCACTCCGTGCGAAGCGC 480
 TTCCCCCACATGAGATTAGAGTACGTTTCCACGGAAACTT
 TCACCAACGAGCTCATCAACCGGCCATCCGCGAGGGACCG
 - GATGACGGAGTTCCGGGAGCGGTACCGCTCCGTGGACCTC 600
 CTGCTGGTGGACGACGTCCAGTTCATCGCCGGAAGGAGC
 GCACCCAGGAGGAGTTTTTCCACACCTTCAACGCCCTTTA
 CGAGGCCCACAAGCAGATCATCCTCTCCTCCGACCGGCCG 720
 CCCAAGGACATCCTCACCCCTGGAGGCGCGCCTGCGGAGCC
 GCTTTGAGTGGGGCCTGATCACCGACAATCCAGCCCCCGA
 CCTGGAAACCCGGATCGCCATCCTGAAGATGAACGCCAGC 840
 AGCGGGCCTGAGGATCCCGAGGACGCCCTGGAGTACATCG
 CCCGGCAGGTCACCTCCAACATCCGGGAGTGGGAAGGGGC
 CCTCATGCGGGCATCGCCTTTTCGCCTCCCTCAACGGCGTT 960
 GAGCTGACCCGCGCCGTGGCGGCCAAGGCTCTCCGACATC
 TTCGCCCCAGGGAGCTGGAGGCGGACCCCTTGGAGATCAT
 CCGCAAAGCGGCGGGACCAGTTCGGCCTGAAACCCCGGGA 1080
 GGAGCTCACGGGGAGCGCCGCAAGAAGGAGGTGGTCCTCC
 CCCGGCAGCTCGCCATGTACCTGGTGCGGGAGCTCACCCC
 GGCCTCCCTGCCCGAGATCGACCAGCTCAACGACGACCGG 1200
 GACCACACCACGGTCCTCTACGCCATCCAGAAGGTCCAGG
 AGCTCGCGGAAAGCGACCGGGAGGTGCAGGGCCTCCTCCG
 CACCTCCGGGAGGCGTGCACATGA

FIG.20A

VSHEAVWQHVLHIRRSITEVEFHTWFERIRPLGIRDGVL
ELAVPTSFALDWIRRHAYAGLIQEGPRLPGAQAPRFELRVV
- PGVVVQEDIFQPPSPPAQAQPEDTFKTSWWGPTTPWPHG 120
GAVAVAESPGRAYNPLFIYGGRGLGKTYLMHAVGPLRAKR
FPHMRLEYVSTETFTNELINRPSARDRMTEFRERYRSVDL
LLVDDVQFIAGKERTQEEFFHTFNALYEAHKQIILSSDRP 240
PKDILTLEARLSRFEWGLITDNPAPDLETRIAILKMNAS
SGPEDPEDALEYIARQVTSNIREWEGALMRASPFASLNGV
ELTRAVAAKALRHLRPRELEADPLEIIRKAAGPVRPETPG 360
GAHGERRKKEVVLPRLAMYLVLRELTPASLPEIDQLNDDR
DHTTVLYAIQKVQELAESDREVQGLLRTLREACT

FIG.20B

ATGAACATAACGGTTCCCAAAAACTCCTCTCGGACCAGC 40
 TTTCCCTCCTGGAGCGCATCGTCCCCTCTAGAAGCGCCAA
 CCCCTCTACACCTACCTGGGGCTTTACGCCGAGGAAGGG 120
 GCCTTGATCCTCTTCGGGACCAACGGGGAGGTGGACCTCG
 AGGTCCGCCTCCCCGCCGAGGCCCAAAGCCTTCCCCGGGT 200
 GCTCGTCCCCGCCAGCCCTTCTTCCAGCTGGTGC GGAGC
 CTTCTTGGGGACCTCGTGGCCCTCGGCCTCGCCTCGGAGC 280
 CGGGCCAGGGGGGGCAGCTGGAGCTCTCCTCCGGGCGTTT
 CCGCACCCGGCTCAGCCTGGCCCCCTGCCGAGGGCTACCCC 360
 GAGCTTCTGGTGCCCGAGGGGGAGGACAAGGGGGCCTTCC
 CCTCCGGACGCGGATGCCCTCCGGGGAGCTCGTCAAGGC 440
 CTTGACCCACGTGCGCTACGCCGCGAGCAACGAGGAGTAC
 CGGGCCATCTTCCGCGGGGTGCAGCTGGAGTTCTCCCCC 520
 AGGGCTTCCGGGCGGTGGCCTCCGACGGGTACCGCCTCGC
 CCTCTACGACCTGCCCCCTGCCCCAAGGGTTCCAGGCCAAG 600
 GCCGTGGTCCCCGCCCGAGCGTGGACGAGATGGTGC GGG
 TCCTGAAGGGGGCGGACGGGGCCGAGGCCGTCTCGCCCT 680
 GGGCGAGGGGGTGTGGCCCTGGCCCTCGAGGGCGGAAGC
 GGGGTCCGGATGGCCCTCCGCCTCATGGAAGGGGAGTTCC 760
 CCGACTACCAGAGGGTCATCCCCCAGGAGTTCGCCCTCAA
 GGTCCAGGTGGAGGGGGAGGCCCTCAGGGAGGCGGTGCGC 840
 CGGGTGAGCGTCTCTCCGACCGGCAGAACCACCGGGTGG
 ACCTCCTTTTGGAGGAAGGCCGGATCCTCCTCTCCGCCGA 920
 GGGGGACTACGGCAAGGGGCAGGAGGAGGTGCCCGCCAG
 GTGGAGGGGGCCGGACATGGCCGTGGCCTACAACGCCCGCT 1000
 ACCTCCTCGAGGCCCTCGCCCCCGTGGGGGACCGGGCCCA
 CCTGGGCATCTCCGGGCCCACGAGCCCGAGCCTCATCTGG 1080
 GGGGACGGGGAGGGGTACCGGGCGGTGGTGGTGCCCCCTCA
 GGGTCTAG 1128

FIG.21A

MNITVPPKLLSDQLSLLERIVPSRSANPLYTYLGGLYAEEG 40
 ALILFGTNGEVDLEVRLPAEAQSLPRVLVPAQPFFQLVRS
 LPGDLVALGLASEPGQGGQLELSSGRFRTRLAPAEGLYP 120
 - ELLVPEGEDKGAFPLRTRMPGELVKALTHVRYAASNEEY
 RAIFRGVQLEFSPQGFRAVASDGYRLALYDLPLPQGFQAK 200
 AVVPARSVDEMVRVLKGADGAEAVLALGEGVLALALEGGS
 GVRMALRLMEGEFPDYQRVIPQEFALKVQVEGEALREAVR 280
 RVSVLSDRQNHVRVDLLLEEGRILLSAEGDYGKGQEEVPAQ
 VEGPDMAVAYNARYLLEALAPVGDRAHLGISGPTSPSLIW 360
 GDGEGYRAVVVPLRVZ

FIG.21B

T.th.beta	MNITVPKLLSDQLSLLEIRIVPSRSANPLYTYLGLYAEAGALILFGTNGEVDLEVRPAE
E.coli.bet	MKFTVEREHLKPLQQVSGPLGGRPTLPILGNLLQVADGTLSTGTDLEMENVARVALV
P.mirab.be	MKFIIEREQLKPLQQVSGPLGGRPTLPILGNLLKVTENTLSLTGTDLEMENMARVSL
H.infl.bet	MQFSISRENLLKPLQQVCGVLSNRPNIPVLNNVLLQIEDYRLTITGTDLEVELSSQTOLS
P.put.beta	MHFTIQREALKPLQVAGVVERRQTLPVLSNVLVQQQLSLTGTDEVELVGRVQLE
B.cap.beta	MKFTIQNDILTKNLKKITRVLVKNISFPILNLIQVEDGTLSTTNLEIELISKIEII
	* . . * * . . * *
T.th.beta	AQSLP-RVLVPAQFFQVLRSLPGDLVALGLASEPGGQQLLELSSGRFRTRLSLAPAEY
E.coli.bet	QPHEPGATTVPARKFFDICRGLP-EGAEIAVQLE---GERMLVRSGRSFRSLSTLPAADF
P.mirab.be	QSHEIGATTVPARKFFDIWRGLP-EGAEISVELD---GDRLLVRSGRSFRSLSTLPASDF
H.infl.bet	SSSENGTFTIPAKKFLDICRTLS-DDSEITVTPE---QDRALVQSGRSRFTLATQPAEEY
P.put.beta	EPAEPGEITVPARKLMDICKSLP-NDALIDIKVD---EQKLLVKAGRSRFTLSTLPANDF
B.cap.beta	TKYIPGKTTISGRKILNICRTLS-EKSKIKMQLK---NKKMYISSENSNYILSTLSADTF
 *
T.th.beta	PELLVPEGEDKGAFLRTPMPSGELVKALTHVRYAASNEEYRAIFRGVQLEFSPQGFRVAV
E.coli.bet	PNLDD--WQSEVEFTLPQAT-----MKRLIEATQFSMAHQDVRYVYINGMLFETEGEELRTV
P.mirab.be	PNLDD--WQSEVEFTLPQAT-----LKRLIESTQFSMAHQDVRYVYINGMLFETENTEELRTV
H.infl.bet	PNLTD--WQSEVDFELPQNT-----LRRLIEATQFSMANQDARYFLNGMKFETEGNLLRTV
P.put.beta	PTVEE--GPGSLTCNLEQSK-----LRRLIERTSFAMAQQDVRYVYINGMLLEVSRLTRAV
B.cap.beta	PNHQN--FDYISKFDISSNI-----LKEMIEKTEFSMGKQDVRYVYINGMLLEKKDKFLRSV
	* *
T.th.beta	ASDGYRLALYDLPLPQGFQA--KAVVPARSVDEMVRVLKGADGAEAVLALGEGVLALALE
E.coli.bet	ATDGHRLAVCSMPIGQSLPS-HSVIVPRKGVIELMRMLDG-GDNPLRVQIGSNNIRAHVG
P.mirab.be	ATDGHRLAVCAMDIGQSLPG-HSVIVPRKGVIELMRLLDGSGESLLQLQIGSNNLRAHVG
H.infl.bet	ATDGHRLAVCTISLEQELQN-HSVILPRKGVLELVRLLLET-NDEPARLQIGTNNLRLVHLK
P.put.beta	STDGHRLALCMSAPIEQEDRHQVIVPRKGILLELRLITD-PEGMVSVIVLGQHHIRATTG
B.cap.beta	ATDGYRLAISYTQLKKDINF-FSIIIPNKAVMELLKLLNT-QPQLNLIIGSNSIRIYTK
	..* ** *

FIG.22A

T.th.beta	GGSGVRMALRLMEGEFPDYQRVI PQEFALKVQVEGEALREAVRVSLSDRQNHVRVDLLL
E.coli.bet	---DFIFTSKLVDGRFPDYRRVL PKNPDKHLEAGCDLLKQAFARAAILSNEKFRGVRLYV
P.mirab.be	---DFIFTSKLVDGRFPDYRRVL PKNPTKTVIAGCDILKQAFSRAAILSNEKFRGVRINL
H.infl.bet	---NTVFTSKLIDGRFPDYRRVL PRNATKIVEGNWEMLKQAFARASILSNERARSVRLSL
P.put.beta	---EFTFTSKLVDGKFPDYERVL PKGGDKLVGDRQALREAFSRTAILSNEKYRGIRLQL
B.cap.beta	---NLIFTTQLIEGEYPDYKSVLFKEKKNPITITNSILLKSLRLRVAILAHEKFCGIEIKI . * . . . * . . . * . . . * . . . * . . .
T.th.beta	EEGRILLSAEGDYGK-GQEEVPAQVEGPDMAVAYNARYLLEALAPVG-DRAHLGISGPTS
E.coli.bet	SENQLKITANNPEQEEAEIILDVTYSGAEMEIGFNVSYLLDVLNALKCENVRMMLTDSVS
P.mirab.be	TNGQLKITANNPEQEEAEIIVDVQYQGEEMEIGFNVSYLLDVLNTILKCEEVKLLLTDAVS
H.infl.bet	KENQLKITASNTHEHEEAEIIVDVNYNGEELEVGFNVTYLLDVLNALKCNQVRMCLTDAFS
P.put.beta	AAGQLKIQANNPEQEEAEIISVDYEGSSLEIGFNVSYLLDVLGVMTEQVRLILSDSNS
B.cap.beta	ENGKFKVLSDNQEEETAEDLFEIDYFGEKIEISINVYLLDVLINNICKSENIALFLNKSKS * . . . * . . . * . . . * . . .
T.th.beta	PSLIWGDG-EGYRAVVVPLRVZ (ID#108)
E.coli.bet	SVQIEDAASQSAAYVVMPMRLZ (ID#109)
P.mirab.be	SVQVENVASAAAAAYVVMPMRL- (ID#110)
H.infl.bet	SCLIENCEDSSCEYVIMPML- (ID#111)
P.put.beta	SALLQEAGNDSSYVVMPMRL- (ID#112)
B.cap.beta	SIQIEAENNSSNAYVVMMLKR- (ID#113) *

FIG.22B





FIG.23


FIG.24A Induction



↓
Lysis
↓
Heat Step
↓

FIG.24B MonoQ Column

Fraction: 5 7 9 11 13 15 17 19 21 23 25







FIG.25

FIG.25A





FIG.25B

FIG. 26A





FIG. 26B



FIG. 27




FIG. 28




FIG. 29




FIG. 30




FIG. 31




FIG. 32

α




FIG. 33A

β




FIG. 33B

$\tau\delta\delta'$




FIG. 33C

SSB




FIG. 33D

Pol III*




FIG. 33E

ATGAGTAAGGATTTTCGTCCACCTTCACCTGCACACCCAGTTCTCACTCCT	
GGACGGGGCTATAAAGATAGACGAGCTCGTGAAAAAGGCAAAGGAGTATG	100
GATACAAAGCTGTCGGAATGTCAGACCACGGAAACCTCTTCGGTTTCGTAT	
AAATTCTACAAAGCCCTGAAGGCGGAAGGAATTAAGCCCATAATCGGCAT	200
GGAAGCCTACTTTACCACGGGTTTCGAGGTTTGACAGAAAGACTAAAACGA	
GCGAGGACAACATAACCGACAAGTACAACCACCACCTCATACTTATAGCA	300
AAGGACGAAAAGGTCTAAAGAACTTAATGAAGCTCTCAACCCTCGCCTAC	
AAAGAAGGTTTTTACTACAAACCCAGAATTGATTACGAACTCCTTGAAAA	400
GTACGGGGAGGGCCTAATAGCCCTTACCGCATGCCTGAAAGGTGTTCCCA	
CCTACTACGCTTCTATAAACGAAGTGAAAAAGGCGGAGGAATGGGTAAAG	500
AAGTTCAAGGATATATTCGGAGATGACCTTTATTTAGAACTTCAAGCGAA	
CAACATTCCAGAACAGGAAGTGGCAAACAGGAACTTAATAGAGATAGCCA	600
AAAAGTACGATGTGAAACTCATAGCGACGCAGGACGCCCCTACCTCAAT	
CCCGAAGACAGGTACGCCCACACGGTTCTTATGGCACTTCAAATGAAAAA	700
GACCATTACGAACTGAGTTCGGGAAACTTCAAGTGTTCAAACGAAGACC	
TTCACTTTTGCTCCACCCGAGTACATGTGGAAAAAGTTTGAAGGTAAGTTC	800
GAAGGCTGGGAAAAGGCACTCCTGAACACTCTCGAGGTAATGGAAAAGAC	
AGCGGACAGCTTTGAGATATTTGAAAACCTCCACCTACCTCCTTCCCAAGT	900
ACGACGTTCCGCCCCGACAAAACCTTGAGGAATACCTCAGAGAACTCGCG	
TACAAAGGTTTTAAGACAGAGGATAGAAAGGGGACAAGCTAAGGATACTAA	1000
AGAGTACTGGGAGAGGCTCGAGTACGAACTGGAAGTTATAAACAAAATGG	
GCTTTGCGGGATACTTCTTGATAGTTCAGGACTTCATAAACTGGGCTAAG	1100
AAAAACGACATACCTGTTGGACCCGGAAGGGGAAGTGCTGGAGGTTCCCT	
CGTCGCATACGCCATCGGAATAACGGACGTTGACCCTATAAAGCACGGAT	1200
TCCTTTTTGAGAGGTTCTTAAACCCCGAAAGGGTTTCCATGCCGGATATA	
GACGTGGATTTCTGTCTCAGGACAACAGGGAAAAGGTCATAGAGTACGTAAG	1300
GAACAAGTACGGACACGACAACGTAGCTCAGATAATCACCTACAACGTAA	
TGAAGGCGAAGCAAACACTGAGAGACGTGCAAGGGCCATGGGACTCCCC	1400
TACTCCACCGCGGACAAACTCGCAAAACTCATTCTCAGGGGGACGTTCA	
GGGAACGTGGCTCAGTCTGGAAGAGATGTACAAAACGCCTGTGGAGGAAC	1500
TCCTTCAGAAGTACGGAGAACACAGAACGGACATAGAGGACAACGTAAAG	
AAGTTCAGACAGATATGCGAAGAAAAGTCCGGAGATAAAACAGCTCGTTGA	1600
GACGGCCCTGAAGCTTGAAGGTCTCACGAGACACACCTCCCTCCACGCCG	
CGGGAGTGGTTATAGCACCAAAGCCCTTGAGCGAGCTCGTTCCCCTCTAC	1700
TACGATAAAGAGGGCGAAGTCGCAACCCAGTACGACATGGTTTCAGCTCGA	
AGAACTCGGTCTCCTGAAGATGGACTTCCTCGGACTCAAACCCCTCACAG	1800
AACTGAAACTCATGAAAGAACTCATAAAGGAAAGACACGGAGTGGATATA	
AACTTCCTTGAACTTCCCCTTGACGACCCGAAAGTTTACAAACTCCTTCA	1900
GGAAGGAAAAACACGGGAGTGTTCCAGCTCGAAAGCAGGGGAATGAAAG	
AACTCCTGAAGAACTAAAGCCCGACAGCTTTGACGACATCGTTGCGGTC	2000
CTCGCACTCTACAGACCCGGACCTCTAAAGAGCGGACTCGTTGACACATA	
CATTAAGAGAAAGCACGGAAAAGAACCCGTTGAGTACCCCTTCCCGGAGC	2100
TTGAACCCGTCCTTAAGGAAACCTACGGAGTAATCGTTTATCAGGAACAG	
GTGATGAAGATGTCTCAGATACTTTCCGGCTTTACTCCCGGAGAGGCGGA	2200
TACCCTCAGAAAGGCGATAGGTAAGAAGAAAGCGGATTTAATGGCTCAGA	
TGAAAGACAAGTTCATACAGGGAGCGGTGGAAAGGGGATACCCTGAAGAA	2300
AAGATAAGGAAGCTCTGGGAAGACATAGAGAAGTTCGTTTCTACTCCTT	
CAACAAGTCTCACTCGGTAGCTTACGGGTACATCTCCTACTGGACCGCCT	2400

FIG. 34A

ACGTTAAAGCCCACTATCCCGCGGAGTTCTTCGCGGTAAAACTCACAACT
 GAAAAGAACGACAACAAGTTCCTCAACCTCATAAAAGACGCTAAACTCTT 2500
 CGGATTTGAGATACTTCCCCCGACATAAACAAGAGTGATGTAGGATTTA
 CGATAGAAGGTGAAAACAGGATAAGGTTCTGGGCTTGCGAGGATAAAGGGA 2600
 GTGGGAGAGGAAACTGCTAAGATAATCGTTGAAGCTAGAAAGAAGTATAA
 GCAGTTCAAAGGGCTTGCGGACTTCATAAACAAAACCAAGAACAGGAAGA 2700
 TAAACAAGAAAGTCGTGGAAGCACTCGTAAAGGCAGGGGCTTTTGACTTT
 ACTAAGAAAAAGAGGAAAGAACTACTCGCTAAAGTGGCAAACCTCTGAAAA 2800
 AGCATTAATGGCTACACAAAACCTCCCTTTTCGGTGCACCGAAAGAAGAAG
 TGGAAGAACTCGACCCCTTAAAGCTTGAAAAGGAAGTTCTCGGTTTTTAC 2900
 ATTTTCAGGGCACCCCTTGACAACTACGAAAAGCTCCTCAAGAACCGCTA
 CACACCCATTGAAGATTTAGAAGAGTGGGACAAGGAAAGCGAAGCGGTGC 3000
 TTACAGGAGTTATCACGGAACCTCAAAGTAAAAAAGACGAAAAACGGAGAT
 TACATGGCGGTCTTCAACCTCGTTGACAAGACGGGACTAATAGAGTGTGT 3100
 CGTCTTCCCGGGAGTTTACGAAGAGGCAAAGGAACTGATAGAAGAGGACA
 GAGTAGTGGTAGTCAAAGGTTTTCTGGACGAGGACCTTGAAACGGAAAAT 3200
 GTCAAGTTTCGTGGTGAAAGAGGTTTTCTCCCCTGAGGAGTTCGCAAAGGA
 GATGAGGAATACCCTTTATATATTCTTAAAAAGAGAGCAAGCCCTAAACG 3300
 GCGTTGCCGAAAAACTAAAGGGAATTATTGAAAACAACAGGACGGAGGAC
 GGATACAACTTGGTTCTCACGGTTGATCTGGGAGACTACTTCGTTGATTT 3400
 AGCACTCCCACAAGATATGAACTAAAGGCTGACAGAAAGGTTGTAGAGG
 AGATAGAAAAACTGGGAGTGAAGGTCATAATTTAGTAAATAACCCTTACT 3500
 TCCGAGTAGTCCCC

FIG. 34B

MSKDFVHLHLHTQFSLLDGAIKIDELVKKAKEYGYKAVGMSDHGNI FGSY	
KFYKALKAEGIKPIIGMEAYFTTGSRFDRKTKTSEDNITDKYNHHLILIA	100
KDDKGLKLNLMKLSTLAYKEGFYKPRIDYELLEKYGEGLIALTACLKGV	
TYYASINEVKKAEWVKKFKDIFGDDLYLELQANNIPEQEVANRNLI EIA	200
KKYDVKLIATQDAHYLNPEDRYAHTVLMALQMKKTIHELSSGNFKCSNED	
LHFAPPEYMWKKFEGKFEGWEKALLNTLEVMEKTADSFEIFENSTYLLPK	300
YDVPPDKTLEEYLRELAYKGLRQRIERGQAKDTKEYWERLEYELEVINKM	
GFAGYFLIVQDFINWAKKNDIPVGPGRGSAGGSLVAYAIGITDVDPIKHG	400
FLFERFLNPERVSMPI DIDVDFCQDNREKVIEYVRNKYGHDNVAQIIITYNV	
MKAKQTLRDVARAMGLPYSTADKLAKLIPQGDVQGTWLSLEEMYKTPVEE	500
LLQKYGEHRDIEDNVKKFRQICEESPEIKQLVETALKLEGLTRHTSLHA	
AGVVIAPKPLSELVPLYDKEGEVATQYDMVQLEELGLLKMDFLGLKTLT	600
ELKLMKELIKERHGVDINFLELPLDDPKVYKLLQEGKTTGVFQLESRGMK	
ELLKKLKPDSFDDIVAVLALYRPGPLKSGLVDTYIKRKHGKEPVEYPFPE	700
LEPVLKETYGVI VYQEQVMKMSQILSGFTPGEADTLRKAIGKKKADLMAQ	
MKDKFIQGAVERGYPEEKIRKLWEDIEKFASYSFNKSHSVAYGYISYWTA	800
YVKAHYPAEFFAVKLTTEKNDNKFLNLIKDAKLFGFEILPPDINKSDVGF	
TIEGENRIRFGLARIKGVGEETAKIIVEARKKYKQFKGLADFINKTKNRK	900
INKKVVEALVKAGAFDFTKKRKELLAKVANSEKALMATQNSLFGAPKEE	
VEELDPLKLEKEVLGFYISGHPLDNYEKLLKNRYTPIEDLEEWDKSEAV	1000
LTGVITELKVKKTKNGDYMAVFNLVDKTGLIECVVFPGVYEEAKELIEED	
RVVVVKGFLEDEDLETENVKFVVKVFSPEEFAKEMRNTLYIFLKREQALN	1100
GVAEKLKGI IENNRTEDGYNLVLTVDLGDYFVDLALPQDMKLLKADRKVVE	
EIEKLGVKVII	1161

FIG. 35

ATGAACTACGTTCCCTTCGCGAGAAAGTACAGACCGAAATTCTTCAGGGA
 AGTAATAGGACAGGAAGCTCCCGTAAGGATACTCAAAAACGCTATAAAAA 100
 ACGACAGAGTGGCTCACGCCTACCTCTTTGCCGGACCGAGGGGGGTTGGG
 AAGACGACTATTGCAAGAATTCTCGCAAAGCTTTGAACTGTAAAAATCC 200
 CTCAAAGGTGAGCCCTGCGGTGAGTGCGAAAAGTGCAGGGAGATAGACA
 GGGGTGTGTTCCCTGACTTAATTGAAATGGATGCCGCCTCAAACAGGGGT 300
 ATAGACGACGTAAGGGCATTAAAAGAAGCGGTCAATTACAAACCTATAAA
 AGGAAAGTACAAGGTTTACATAATAGACGAAGCTCACATGCTCACGAAAG 400
 AAGCTTTCAACGCTCTCTTAAAAACCCTCGAAGAGCCCCCTCCCAGAACT
 GTTTTCGTCCTTTGTACCACGGAGTACGACAAAATTCTTCCCACGATACT 500
 CTCAAGGTGTGAGAGGATAATCTTCTCAAAGGTAAGAAAGGAAAAAGTAA
 TAGAGTATCTAAAAAGATATGTGAAAAGGAAGGGATTGAGTGCGAAGAG 600
 GGAGCCCTTGAGGTTCTGGCTCATGCCTCTGAAGGGTGCATGAGGGATGC
 AGCCTCTCTCCTGGACCAGGCGAGCGTTTACGGGGAAGGCAGGGTAACAA 700
 AAGAAGTAGTGGAGAACTTCCTCGGAATTCTCAGTCAGGAAAGCGTTAGG
 AGTTTTCTGAAATTGCTTCTGAACTCAGAAGTGGACGAAGCTATAAAGTT 800
 CCTCAGAGAACTCTCAGAAAAGGGCTACAACCTGACCAAGTTTTTGGGAGA
 TGTTAGAAGAGGAAGTGAGAAACGCAATTTTAGTAAAGAGCCTGAAAAAT 900
 CCCGAAAGCGTGGTTCAGAACTGGCAGGATTACGAAGACTTCAAAGACTA
 CCCTCTGGAAGCCCTCCTCTACGTTGAGAACCTGATAAACAGGGGTAAAG 1000
 TTGAAGCGAGAACGAGAGAACCCTTAAGAGCCTTTGAACTCGCGGTAATA
 AAGAGCCTTATAGTCAAAGACATAATTCCTGTATCCCAGCTCGGAAGTGT 1100
 GGTAAGGAAACCAAAAAGGAAGAAAAGAAAGTTGAAGTAAAAGAAGAGC
 CAAAAGTAAAAGAAGAAAAACCAAAGGAGCAGGAAGAGGACAGGTTCCAG 1200
 AAAGTTTTTAAACGCTGTGGACGGCAAAATCCTTAAAAGAATACTTGAAGG
 GGCAAAAAGGGAAGAAAGAGACGGAAAAATCGTCCTAAAGATAGAAGCCT 1300
 CTTATCTGAGAACCATGAAAAAGGAATTTGACTCACTAAAGGAGACTTTT
 CCTTTTTTTAGAGTTTGAACCCGTGGAGGATAAAAAAAAACCTCAGAAGTC 1400
 CAGCGGGACGAGGCTGTTTTAAAGGTAAAGGAGCTCTTCAATGCAAAAAT
 ACTCAAAGTACGAAGTAAAAGCTAAGGTCATAAAGGTGAGAATGCCCGTG 1500
 GAAGAGATAGGGCTGTTTAACGCACTAATAGACGGCTTGCCCAGGTACGC
 ACTCACGAGGACGAAGGAAAAGGGAAAGGGAGAAGTTTTTCGTTTTAGCGA 1600
 CTCCTTATAAAGTCAAGGAATTGATGGAAGCTATGGAGGGTATGAAAAAA
 CACATAAAGGATTTAGAAATCCTCGGAGAGACGGATGAGGATTTAACTTT 1700
 TTAAAGTATGGGTGTATCTGAGCAAAGGTTTAAGCTAAAAACAAACCTGA
 AACCCGCAGGGGACCAGCCGAAAGCCATAAAAAAACTCCTTGAAAACCTA 1800
 AGGAAAGGCGTAAAAGAACAAACACTTCTCGGAGTCACGGGAAGCGGAAA
 GACTTTTACTCTAGCAAACGTAATAGCGAAGTACAACAAACCAACTCTTG 1900
 TGGTAGTTCACAACAAAATTCTCGCGGCACAGCTATACAGGGAGTTTAAA
 GAACTATTCCCTGAAAACGCTGTAGAGTACTTTGTCTCTTACTACGACTA 2000
 TTACCAACCTGAAGCCTACATTCCCGAAAAAGATTTATACATAGAAAAGG
 ACGCGAGTATAAACGAAAGCTGGAACGTTTCAGACACTCCGCCACGATAT 2100
 CCGTTCTAGAAAGGAGGGACGTTATAGTAGTTGCTTCAGTTTCTTGCATA
 TACGGACTCGGGAAACCTGAGCACTACGAAAACCTGAGGATAAACTCCA 2200
 AAGGGGAATAAGACTGAACTTGAGTAAGCTCCTGAGGAACTCGTTGAGC
 TAGGATATCAGAGAAATGACTTTGCCATAAAGAGGGCTACCTTCTCGGTT 2300
 AGGGGAGACGTGGTTGAGATAGTCCCTTCTCACACGGAAGATTACCTCGT
 GAGGGTAGAGTTCTGGGACGACGAAGTTGAAAGAATAGTCCTCATGGACG 2400
 CTCTGAAC

FIG. 36

ATGCGCGTTAAGGTGGACAGGGAGGAGCTTGAAGAGGTTCTTAAAAAAGC	
AAGAGAAAGCACGGAAAAAAAAGCCGCACTCCCGATACTCGCGAACTTCT	100
TACTCTCCGCAAAAGAGGAAAACTTAATCGTAAGGGCAACGGACTTGGAA	
AACTACCTTGTAGTCTCCGTAAAGGGGGAGGTTGAAGAGGAAGGAGAGGT	200
TTGCGTCCACTCTCAAAAACTCTACGATATAGTCAAGAACTTAAATTCCG	
CTTACGTTTACCTTCATACGGAAGGTGAAAACTCGTCATAACGGGAGGA	300
AAGAGTACGTACAAACTTCCGACAGCTCCCGCGGAGGACTTTCCCGAATT	
TCCAGAAATCGTAGAAGGAGGAGAAACACTTTCGGGAAACCTTCTCGTTA	400
ACGGAATAGAAAAGGTAGAGTACGCCATAGCGAAGGAAGAAGCGAACATA	
GCCCTTCAGGGAATGTATCTGAGAGGATACGAGGACAGAATTCACCTTGT	500
GTTTCGGACGGTCACAGGCTTGCACTTTATGAACCTCTACGTAAACATTGA	
AAAGAGTGAAGACGAGTCTTTTGCTTACTTCTCCACTCCCGAGTGGAAAC	600
TCGCCGTTAGCTCCTGGAAGGAGAATTCCCGGACTACATGAGTGTATCC	
CTGAGGAGTTTTTCGGCGGAAGTCTTGTTTGAGACAGAGGAAGTCTTAAAG	700
GTTTTAAAGAGGTTGAAGGCTTTAAGCGAAGGAAAAGTTTTTCCCGTGAA	
GATTACCTTAAGCGAAAACCTTGCCATCTTTGAGTTCGCGGATCCGGAGT	800
TCGGAGAAGCGAGAGAGGAAATTGAAGTGGAGTACACGGGAGAGCCCTTT	
GAGATAGGATTCAACGGAAATACCTTATGGAGGCGCTTGACGCCTACGAC	900
AGCGAAAGAGTGTGGTTCAAGTTCACAACCCCGACACGGCCACTTTATT	
GGAGGCTGAAGATTACGAAAAGGAACCTTACAAGTGCATAATAATGCCGA	1000
TGAGGGTGTAGCCATGAAAAAGCTTTAATCTTTTATTGAGCTTGAGCC	
TTTTAATTCTGCGTTTAGCGAAGCCAAACCCAAGTCTTC	1090

FIG. 38

MRVKVDREELEEVLLKKARESTEKKAALPILANFLLSAKEENLIVRATDLE	
NYLVSVKGEVEEEGEVVCVHSQKLYDIVKNLNSAYVYLHTEGEKLVITGG	100
KSTYKLPTAPAEDFPEFPEIVEGGETLSGNLLVNGIEKVEYAIKKEANI	
ALQGMYLRGYEDRIHFVGS DGHRLALYEPLGEFSKELLI PRKSLKVLKKL	200
ITGIEDVNIEKSEDES FAYFSTPEWKLAVRLLEGEFPDYMSV IPEEFSAE	
VLFEETEEVLKVLKRLKALSEGKVFPVKITLSENLAIFE FADPEFGEAREE	300
IEVEYTGEPFEIGFNGKYLMEALDAYDSERVWFKFTTPDTATLLEAEDYE	
KEPYKCIIMPMRV	363

FIG. 39

GTGGAAACCACAATATTCCAGTTCAGAAAACCTTTTTTCACAAAACCTCC
 GAAGGAGAGGGTCTTCGTCCTTCATGGAGAAGAGCAGTATCTCATAAGAA 100
 CCTTTTTGTCTAAGCTGAAGGAAAAGTACGGGGAGAATTACACGGTTCTG
 TGGGGGGATGAGATAAGCGAGGAGGAATTCTACACTGCCCTTTCCGAGAC 200
 CAGTATATTCCGCGGTTCAAAGGAAAAAGCGGTGGTCATTTACAACTTCG
 GGGATTTCTGAAGAAGCTCGGAAGGAAGAAAAAGGAAAAAGAAAGGCTT 300
 ATAAAAGTCCTCAGAAACGTAAAGAGTAACTACGTATTTATAGTGTACGA
 TGCGAAACTCCAGAAACAGGAACTTTCTTCGGAACCTCTGAAATCCGTAG 400
 CGTCTTTTCGGCGGTATAGTGGTAGCAAACAGGCTGAGCAAGGAGAGGATA
 AAACAGCTCGTCCTTAAGAAGTTCAAAGAAAAAGGGATAAACGTAGAAAA 500
 CGATGCCCTTGAATACCTTCTCCAGCTCACGGGTTACAACCTGATGGAGC
 TCAAACCTTGAGGTTGAAAACTGATAGATTACGCAAGTGAAAAGAAAATT 600
 TTAACACTCGATGAGGTAAAGAGAGTAGCCTTCTCAGTCTCAGAAAACGT
 AAACGTATTTGAGTTCGTTGATTTACTCCTCTTAAAAGATTACGAAAAGG 700
 CTCTTAAAGTTTTGGACTCCCTCATTTCCTTCGGAATACACCCCTCCAG
 ATTATGAAAATCCTGTCCTCCTATGCTCTAAAACCTTACACCCTCAAGAG 800
 GCTTGAAGAGAAGGGAGAGGACCTGAATAAGGCGATGGAAAGCGTGGGAA
 TAAAGAACAACCTTCTCAAGATGAAGTTCAAATCTTACTTAAAGGCAAAC 900
 TCTAAAGAGGACTTGAAGAACCTAATCCTCTCCCTCCAGAGGATAGACGC
 TTTTCTAAACCTTACTTTTCAAGACACAGTGCAGTTGCTGGGGATTTCTT 1000
 GACCTCAAGACTGGAGAGGGAAGTTGTGAAAAATACTTCTCATGGTGGAT
 AATCTTTTTTATGAAGTTTGCGGTTTGCGTTTTTCCCGGTTCT 1093

FIG. 40

VETTIFQFQKTFFTKPPKERVFLHGEEQYLIRTFLSKLKEKYGENYTVL
 WGDEISEEEFYTALSETSIFFGSKEKAVVIYNFGDFLKKLGRKKKEKERL 100
 IKVLRNVKSNYVFIVYDAKLQKQELSSEPLKSVASF GGIVVANRLSKERI
 KQLVLKKFKEKGINVENDALEYLLQLTGYNLMELKLEVEKLIDYASEKKI 200
 LTLDEVKRVAFSVSENVNVFEFVDLLLLKDYEKALKVLDLISFGIHPLO
 IMKILSSYALKLYTLKRLEEKGEDLNKAMESVGIKNNFLKMKFKSYLKAN 300
 SKEDLKNLILSLQRIDAFSKLYFQDTVQLLRDFLTSRLEREVVKNTSHGG

FIG. 41

ATGGAAAAAGTTTTTTTGGAAAACTCCAGAAAACCTTGACATACCCGG
 AGGACTCCTTTTTTACGGCAAAGAAGGAAGCGGAAAGACGAAAACAGCTT 100
 TTGAATTTGCAAAGGTATTTTATGTAAGGAAAACGTACCTGGGGATGCG
 GAAGTTGTCCCTCCTGCAAACACGTAAACGAGCTGGAGGAAGCCTTCTTT 200
 AAAGGAGAAATAGAAGACTTTAAAGTTTATAAGACAAGGACGGTAAAAAG
 CACTTCGTTTACCTTATGGGCGAACATCCCGACTTTGTGGTAATAATCCC 300
 GAGCGGACATTACATAAAGATAGAACAGATAAGGGAAGTTAAGAACTTTG
 CCTATGTGAAGCCCGCACTAAGCAGGAGAAAAAGTAATTATAATAGACGAC 400
 GCCCACGCGATGACCTCTCAGGCGGCAAACGCTCTTTTAAAGGTATTGGA
 AGAGCCACCTGCGGACACCACCTTTATCTTGACCACGAACAGGCGTTCTG 500
 CAATCCTGCCGACTATCCTCTCCAGAACTTTTCAAGTGGAGTTCAAGGGC
 TTTTCAGTAAAAGAGGTTATGGAAATAGCGAAAGTAGACGAGGAAATAGC 600
 GAAACTCTCTGGAGGCAGTCTAAAAAGGGCTATCTTACTAAAGGAAAACA
 AAGATATCCTAAACAAAGTAAAGGAATTCTTGAAAACGAGCCGTAAAAA 700
 GTTTACAAGCTTGCAAGTGAATTCGAAAAGTGGGAACCTGAAAAGCAAAA
 ACTCTTCCTTGAAATTATGGAAGAATTGGTATCTCAAAAATTGACCGAAG 800
 AGAAAAAAGACAATTACACCTACCTTCTTGATACGATCAGACTCTTTAAA
 GACGGACTCGCAAGGGGTGTAAACGAACCTCTGTGGCTGTTTACGTTAGC 900
 CGTTCAGGCGGATTAATAAACCGTTATTGATTCCGTAACATTTAAACCTT
 AATCTAAATTATGAGAGCCTTTGAAGGAGGTCTGGTATGGAAAATTTGAA 1000
 GATTAGATATATAGATACGAGGAAGATAGGAACCGTGAGCGGTGTAAAAG
 T 1051

FIG. 42

MEKVFLEKLQKTLHIPGGLLFYKGESGKTKTAFEFAKGILCKENVPWGC
 GSCPSCKHVNELEEAFKGEIEDFKVYKDKDGKKHFVYLMGEHPDFVVI 100
 PSGHYIKIEQIREVKNFAYVKPALSRRKVIIIDDAHAMTSQAANALLKVL
 EEPPADTTFILTTNRRSAILPTILSRFQVEFKGFSVKEVMEIAKVDEEI 200
 AKLSGGSCLKRAILLKENKDI LNKVKEFLENEPLKVYKLASEFEKWEPEKQ
 KLFLEIMEELVSQKLTEKKDNYTYLLDTIRLFKDGLARGVNEPLWLFTL 300
 AVQAD

FIG. 43

ATGAACTTCCTGAAAAAGTTCCTTTTACTGAGAAAAGCTCAAAAGTCTCC	
TTACTTCGAAGAGTTCTACGAAGAAATCGATTTGAACCAGAAGGTGAAAG	100
ATGCAAGGTTTGTAGTTTGTGACTGCGAAGCCACAGAACTCGACGTAAAG	
AAGGCAAACTCCTTTCAATAGGTGCGGTTGAGGTAAAAACCTGGAAAT	200
AGACCTCTCTAAATCTTTTACGAGATACTCAAAAGTGACGAGATAAAGG	
CGGCGGAGATACATGGAATAACCAGGGAAGACGTTGAAAAGTACGGAAAG	300
GAACCAAAGGAAGTAATATACGACTTTCTGAAGTACATAAAGGGAAGCGT	
TCTCGTTGGCTACTACGTGAAGTTTGACGTCTCACTCGTTGAGAAGTACT	400
CCATAAAGTACTTCCAGTATCCAATCATCAACTACAAGTTAGACCTGTTT	
AGTTTCGTGAAGAGAGAGTACCAGAGTGGCAGGAGTCTTGACGACCTTAT	500
GAAGGAACTCGGTGTAGAAATAAGGGCAAGGCACAACGCCCTTGAAGATG	
CCTACATAACCGCTCTTCTTTTCTAAAGTACGTTTACCCGAACAGGGAG	600
TACAGACTAAAGGATCTCCCGATTTTCCTT	

FIG. 44

MNFLKKFLLLRKAQKSPYFEEFYEEIDLNQKVKDARFVVFDCEATELDVK	
KAKLLSIGAVEVKNLEIDLSKSFYEILKSDEIKAAEIHGITREDVEKYGK	100
EPKEVIYDFLKYIKGSVLVGYYVKFDVSLVEKYSIKYFQYPIINYKLDLF	
SFVKREYQSGRSLDDLMKELGVEIRARHNALEDAYITALLFLKYVYPNRE	200
YRLKDLPIFL	

FIG. 45

ATGCTCAATAAGGTTTTTATAATAGGAAGACTTACGGGTGACCCCGTTAT	
AACTTATCTACCGAGCGGAACGCCCCTAGTAGAGTTTACTCTGGCTTACA	100
ACAGAAGGTATAAAAACCAGAACGGTGAATTTTACAGAGGAAAGTCACTTC	
TTTGACGTAAAGGCGTACGGAAAAATGGCTGAAGACTGGGCTACACGCTT	200
CTCGAAAGGATACCTCGTACTCGTAGAGGGAAGACTCTCCAGGAAAAGT	
GGGAGAAAGAAGGAAAGAAGTTCTCAAAGGTCAGGATAATAGCGGAAAAC	300
GTAAGATTAATAAACAGGCCGAAAGGTGCTGAACTTCAAGCAGAAGAAGA	
GGAGGAAGTTCCTCCCATTTGAGGAGGAAATTGAAAAACTCGGTAAAGAGG	400
AAGAGAAGCCTTTTACCGATGAAGAGGACGAAATACCTTTTTTAATTTTGA	
GGAGGTTAAAGTATGGTAGTGAGAGCTCCTAAGAAGAAAGTTTGTATGTA	500
CTGTGAACAAAAGAGAGAGCCAGATT	

FIG. 46

MLNKVFIIGRLTGDPVITYLPSGTPVVEFTLAYNRRYKNQNGEFQEESHF	
FDVKAYGKMAEDWATRFSGYLVLEGRLSQEKWEKEGKKFSKVRIIAEN	100
VRLINRPGKGAELQAEIEEEVPPIEEEIEKLGKEEEKPFTDEEDEIIPF	

FIG. 47

ATGCAATTTGTGGATAAACTTCCCTGTGACGAATCCGCCGAGAGGGCGGT
TCTTGGCAGTATGCTTGAAGACCCCGAAAACATACCTCTGGTACTTGAAT 100
ACCTTAAAGAAGAAGACTTCTGCATAGACGAGCACAAGCTACTTTTCAGG
GTTCTTACAAACCTCTGGTCCGAGTACGGCAATAAGCTCGATTTTCGTATT 200
AATAAAGGATCACCTTGAAAAGAAAACTTACTCCAGAAAATACCTATAG
ACTGGCTCGAAGAACTCTACGAGGAGGCGGTATCCCCTGACACGCTTGAG 300
GAAGTCTGCAAAATAGTAAAACAACGTTCCGCACAGAGGGCGATAATTCA
ACTCGGTATAGAACTCATTACAAAAGGAAAGGAAAAACAAGACTTTTACA 400
CATTAATCGAGGAAGCCCAGAGCAGGATATTTTCCATAGCGGAAAGTGCT
ACATCTACGCAGTTTTTACCATGTGAAAGACGTTGCGGAAGAAGTTATAGA 500
ACTCATTTATAAATTCAAAGCTCTGACAGGCTAGTCACGGGACTCCCAA
GCGGTTTTACGGAACTCGATCTAAAGACGACGGGATTCCACCCTGGAGAC 600
TTAATAATACTCGCCGCAAGACCCGGTATGGGGAAAACCGCCTTTATGCT
CTCCATAATCTACAATCTCGCAAAAGACGAGGGAAAACCTCAGCTGTAT 700
TTTCCTTGGAATGAGCAAGGAACAGCTCGTTATGAGACTCCTCTCTATG
ATGTCGGAGGTCCCACTTTTCAAGATAAGGTCTGGAAGTATATCGAATGA 800
AGATTTAAAGAAGCTTGAAGCAAGCGCAATAGAAGTACGAAAGTACGACA
TATACCTCGACGACACACCCGCTCTCACTACAACGGATTTAAGGATAAGG 900
GCAAGAAAGCTCAGAAAGGAAAAGGAAGTTGAGTTCGTGGCGGTGGACTA
CTTGCAACTTCTGAGACCGCCAGTCCGAAAGAGTTCAAGACAGGAGGAAG 1000
TGGCAGAGGTTTTCAAGAACTTAAAAGCCCTTGCAAAGGAACTTCACATT
CCCGTTATGGCACTTGCGCAGCTCTCCCGTGAGGTGGAAAAGAGGAGTGA 1100
TAAAAGACCCCAGCTTGCGGACCTCAGAGAATCCGGACAGATAGAACAGG
ACGCAGACCTAATCCTTTTCCCTCCACAGACCCGAGTACTACAAGAAAAAG 1200
CCAAATCCCAGAGAGCAGGGTATAGCGGAAGTGATAATAGCCAAGCAAAG
GCAAGGACCCACGGACATTGTGAAGCTCGCATTTATTAAGGAGTACACTA 1300
AGTTTGCAAACCTAGAAGCCCTTCTGAACAACCTCCTGAAGAAGAGGAA
CTTTCCGAAATTATTGAAACACAGGAGGATGAAGGATTGAAGATATTGA 1400
CTTCTGAAAATTAAGTTTTATAATTTTATCTTGGCTATCCGGGGTAGCT
CAATCGGCAGAGCGGGTGGCTG 1472

FIG. 48

MQFVDKLPCEESAERAVLGSMLEDPENIPLVLEYLKEEDFCIDEHKLLFR
VLTNLWSEYGNKLDVFLIKDHLEKKNLLQKIPIDWLEELYEEAVSPDTLE 100
EVCKIVKQRSAQRAIIQLGITSTQFYHVKDVAEEVIELIYKFKSSDRLVT
GLPSGFTELDLKTTFHPPGDLIIILAAPGMGKTAFMLSIIYNLAKDEGKP 200
SAVFSLEMSKEQLVMRLLSMMSEVPLFKIRSGSISNEDLKKLEASAIELA
KYDIYLDLDDTPALTTDLRIRARKLRKEKEVEFVAVDYLQLLRPPVRKSSR 300
QEEVAEVSRLKALAKELHIPVMALAQLSREVEKRSKRPQLADLRESGQ
IEQDADLILFLHRPEYYKKKPNPEEQGIAEVIIAKQRQGPTDIVKLAFIK 400
EYTKFANLEALPEQPPEEEELSEIIETQEDEGFEDIDF

FIG. 49

ATGTCCTCGGACATAGACGAACTTAGACGGGAAATAGATATAGTAGACGT	
CATTTCCGAATACTTAAACTTAGAGAAGGTAGGTTCCAATTACAGAACGA	100
ACTGTCCCTTTTACCCTGACGATACACCCTCCTTTTACGTGTCTCCAAGT	
AAACAAATATTCAAGTGTTTCGGTTGCGGGGTAGGGGGAGACGCGATAAA	200
GTTTCGTTTCCCTTTACGAGGACATCTCCTATTTTGAAGCCGCCCTTGAAC	
TCGCAAAACGCTACGGAAGAAATTAGACCTTGAAAAGATATCAAAAGAC	300
GAAAAGGTATACGTGGCTCTTGACAGGGTTTGTGATTTCTACAGGGAAAG	
CCTTCTCAAAAACAGAGAGGCAAGTGAGTACGTAAAGAGTAGGGGAATAG	400
ACCCTAAAGTAGCGAGGAAGTTTGATCTTGGGTACGCACCTTCCAGTGAA	
GCACTCGTAAAAGTCTTAAAAGAGAACGATCTTTTAGAGGCTTACCTTGA	500
AACTAAAAACCTCCTTTCTCCTACGAAGGGTGTTTACAGGGATCTCTTTC	
TTCGGCGTGTCGTGATCCCGATAAAGGATCCGAGGGGAAGAGTTATAGGT	600
TTCGGTGGAAGGAGGATAGTAGAGGACAAATCTCCCAAGTACATAAACTC	
TCCAGACAGCAGGGTATTTAAAAGGGGGAGAACTTATTCGGTCTTTACG	700
AGGCAAAGGAGTATATAAAGGAAGAAGGATTTGCGATACTTGTGGAAGGG	
TACTTTGACCTTTTGAGACTTTTTTCCGAGGGAATAAGGAACGTTGTTGC	800
ACCCCTCGGTACAGCCCTGACCCAAAATCAGGCAAACCTCCTTTCCAAGT	
TCACAAAAAAGGTCTACATCCTTTACGACGGAGATGATGCGGGAAGAAAG	900
GCTATGAAAAGTGCCATTCCCCTACTCCTCAGTGCAGGAGTGGAAGTTTA	
TCCCGTTTACCTCCCCGAAGGATACGATCCCGACGAGTTTATAAAGGAAT	1000
TCCGGAAAGAGGAATTAAGAAGACTGATAAACAGCTCAGGGGAGCTCTTT	
GAAACGCTCATAAAAACCGCAAGGGAAAACCTTAGAGGAGAAAACGCGTGA	1100
GTTTCAGGTATTATCTGGGCTTTATTTCCGATGGAGTAAGGCGCTTTGCTC	
TGGCTTCGGAGTTTTCACACCAAGTACAAAGTTTCTATGGAAATTTTATTA	1200
ATGAAAATTGAAAAAATTCTCAAGAAAAAGAAATTAACTCTCCTTTAA	
GGAAAAAATCTTCTGAAAGGACTGATAGAATTAAAACCAAAAATAGACC	1300
TTGAAGTCCTGAACTTAAGTCCTGAGTTAAAGGAACTCGCAGTTAACGCC	
TTAAACGGAGAGGAGCATTTACTTCCAAAAGAAGTTCTCGAGTACCAGGT	1400
GGATAACTTGGAGAACTTTTTTAACAACATCCTTAGGGATTTACAAAAT	
CTGGGAAAAAGAGGAAGAAAAGAGGGTTGAAAAATGTAAATACTTAATTA	1500
ACTTTAATAAATTTTTTAGAGTTAGGA	

FIG. 50

MSSDIDELRREIDIVDVISEYLNLEKVGSNYRTNCPFHPDDTPSFYVSPS	
KQIFKCFGCGVGDAIKFVSLYEDISYFEAALELAKRYGKKLDLEKISKD	100
EKVYVALDRVCDIFYRESLLKNREASEYVKSREGIDPKVARKFDLGYAPSE	
ALVKVLKENDLLEAYLETKNLLSPTKGVYRDLFLRRVVIPIKDPRGRVIG	200
FGGRRIVEDKSPKYINSPDSRVFKKGENLFGLYEAKKEYIKEEGFAILVEG	
YFDLLRLFSEGIRNVVAPLGTALTQNQANLLSKFTKKVYIILYDGDDAGRK	300
AMKSAIPLLLSAGVEVYPVYLPEGYDPDEFIKEFGKEELRRLINSSGELF	
ETLIKTAARENLEEKTRFRYYLGFISDGVRRFALASEFHTKYKVPMEILL	400
MKIEKNSQEKEIKLSFKEKIFLKGLIELKPKIDLEVLNLSPELKELAVNA	
LNGEEHLLPKEVLEYQVDNLEKLFNNILRDLQKSGKKRKKRGLKNVNT	498

FIG. 51

ATGAAAAAGATTGAAAATTTGAAGTGGAAAAATGTCTCGTTTAAAAGCCT
 GGAAATAGATCCCGATGCAGGTGTGTTTCTCGTTTCCGTGGAAAAATTCT 100
 CCGAAGAGATAGAAGACCTTGTGCGTTTACTGGAGAAGAAGACGCGGTTT
 CGAGTCATCGTGAACGGTGTTCAAAAAAGTAACGGGGATCTAAGGGGAAA 200
 GATACTTTCCCTTCTCAACGGTAATGTGCCTTACATAAAAGATGTTGTTT
 TCGAAGGAAACAGGCTGATTCTGAAAGTGCTTGGAGATTTTCGCGCGGGAC 300
 AGGATCGCCTCCAAACTCAGAAGCACGAAAAAACAGCTCGATGAACTGCT
 GCCTCCCGGAACAGAGATCATGCTGGAGGTTGTGGAGCCTCCGGAAGATC 400
 TTTTGA AAAAGGAAGTACCACAACCAGAAAAGAGAGAAGAACCAAAGGCT
 GAAGAATTGAAGATCGAGGATGAAAACCACATCTTTGGACAGAAAACCCAG 500
 AAAGATCGTCTTCACCCCTCAAAAATCTTTGAGTACAACAAAAAGACAT
 CCGTGAAGGGCAAGATCTTCAAAATAGAGAAGATCGAGGGGAAAAGAACG 600
 GTCCTTCTGATTTACCTGACAGACGGAGAAGATTCTCTGATCTGCAAAGT
 CTTCAACGACGTTGAAAAGGTGGAAGGGAAAAGTATCGGTGGGAGACGTGA 700
 TCGTTGCCACAGGAGACCTCCTTCTCGAAAACGGGGAGCCCACCTTTAC
 GTGAAGGGAATCACAAAACCTCCCGAAGCGAAAAGGATGGACAAATCTCC 800
 GGTTAAGAGGGTGGAGCTCCACGCCCATAACCAAGTTCAGCGATCAGGACG
 CAATAACAGATGTGAACGAATATGTGAAACGAGCCAAGGAATGGGGCTTT 900
 CCCGCGATAGCCCTCACGGATCATGGGAACGTTTCAAGGCCATACCTTACTT
 CTACGACGCGGCGAAAGAAGCTGGAATAAAGCCCATTTTCGGTATCGAAG 1000
 CGTATCTGGTGAGTGACGTGGAGCCCGTCATAAGGAATCTCTCCGACGAT
 TCGACGTTTGGAGATGCCACGTTTCGTCTCGTCTCGACTTCGAGACGACGGG 1100
 TCTCGACCCCGCAGGTGGATGAGATCATCGAGATAGGAGCGGTGAAGATAC
 AGGGTGGCCAGATAGTGGACGAGTACCACACTCTCATAAAGCCTTCCAGG 1200
 GAGATCTCAAGAAAAAGTTCGGAGATCACCGGAATCACTCAAGAGATGCT
 GGAAAACAAGAGAAGCATCGAGGAAGTTCTGCCGGAGTTCTCGGTTTTTC 1300
 TGGAAGATTCCATCATCGTAGCACACAACGCCAACTTCGACTACAGATTT
 CTGAGGCTGTGGATCAAAAAAGTGATGGGATTTGGACTGGGAAAGACCTTA 1400
 CATAGATACGCTCGCCCTCGCAAAGTCCCTTCTCAAACCTGAGAAGCTACT
 CTCTGGATTCCGTTGTGGA AAAAGCTCGGATTGGGTCCCTTCCGGCACCCAC 1500
 AGGGCCCTGGATGACGCGAGGGTCCACCGCTCAGGTTTTCTCAGGTTTCGT
 TGAGATGATGAAGAAGATCGGTATCACGAAGCTTTTCAGAAATGGAGAAGT 1600
 TGAAGGATACGATAGACTACACCGCGTTGAAACCCTTCCACTGCACGATC
 CTCGTTTCAGAACAAAAAGGGATTGAAAAACCTATACAAACTGGTTTTCTGA 1700
 TTCCTATATAAAGTACTTCTACGGTGTTCGAGGATCCTCAAAAGTGAGC
 TCATCGAGAACAGAGAAGGACTGCTCGTGGGTAGCGCGTGTATCTCCGGT 1800
 GAGCTCGGACGTGCCGCCCTCGAAGGAGCGAGTGATTGAGAACTCGAAGA
 GATCGCGAAGTTCTACGACTACATAGAAGTCATGCCGCTCGACGTTATAG 1900
 CCGAAGATGAAGAAGACCTAGACAGAGAAAAGACTGAAAGAAGTGTACCGA
 AAAC'TCTACAGAATAGCGAAAAAATTGAACAAGTTTCGTCTCATGACCGG 2000
 TGATGTTTCAATTTCTCGATCCCGAAGATGCCAGGGGCAGAGCTGCACTTC
 TGGCACCTCAGGGAAACAGAAACTTCGAGAATCAGCCCGCACTCTACCTC 2100
 AGAACGACCGAAGAAATGCTCGAGAAGGCGATAGAGATATTCGAAGATGA
 AGAGATCGCGAGGGAAGTCGTGATAGAGAATCCCAACAGAATAGCCGATA 2200
 TGATCGAGGAAGTGCAGCCGCTCGAGAAAAAACTTCACCCGCCGATCATA
 GAGAACGCCGATGAAATAGTGAGAAACCTCACCATGAAGCGGGCGTACGA 2300
 GATCTACGGTGATCCGCTTCCCGAAATCGTCCAGAAGCGTGTGGA AAAGG

FIG. 54A

AACTGAACGCCATCATAAATCATGGATACGCCGTTCTCTATCTCATCGCT 2400
 CAGGAGCTCGTTTCAGAAATCTATGAGCGATGGTTACGTGGTTGGATCCAG
 AGGATCCGTCGGGTCTTCACTCGTGGCCAATCTCCTCGGAATAACAGAGG 2500
 TGAATCCCCCTACCACCACATTACAGGTGTCCAGAGTGCAAATACTTTGAA
 GTTGTCTGAAGACGACAGATACGGAGCGGGTTACGACCTTCCCAACAAGAA 2600
 CTGTCCAAGATGTGGGGCTCCTCTCAGAAAAGACGGCCACGGCATAACCGT
 TTGAAACGTTTCATGGGGTTCGAGGGTGACAAGGTCCCCGACATAGATCTC 2700
 AACTTCTCAGGAGAGTATCAGGAACGTGCTCATCGTTTTTGTGGAAGAACT
 CTTTCGGTAAAGACCACGTCTATAGGGCGGGAACCATAAACACCATCGCGG 2800
 AAAGAAGTGCGGTGGGTACGTGAGAAGCTACGAAGAGAAAACCGGAAAG
 AAGCTCAGAAAGGCGGAAATGGAAAGACTCGTTTTCCATGATCACGGGAGT 2900
 GAAGAGAACGACGGGTGAGCACCAGGGGGGCTCATGATCATACCGAAAG
 ACAAAGAAGTCTACGATTTCACTCCCATACAGTATCCAGCCAACGATAGA 3000
 AACGCAGGTGTGTTACACACGCACTTCGCATACGAGACGATCCATGATGA
 CCTGGTGAAGATAGATGCGCTCGGCCACGATGATCCCACTTTCATCAAGA 3100
 TGCTCAAGGACCTCACCGGAATCGATCCCATGACGATTCCCATGGATGAC
 CCGGATACGCTCGCCATATTCACTTCTGTGAAGCCTCTTGGTGTGGATCC 3200
 CGTTGAGCTGGAAAGCGATGTGGGAACGTACGGAATTCGGGAGTTCGGAA
 CCGAGTTTGTGAGGGGAATGCTCGTTGAAACGAGACCAAAGAGTTTCGCC 3300
 GAGCTTGTGAGAATCTCAGGACTGTACACGGTACGGACGTCTGGTTGAA
 CAACGCACGTGATTGGATAAACCTCGGCTACGCCAAGCTCTCCGAGGTTA 3400
 TCTCGTGTAGGGACGACATCATGAACTTCCTCATAACAAAGGAATGGAA
 CCGTCACTTGCCTTCAAGATCATGGAAAACGTGAGGAAGGGAAAGGGTAT 3500
 CACAGAAGAGATGGAGAGCGAGATGAGAAGGCTGAAGGTTCCAGAATGGT
 TCATCGAATCCTGTAAAAGGATCAAATATCTCTTCCCGAAAGCTCACGCT 3600
 GTGGCTTACGTGAGTATGGCCTTCAGAATTGCTTACTTCAAGGTTCACTA
 TCCTCTTCAGTTTTTACGCGGCGTACTTCACGATAAAAGGTGATCAGTTCG 3700
 ATCCGGTTCTCGTACTCAGGGGAAAAGAAGCCATAAAGAGGCGCTTGAGA
 GAACTCAAAGCGATGCCTGCCAAAGACGCCCAGAAGAAAAACGAAGTGAG 3800
 TGTTCCTGGAGGTTGCCCTGGAAATGATACTGAGAGGTTTTTTCCTTCCTAC
 CGCCCGACATCTTCAAATCCGACGCGAAGAAATTTCTGATAGAAGGAAAC 3900
 TCGCTGAGAATTCCGTTCAACAAACTTCCAGGACTGGGTGACAGCGTTGC
 CGAGTCGATAATCAGAGCCAGGGAAGAAAAGCCGTTCACTTCGGTGGAAG 4000
 ATCTCATGAAGAGGACCAAGGTCAACAAAAATCACATAGAGCTGATGAAA
 AGCCTGGGTGTTCTCGGGGACCTTCAGAGACGGAACAGTTCACGCTTTT
 C 4100

FIG. 54B

MKKIENLKWKNVSFKSLEIDPDAGVVLVSVEKFSEEIEDLVRLLLEKKTRF	
RVIVNGVQKSNGDLRGKILSLLNGNVPYIKDVVFEGNRLILKVLGDFARD	100
RIASKLRSTKKQLDELLPPGTEIMLEVVEPPEDLLKKEVPQPEKREEPKG	
EELKIEDENHIFGQKPRKIVFTPSKIFEYNKKTSVKGKIFKIEKIEGKRT	200
VLLIYLTGDEDSLICKVFNDVEKVEGKVSVDVIVATGDLLLLLENGETLY	
VKGITKLPEAKRMDKSPVKRVELHAHTKFSDQDAITDVNEYVKRAKEWGF	300
PAIALTDHGNVQAIPYFYDAAKEAGIKPIFGIEAYLVSDVEPVIRNLSDD	
STFGDATFVVLDFETTGLDPQVDEIIIEIGAVKIQGGQIVDEYHTLIKPSR	400
EISRKSSSEITGITQEMLNKRSEIEEVLPEFLGFLEDSIIIVAHNANFDYRF	
LRLWIKKVMGLDWERPYIDTLALAKSLLKLRSYSLDSVVEKLGGLPFRHH	500
RALDDARVTAQVFLRFVEMMKKIGITKLSEMEKLKDTIDYTALKPFHCTI	
LVQNKKGKLNLYKLVSDSYIKYFYGVPRILKSELINREGLLVGSACISG	600
ELGRAALEGASDSELEEIAKFYDYIEVMPLDVIAEDEEDLDRERLKEVYR	
KLYRIAKKLNKFVVMGTGDVHFLDPEDARGRAALLAPQGNRNFNENQPALYL	700
RTTEEMLEKAIEIFEDEEIAREVVIENPNRIADMIEEVQPLEKKLHPPII	
ENADEIVRNLTMKRAYEIIYGDPLPEIVQKRVEKELNAIINHGYAVLYLIA	800
QELVQKSMDSGYVVGSRGSSLVANLLGITEVNPLPPHYRCPECKYFE	
VVEDDRYGAGYDLPNKNCPCRGAPLRKDGHGIPFETFMGFEGDKVPDIDL	900
NFSGEYQERAHRFVEELFGKDHVYRAGTINTIAERSAVGYVRSYEEKTGK	
KLRKAEMERLVSMITGVKRTTGQHPGGLMIIPKDKEVYDFTPIQYPANDR	1000
NAGVFTTHFAYETIHDDLVKIDALGHDDPTFIKMLKDLTGIDPMTIPMDD	
PDTLAI FSSVKPLGVDPELESDVGTYGIPEFGTEFVRGMLVETRPKSFA	1100
ELVRISGLSHGTDVWLN NARDWINLGYAKLSEVISCRDDIMNFLIHKGME	
PSLAFKIMENVRKGKGITEEMESEMRLKVPEWFIESCKRIKYLFPKAHA	1200
VAYVSMAFRIAYFKVHYPLQFYAAYFTIKGDQFDPVLVLRGKEAIKRRLR	
ELKAMPAKDAQKKNEVSVLEVALEMILRGFSFLPPDIFKSDAKKFLIEGN	1300
SLRIPFNKLPGLGDSVAESIIRAREEKPFTSVEDLMKRTKVKNKHIELMK	
SLGVLGDLPETEQFTLF	1367

FIG. 55

GTGCTCGCCATGATATGGAACGACACCGTTTTTTGCGTCGTAGACACAGA	
AACCACGGGAACCGATCCCTTTGCCGGAGACCGGATAGTTGAAATAGCCG	100
CTGTTCTGTCTTCAAGGGGAAGATCTACAGAAACAAAGCGTTTCACTCT	
CTCGTGAATCCCAGAATAAGAATCCCTGCGCTGATTCAGAAAGTTCACGG	200
TATCAGCAACATGGACATCGTGGAAGCGCCAGACATGGACACAGTTTACG	
ATCTTTTCAGGGATTACGTGAAGGGAAACGGTGCTCGTGTTTCACAACGCC	300
AACTTCGACCTCACTTTTCTGGATATGATGGCAAAGGAAACGGGAACTT	
TCCAATAACGAATCCCTACATCGACACACTCGATCTTTCAGAAGAGATCT	400
TTGGAAGGCCTCATTCTCTCAAATGGCTCTCCGAAAGACTTGGAATAAAA	
ACCACGATACGGCACCGTGCTCTTCCAGATGCCCTGGTGACCGCAAGAGT	500
TTTTGTGAAGCTTGTTGAATTTCTTGGTGAAAACAGGGTCAACGAATTCA	
TACGTGGAAAACGGGGG	567

FIG. 56

MLAMIWNDTVFCVVDTEETTGTDPFAGDRIVEIAAVPVFKGKIYRNKAFHS	
LVNPRIRIPALIQKVHGISNMDIVEAPDMDTVYDLFRDYVKGTVLVFHNA	100
NFDLTFLDMMAKETGNFPITNPYIDTLDLSEEIFGRPHSLKWLSERLGIK	
TTIRHRALPDALVTARVFVKLVEFLGENRVNEFIRGKRG	189

FIG. 57

GTGGAAGTTCTTTACAGGAAGTACAGGCCAAAGACTTTTTCTGAGGTTGT	
CAATCAGGATCATGTGAAGAAGGCAATAATCGGTGCTATTTCAGAAGAACA	100
GCGTGGCCACCGGATACATATTCGCCGGTCCGAGGGGAACGGGGAAGACT	
ACTCTTGCCAGAATTCTCGCAAATCCCTGAACTGTGAGAACAGAAAGGG	200
AGTTGAACCCTGCAATTCTGCAGAGCCTGCAGAGAGATAGACGAGGGAA	
CCTTCATGGACGTGATAGAGCTCGACGCGGCCTCCAACAGAGGAATAGAC	300
GAGATCAGAAGAATCAGAGACGCCGTTGGATACAGGCCGATGGAAGGTAA	
ATACAAAGTCTACATAATAGACGAAGTTCACATGCTCACGAAAGAAGCCT	400
TCAACGCGCTCCTCAAAACACTCGAAGAACCTCCTTCCCACGTCGTGTTT	
GTGCTGGCAACGACAAACCTTGAGAAGGTTCTTCCCACGATTATCTCGAG	500
ATGTCAGGTTTTTCGAGTTCAGAAACATTCCCGACGAGCTCATCGAAAAGA	
GGCTCCAGGAAGTTGCGGAGGCTGAAGGAATAGAGATAGACAGGGAAAGCT	600
CTGAGCTTCATCGCAAAAAGAGCCTCTGGAGGCTTGAGAGACGCGCTCAC	
CATGCTCGAGCAGGTGTGGAAGTTCTCGGAAGGAAAGATAGATCTCGAGA	700
CGGTACACAGGGCGCTCGGGTTGATACCGATACAGGTTGTTTCGCGATTAC	
GTGAACGCTATCTTTTCTGGTGATGTGAAAAGGGTCTTCACCGTTCTCGA	800
CGACGTCTATTACAGCGGGAAGGACTACGAGGTGCTCATTTCAGGAAGCAG	
TCGAGGATCTGGTCGAAGACCTGGAAAGGGAGAGAGGGGTTTACCAGGTT	900
TCAGCGAACGATATAGTTTCAGGTTTCGAGACAACTTCTGAATCTTCTGAG	
AGAGATAAAGTTTCGCCGAAGAAAAACGACTCGTCTGTAAAGTGGGTTTCGG	1000
CTTACATAGCGACGAGGTTCTCCACCACAAACGTTTCAGGAAAACGATGTC	
AGAGAAAAAACGATAATTCAAATGTACAGCAGAAAAGAGAAGAAAGA	1100
AACGGTGAAGGCAAAAGAAGAAAAACAGGAAGACAGCGAGTTCGAGAAAC	
GCTTCAAAGAACTCATGGAAGAACTGAAAGAAAAGGGCGATCTCTCTATC	1200
TTTGTGCTCTCAGCCTCTCAGAGGTGCAGTTTGACGGAGAAAAGGTGAT	
TATTTCTTTTGATTTCATCGAAAGCTATGCATTACGAGTTGATGAAGAAAA	1300
AACTGCCTGAGCTGGAAGAACATTTTTTCTAGAAAACTCGGGAAAAAAGTA	
GAAGTTGAACTTCGACTGATGGGAAAAGAAGAAACAATCGAGAAGGTTTC	1400
TCAGAAGATCCTGAGATTGTTTGAACAGGAGGGA	

FIG. 58

MEVLYRKYPKTFSEVVNQDHVKKAIIGAIQKNSVAHGYIFAGPRGTGKT	
TLARILAKSLNCENRKGVEPCNSCRACREIDEGTFMDVIELDAASNRGID	100
EIRRIRDAVGYPMEGKYKVYIIDEVHMLTKEAFNALLKTLEPPSHVVF	
VLATTNLEKVPPTIISRCQVFEFRNIPDELIEKRLQEVAAEAGIEIDREA	200
LSFIAKRASGGLRDALTMLEQVWKFSEKIDLETVHRALGLIPIQVVRDY	
VNAIFSGDVKRVFTVLDDVYYSGKDYEVLIQEAVEDLVEDLERERGVYQV	300
SANDIVQVSRQLLNLLREIKFAEEKRLVCKVGSAYIATRFSTTNVQENDV	
REKNDNSNVQQKEEKETVKAKEEKQEDSEFEKRFKELMEELKEKGDLSI	400
FVALSLSEVQFDGEKVIISFDSSKAMHYELMKKKLPELENIFSRKLGGKV	
EVELRLMGKEETIEKVSQKILRLFEQEG	478

FIG. 59

ATGAAAGTAACCGTCACGACTCTTGAATTGAAAGACAAAATAACCATCGC	
CTCAAAAGCGCTCGCAAAGAAATCCGTGAAACCCATTCTTGCTGGATTTC	100
TTTTCGAAGTGAAAGATGGAAATTTCTACATCTGCGCGACCGATCTCGAG	
ACCGGAGTCAAAGCAACCGTGAATGCCGCTGAAATCTCCGGTGAGGCACG	200
TTTTGTGGTACCAGGAGATGTCATTGAGAGATGGTCAAGGTTCTCCAG	
ATGAGATAACGGAACCTTTCTTTAGAGGGGGATGCTCTTGTTATAAGTTCT	300
GGAAGCACCGTTTTTCAGGATCACCACCATGCCCGCGGACGAATTTCCAGA	
GATAACGCCTGCCGAGTCTGGAATAACCTTCGAAGTTGACACTTCGCTCC	400
TCGAGGAAATGGTTGAAAAGGTCATCTTCGCCGCTGCCAAAGACGAGTTC	
ATGCGAAATCTGAATGGAGTTTTCTGGGAACTCCACAAGAATCTTCTCAG	500
GCTGGTTGCAAGTGATGGTTTCAGACTTGCACTTGCTGAAGAGCAGATAG	
AAAACGAGGAAGAGGCGAGTTTCTTGCTCTCTTTGAAGAGCATGAAAGAA	600
GTTCAAAACGTGCTGGACAACACAACGGAGCCGACTATAACGGTGAGGTA	
CGATGGAAGAAGGGTTTTCTCTGTCGACAAATGATGTAGAAACGGTGATGA	700
GAGTGGTCGACGCTGAATTTCCCGATTACAAAAGGGTGATCCCCGAAACT	
TTCAAAACGAAAGTGTTGGTTTCCAGAAAAGAACTCAGGGAATCTTTGAA	800
GAGGGTGATGGTGATTGCCAGCAAGGGAAGCGAGTCCGTGAAGTTCGAAA	
TAGAAGAAAACGTTATGAGACTTGTGAGCAAGAGCCCGGATTATGGAGAA	900
GTGGTCGATGAAGTTGAAGTTCAAAAAGAAGGGGAAGATCTCGTGATCGC	
TTTCAACCCGAAGTTCATCGAGGACGTTTTGAAGCACATTGAGACTGAAG	1000
AAATCGAAATGAACTTCGTTGATTCTACCAGTCCATGTCAGATAAATCCA	
CTCGATATTTCTGGATACCTTTACATAGTGATGCCCATCAGACTGGCA	1098

FIG. 60

MKVTVTTLLELKDKITIASKALAKKSVKPILAGFLFEVKDGNFYICATDLE	
TGVKATVNAAEISGEARFVVPGDVVIQKMKVLPDEITELSLGDALVISS	100
GSTVFRITTMPADEFPEITPAESGITFEVDTSLLLEEMVEKVIFAAAKDEF	
MRNLNGVFWELHKNLLRLVASDGFRLALAEQIENEEASFLSLKSMKE	200
VQNVLDNTTEPTITVRYDGRRVSLSTNDVETVMRVVDAEFDPYKRVIPET	
FKTKVVVSRKELRESLKRVMVIASKGSESVKFEIEENVMLVSKSPDYGE	300
VVDEVEVQKEGEDLVIAFNPKFIEDVLKHIEETEEIEMNFVDSTSPCQINP	
LDISGYLYIVMPIRLA	366

FIG. 61

ATGCCAGTCACGTTTCTCACAGGTACTGCAGAACTCAGAAGGAAGAATT	
GATAAAGAACTCCTGAAGGATGGTAACGTGGAGTACATAAGGATCCATC	100
CGGAGGATCCCGACAAGATCGATTTTCATAAGGTCTTTACTCAGGACAAAG	
ACGATCTTTTCCAACAAGACGATCATTTGACATCGTCAATTTTCGATGAGTG	200
GAAAGCACAGGAGCAGAAGCGTCTCGTTGAACTTTTGAAAAACGTACCGG	
AAGACGTTTCATATCTTCATCCGTTCTCAAAAAACAGGTGGAAAGGGAGTA	300
GCGCTGGAGCTTCCGAAGCCATGGGAAACGGACAAGTGGCTTGAGTGGAT	
AGAAAAGCGCTTCAGGGAGAATGGTTTGCTCATCGATAAAGATGCCCTTC	400
AGCTGTTTTTCTCCAAGGTTGGAACGAACGACCTGATCATAGAAAGGGAG	
ATTGAAAAACTGAAAGCTTATTCAGAGGACAGAAAGATAACGGTAGAAGA	500
CGTGGAAGAGGTCGTTTTTACCTATCAGACTCCGGGATACGATGATTTTT	
GCTTTGCTGTTTTCCGAAGGAAAAAGGAAGCTCGCTCACTCTCTTCTGTCG	600
CAGCTGTGGAACACAGAGTCCGTGGTGATTGCCACTGTCCTTGCGAA	
TCACTTCTTGATCTCTTCAAATCCTCGTTCTTGTGACAAAGAAAAGAT	700
ACTACACCTGGCCTGATGTGTCCAGGGTGTCCAAAGAGCTGGGAATTCCC	
GTTCCCTCGTGTGGCTCGTTTCCTCGGTTTCTCCTTTAAGACCTGGAAATT	800
CAAGGTGATGAACCACCTCCTCTACTACGATGTGAAGAAGGTTAGAAAGA	
TACTGAGGGATCTCTACGATCTGGACAGAGCCGTGAAAAGCGAAGAAGAT	900
CCAAAACCGTTCTTCCACGAGTTCATAGAAGAGGTGGCACTGGATGTATA	
TTCTCTTCAGAGAGATGAAGAA	972

FIG. 62

MPVTFLTGTAEQKEELIKKLLKDG NVEYIRIHPEDDPKIDFIRSLLR TK	
TIFSNKTIIDIVNFDEWKAQEQR LVELLKNVPEDVHIFIRSQKTGGKGV	100
ALELPKPWETDKWLEWIEKRFR ENGLLIDKDALQLFFSKVGTNDLI IERE	
IEKLKAYSEDRKITVEDVEEVVFTYQTPGYDDFCFAVSEGKRKL AHSLLS	200
QLWKTTESVVIATVLANHF LDFKILVLVTKKRYYTWPDVSRVSKELGIP	
VPRVARFLGFSFKTWKFKVMNHL LYYDVKKVRKILRDLYDLDR AVKSEED	300
PKPFFHEFIEEVALDVYSLQR DEE	

FIG. 63

ATGAACGATTTGATCAGAAAAGTACGCTAAAGATCAACTGGAAACTTTGAA	
AAGGATCATAGAAAAGTCTGAAGGAATATCCATCCTCATAAATGGAGAAG	100
ATCTCTCGTATCCGAGAGAAGTATCCCTTGAAC TTCCCGAGTACGTGGAG	
AAATTTCCCCCGAAGGCCTCGGATGTTCTGGAGATAGATCCCGAGGGGGA	200
GAACATAGGCATAGACGACATCAGAACGATAAAGGACTTCCTGAACTACA	
GCCCCGAGCTCTACACGAGAAAAGTACGTGATAGTCCACGACTGTGAAAGA	300
ATGACCCAGCAGGCGGCGAACGCGTTTCTGAAGGCCCTTGAAGAACCACC	
AGAATACGCTGTGATCGTTCTGAACACTCGCCGCTGGCATTATCTACTGC	400
CGACGATAAAGAGCCGAGTGTTTCAGAGTGTTGTGAACGTTCCAAAGGAG	
TTCAGAGATCTCGTGAAAGAGAAAATAGGAGATCTCTGGGAGGAACTTCC	500
ACTTCTTGAGAGAGACTTCAAAACGGCTCTCGAAGCCTACAAACTTGGTG	
CGGAAAAA CTTTCTGGATTGATGGAAAGTCTCAAAGTTTTTGGAGACGGAA	600
AAACTCTTGAAAAAGGTCTTTTCAAAGGCCTCGAAGGTTATCTCGCATG	
TAGGGAGCTCCTGGAGAGATTTTCAAAGGTGGAATCGAAGGAATTCTTTG	700
CGCTTTTTTGATCAGGTGACTAACACGATAACAGGAAAAGACGCGTTTCTT	
TTGATCCAGAGACTGACAAGAATCATTTCTCCACGAAAACACATGGGAAAG	800
CGTTGAAGATCAAAAAAGCGTGTCTTTCTCGATTCAATTCTCAGGGTGA	
AGATAGCGAATCTGAACAACAACTCACTCTGATGAACATCCTCGCGATA	900
CACAGAGAGAGAAAGAGAGGTGTCAACGCTTGGAGC	

FIG. 64

MNDLIRKYAKDQLETLKRIIEKSEGISILINGEDLSYPREVSLELPEYVE	
KFPPKASDVLEIDPEGENIGIDDIRTIKDFLNYSPELYTRKYVIVHDCER	100
MTQQAANAFLKALEEPPEYAVIVLNTRRWYLLPTIKSRVFRVVVNPKE	
FRDLVKEKIGDLWEELPLLERDFKTALEAYKLGAEKLSGLMESLKVLETE	200
KLLKKVLSKGLEGYLACRELLERFSKVESKEFFALFDQVTNTITGKDAFL	
LIQRLTRIILHENTWESVEDKSVSFLDSILRVKIANLNNKLTLMNILAIH	300
RERKRGVNAWS	

FIG. 65

ATGCGTGTTCCCCCGCACAACTTAGAGGCCGAAGTTGCTGTGCTCGGAAG
CATATTGATAGATCCGTCGGTAATAAACGACGTTCTTGAAATTTTGAGCC 100
ACGAAGATTTCTATCTGAAAAAACACCAACACATCTTCAGAGCGATGGAA
GAGCTTTACGACGAAGGAAAACCGGTGGACGTGGTTTCCGTCTGTGACAA 200
GCTTCAAAGCATGGGAAAACCTCGAGGAAGTAGGTGGAGATCTGGAAGTGG
CCCAGCTCGCTGAGGCTGTGCCCAGTTCTGCACACGCACTTCACTACGCG
GAGATCGTCAAGGAAAAATCCATTCTGAGGAAACTCATTGAGATCTCCAG
AAAAATCTCAGAAAGTGCCTACATGGAAGAAGATGTGGAGATCCTGCTCG 400
ACAACGCAGAAAAGATGATCTTCGAGATCTCAGAGATGAAAACGACAAAA
TCCTACGATCATCTGAGAGGCATCATGCACCGGGTGTTTGAAAACCTGGA
GAACTTCAGGGAAAGAGCCAACCTTATAGAACCCGGTGTGCTCATAACGG
GACTACCAACGGGATTCAAAGTCTGGACAAACAGACCACAGGGTTCCAC 600
AGCTCCGATCTGGTGATAATAGCAGCGAGACCCTCCATGGGAAAAACCTC
CTTCGCACTCTCAATAGCGAGGAACATGGCTGTCAATTTTCGAAATCCCCG
TCGGAATATTCAGTCTCGAGATGTCCAAGGAACAGCTCGCTCAAAGACTA
CTCAGCATGGAGTCCGGTGTGGATCTTTACAGCATCAGAACAGGATACCT 800
GGATCAGGAGAAGTGGGAAAGACTCACAATAGCGGCTTCTAAACTCTACA
AAGCACCCATAGTTGTGGACGATGAGTCACTCCTCGATCCGCGATCGTTG
AGGGCAAAAGCGAGAAGGATGAAAAAGAATACGATGTAAAAGCCATTTT 900
TGTCGACTATCTCCAGCTCATGCACCTGAAAGGAAGAAAAGAAAGCAGAC 1000
AGCAGGAGATATCCGAGATCTCGAGATCTCTGAAGCTCCTTGCGAGGGAA
CTCGACATAGTGGTGATAGCGCTTTCACAGCTTTCGAGGGCCGTAGAACA 1100
GAGAGAAGACAAAAAGACCGAGGCTGAGTGACCTCAGGGAATCCGGTGCGA
TAGAACAGGACGCAGACACAGTCATCTTCATCTACAGGGAGGAATATTAC 1200
AGGAGCAAAAAATCCAAGAGGAAAGCAAGCTTCACGAACCTCACGAAGC
TGAAATCATAATAGGTAAACAGAGAAACGGTCCCGTTGGAACGATCACTC 1300
TGATCTTCGACCCCAGAACGGTTACGTTCCATGAAGTCGATGTGGTGCAT
TCA 1353

FIG. 68

MRVPPHNLEAEVAVLGSILIDPSVINDVLEILSHEDFYLKKHQHIFRAME
ELYDEGKPVVDVSVCDKLQSMGKLEEVGGDLEVAQLAEAVPSSAHALHYA 100
EIVKEKSILRKLIEISRKISESAYMEEDVEILLDNAEKMIFEISEMKTTK
SYDHLRGIMHRVFENLENFRERANLIEPGVLITGLPTGFKSLDKQTTGFH 200
SSDLVIAARPSMGKTSFALSIARNMAVNFEIPVGIFSLEMSKEQLAQRL
LSMESGVDLYSIRTGYLDQEKWERLTIAASKLYKAPIVVDDESLLDPRSL 300
RAKARRMKKEYDVKAIFVDYLQLMHLKGRKESRQQEISEISRSLKLLARE
LDIVVIALSQLSRAVEQREDKRPRLSDLRESGAIEQDADTVIFIYREEYY 400
RSKKSKEESKLHEPHEAEIIIGQRNGPVGTTITLIFDPRTVTFHEVDVH
S 451

FIG. 69

GTGATTCCTCGAGAGGTCATCGAGGAAATAAAAAGAAAAGGTTGACATCGT	
AGAGGTCATTTCCGAGTACGTGAATCTTACCCGGGTAGGTTCTCTCTACA	100
GGGCTCTCTGTCCCTTTTCATTGAGAAACCAATCCTTCTTTCTACGTTTAT	
CCGGGTTTGAAGATATACCATTTGTTTCGGCTGCGGTGCGAGTGGAGACGT	200
CATCAAATTTCTTCAAGAAATGGAAGGGATCAGTTTCCAGGAAGCGCTGG	
AAAGACTTGCCAAAAGAGCTGGGATTGATCTTTCTCTCTACAGAACAGAA	300
GGGACTTCTGAATACGGAAAATACATTCGTTTGTACGAAGAAACGTGGAA	
AAGGTACGTCAAAGAGCTGGAGAAATCGAAAGAGGCAAAAGACTATTTAA	400
AAAGCAGAGGCTTCTCTGAAGAAGATATAGCAAAGTTTCGGCTTTGGGTAC	
GTCCCAAGAGATCCAGCATCTCTATAGAAGTTGCAGAAGGCATGAACAT	500
AACACTGGAAGAACTTGTGAGATACGGTATCGCGCTGAAAAGGGTGATC	
GATTCGTTGATAGATTGGAAGGAAGAATCGTTGTTCCAATAAAGAACGAC	600
AGTGGTCATATTGTGGCTTTTGGTGGGCGTGCTCTCGGCAACGAAGAACC	
GAAGTATTTGAACTCTCCAGAGACCAGGTATTTTTCGAAGAAGAAGACCC	700
TTTTTCTCTTCGATGAGGCGAAAAAAGTGGCAAAAGAGGTTGGTTTTTTC	
GTCATCACCGAAGGCTACTTCGACGCGCTCGCAATTCAGAAAGGATGGAAT	800
ACCAACGGCGGTGCTGTTCTTGGGGCGAGTCTTTCAAGAGAGGCGATTCT	
TAAAACTTTCGGCGTATTGAAAAACGTCATACTGTGTTTCGATAATGAC	900
AAAGCAGGCTTCAGAGCCACTCTCAAATCCCTCGAGGATCTCCTAGACTA	
CGAATTCAACGTGCTTGTGGCAACCCCTCTCCTTACAAAGACCCAGATG	1000
AACTCTTTCAGAAAGAAGGAGAAGGTTTATTGAAAAAGATGCTGAAAAAC	
TCGCGTTTCGTTTGAATATTTTCTGGTGACGGCTGGTGAGGTCTTCTTTGA	1100
CAGGAACAGCCCCGCGGGTGTGAGATCCTACCTTTCTTTCCTCAAAGGTT	
GGGTCCAAAAGATGAGAAGGAAAGGATATTTGAAACACATAGAAAATCTC	1200
GTGAATGAGGTTTCATCTTCTCTCCAGATACCAGAAAACCAGATTTTGAA	
CTTTTTTTGAAAGCGACAGGTCTAACACTATGCCTGTTTATGAGACCAAGT	1300
CGTCAAAGGTTTACGATGAGGGGAGAGGACTGGCTTATTTGTTTTTTGAAC	
TACGAGGATTTGAGGGAAAAGATTCTGGAACCTGGACTTAGAGGTACTGGA	1400
AGATAAAAACGCGAGGGAGTTTTTCAAGAGAGTCTCACTGGGAGAAGATT	
TGAACAAAGTCATAGAAAACCTCCCAAAGAGCTGAAAGACTGGATTTTTT	1500
GAGACAATAGAAAGCATTCTCCTCCAAAGGATCCCGAGAAATTCCTCGG	
TGACCTCTCCGAAAAGTTGAAAATCCGACGGATAGAGAGACGTATCGCAG	1600
AAATAGATGATATGATAAAGAAAGCTTCAAACGATGAAGAAAGGCGTCTT	
CTTCTCTCTATGAAAGTGGATCTCCTCAGAAAAATAAAGAGGAGG	1695

FIG. 70

MIPREVIEEIKEKVDIVEVISEYVNLTRVGSSYRALCPFHSETNPSFYVH	
PGLKIYHCFGCGASGDVIKFLQEMEGISFQEALERLAKRAGIDLSLYRTE	100
GTSEYGKYIRLYEETWKRYVKELEKSKEAKDYLSRGFSEEDIAKFGFGY	
VPKRSSISIEVAEGMNITLEELVRYGIALKKGDRFVDRFEGRIVVPIKND	200
SGHIVAFGGRALGNEEPKYLNSPETRYFSKKKTLFLFDEAKKVAKEVGFF	
VITEGYFDALAFRKDGIPTAVAVLGASLSREAILKLSAYSKNVILCFDND	300
KAGFRATLKSLEDLLDYEFNVLVATPSPYKDPDELFOKEGEGSLKKMLKN	
SRSFEYFLVTAGEVFFDRNSPAGVRSYLSFLKGWVQKMRRKGYLKHIEENL	400
VNEVSSSLQIPENQILNFFESDRSNTMPVHETKSSKVYDEGRGLAYLFLN	
YEDLREKILELDLEVLEDKNAREFFKRVSLGEDLNKVIENFPKELKDWIF	500
ETIESIPPPKDPEKFLGDLSEKLIKIRRIERRIAEIDDMIKKASNDEERRL	
LLSMKVDLLRKIKRR	565

FIG. 71

ATGGCTCTACACCCGGCTCACCTGGGGCAATAATCGGGCACGAGGCCGT	
TCTCGCCCTCCTTCCCCGCCTCACCGCCCAGACCCTGCTCTTCTCCGGCC	100
CCGAGGGGGTGGGGCGGCGCACCGTGGCCCGCTGGTACGCCTGGGGGCTC	
AACCGCGGCTTCCCCCGCCCTCCCTGGGGGAGCACCCGGACGTCTCGA	200
GGTGGGGCCCAAGGCCCGGGACCTCCGGGGCCGGGCCGAGGTGCGGCTGG	
AGGAGGTGGCGCCCCTCTTGGAGTGGTGCTCCAGCCACCCCCGGGAGCGG	300
GTGAAGGTGGCCATCCTGGACTCGGCCACCTCCTCACCGAGGCCGCCGC	
CAACGCCCTCCTCAAGCTCCTGGAGGAGCCCCCTTCTACGCCCCGCATCG	400
TCCTCATCGCCCCAAGCCGCGCCACCCTCCTCCCCACCCTGGCCTCCCGG	
GCCACGGAGGTGGCATTGCCCCCGTGCCCGAGGAGGCCCTGCGCGCCCT	500
CACCCAGGACCCGGAGCTCCTCCGCTACGCCGCGGGGCCCGGGCCGCC	
TCCTTAGGGCCCTCCAGGACCCGGAGGGGTACCGGGCCCGCATGGCCAGG	600
GCGCAAAGGGTCTTGAAAGCCCCGCCCCTGGAGCGCCTCGCTTTGCTTCG	
GGAGCTTTTGGCCGAGGAGGAGGGGTCCACGCCCTCCACGCCGTCTTAA	700
AGCGCCCGGAGCACCTCCTTGCCCTGGAGCGGGCGCGGGAGGCCCTGGAG	
GGGTACGTGAGCCCCGAGCTGGTCCTCGCCCGGCTGGCCTTAGACTTAGA	800
GACA	

FIG. 72

MALHPAHPGAIIGHEAVLALLPRLTAQTLLFSGPEGVGRRTVARWYAWGL	
NRGFPPPSLGEHPDVLEVGPKARDLRGRAEVRLEEVAPLLEWCSSHPRER	100
VKVAILDSAHLLTEAAANALLKLLLEPPSYARIVLIAPSRATLLPTLASR	
ATEVAFAPVPEEALRALTDPELLRYAAGAPGRLLRALQDPEGYRARMAR	200
AQRVLKAPPLERLALLRELLAEEEGVHALHAVLKRPEHLLALERAREALE	
GYVSPELVLARLALDLET	268

FIG. 73

ATGCTGGACCTGAGGGAGGTGGGGGAGGCGGAGTGGAAGGCCCTAAAGCC	
CCTTTTGGAAAGCGTGCCCGAGGGCGTCCCGTCCTCCTCCTGGACCCTA	100
AGCCAAGCCCCTCCCGGGCGGCCTTCTACCGGAACCGGGAAAGGCGGGAC	
TTCCCCACCCCCAAGGGGAAGGACCTGGTGCGGCACCTGGAAAACCGGGC	200
CAAGCGCCTGGGGCTCAGGCTCCCGGGCGGGGTGGCCCAGTACCTGGCCT	
CCCTGGAGGGGGACCTCGAGGCCCTGGAGCGGGAGCTGGAGAAGCTTGCC	300
CTCCTCTCCCCACCCCTCACCTGGAGAAGGTGGAGAAGGTGGTGGCCCT	
GAGGCCCCCCTCACGGGCTTTGACCTGGTGCGCTCCGTCTGGAGAAGG	400
ACCCAAGGAGGCCCTCCTGCGCCTAGGCGGCCTCAAGGAGGAGGGGGAG	
GAGCCCCCTCAGGCTCCTCGGGGCCCTCTCCTGGCAGTTCGCCCTCCTCGC	500
CCGGGCCTTCTTCTCCTCCGGGAAAACCCAGGCCCAAGGAGGAGGACC	
TCGCCCCCTCGAGGCCACCCCTACGCCGCCCGCCGCGCCCTGGAGGCG	600
GCGAAGCGCCTCACGGAAGAGGCCCTCAAGGAGGCCCTGGACGCCCTCAT	
GGAGGCGGAAAAGAGGGCCAAGGGGGGAAAGACCCGTGGCTCGCCCTGG	700
AGGCGGCGGTCTCCGCCTCGCCCCTTGA	

FIG. 74

MVIAFTGDPFLAREALLEEARLRGLSRFTEPTPEALAQALAPGLFGGGGA	
MLDLREVGEAEWKALKPLLESVPEGVPVLLDPKPSPSRAAFYRNRERRD	100
FPTPKGKDLVRHLENRAKRLGLRLPGGVAQYLASLEGDLEALERELEKLA	
LLSPPLTLEKVEKVVALRPPLTGFDLVRVLEKDPKEALLRLGGLKEEGE	200
EPLRLLGALSWQFALLARAFFLLRENPRPKEEDLARLEAHPYAARRALEA	
AKRLTEEALKEALDALMEAEKRAKGGKDPWLALEAAVLRRLAR	292

FIG. 75

ATGGCTCGAGGCCTGAACCGCGTTTTCTCATCGGCGCCCTCGCCACCCG	
GCCGGACATGCGCTACACCCCGGCGGGGCTCGCCATTTTGACCTGACCC	100
TCGCCGGTCAGGACCTGCTTCTTTCCGATAACGGGGGGGAACCGGAGGTG	
TCCTGGTACCACCGGTGAGGCTCTTAGGCCGCCAGGCGGAGATGTGGGG	200
CGACCTCTTGACCAAGGGCAGCTCGTCTTCGTGGAGGGCCGCCTGGAGT	
ACCGCCAGTGGGAAAGGGAGGGGGAGAAGCGGAGCGAGCTCCAGATCCGG	300
GCCGACTTCCGGACCCCCTGGACGACCGGGGGAAGAAGCGGGCGGAGGAC	
AGCCGGGGCCAGCCCAGGCTCCGCGCCGCCCTGAACCAGGTCTTCCTCAT	400
GGGCAACCTGACCCGGGACCCGGAACCTCCGCTACACCCCCCAGGGCACCG	
CGGTGGCCCCGGCTGGGCCTGGCGGTGAACGAGCGCCGCCAGGGGGCGGAG	500
GAGCGCACCCACTTCGTGGAGGTTCAAGCCTGGCGCGACCTGGCGGAGTG	
GGCCGCCGAGCTGAGGAAGGGCGACGGCCTTTTCGTGATCGGCAGGTTGG	600
TGAACGACTCCTGGACCAGCTCCAGCGGCGAGCGGCGCTTCCAGACCCGT	
GTGGAGGCCCTCAGGCTGGAGCGCCCCACCCGTGGACCTGCCAGGCCCTG	700
CCCAGGCCGGCGGAACAGGTCCCGCGAAGTCCAGACGGGTGGGGTGGACA	
TTGACGAAGGCTTGGAAGACTTTCCGCCGGAGGAGGATTTGCCGTTTTGA	800
GCACGAA	

FIG. 76

MARGLNRVFLIGALATRPDMRYTPAGLAILDLTLAQDLLLLSDNGGEPEV	
SWYHRVRLLRQAEMWGDLLDQGQLVFVEGRLEYRQWEREGEKRSSELQIR	100
ADFLDPLDDRGGKKRAEDSRGQPRRLRAALNQVFLMGNLTRDPELRYTPQGT	
AVARLGLAVNERRQGAEERTHFVEVQAWRDLAEWAAELRKGDGLFVIGRL	200
VNDSWTSSSGERRFQTRVEALRLERPTRGPAQACPGRNRNSREVQTTGGVD	
IDEGLEDFFPEEDLPF	266

FIG. 77

AATTCCGACATTTCAATTGAATCGTTTATTCCGCTTGAAAAAGAAGGCAA	
GTTGCTCGTTGATGTGAAAAGACCGGGGAGCATCGTACTGCAGGCGCGCT	100
TTTTCTCTGAAATCGTGAAAAAAGTCCGCAACAAACGGTGGAATCGAA	
ACGGAAGACAACCTTTTGGACGATCATCCGCTCGGGGCACTCAGAATTCGG	200
CCTCAATGGGCTAAACGCCGACGAATATCCGCGCCTGCCGCAAATTGAAG	
AAGAAAACGTGTTTCAAATCCCGGCTGATTTATTGAAAACCGTGATTCGG	300
CAAACGGTGTTTCGCCGTTTCTACATCGGAAACGCGCCCAATCTTGACAGG	
TGTCAACTGGAAAGTTGAACATGGCGAGCTTGTCTGCACAGCGACCGACA	400
GTCATCGCTTAGCCATGCGCAAAGTGAAAATTGAGTCGGAAAATGAAGTA	
TCATACAACGTCGTCATCCCTGGAAAAAGTCTTAATGAGCTCAGCAAAAT	500
TTTGGATGACGGCAACCACCCGGTGGACATCGTCATGACAGCCAATCAAG	
TGCTATTTAAGGCCGAGCACCTTCTCTTCTTTTCCCGGCTGCTTGACGGC	600
AACTATCCGGAGACGGCCCGCTTGATTCCAACAGAAAGCAAAACGACCAT	
GATCGTCAATGCAAAGAGTTTCTGCAGGCAATCGACCGAGCGTCCTTGC	700
TTGCTCGAGAAGGAAGGAACAACGTTGTGAAACTGACGACGCTTCCTGGA	
GGAATGCTCGAAATTTCTTCGATTTCTCCGAGATCGGGAAAGTGACGGAG	800
CAGCTGCAAACGGAGTCTCTTGAAGGGGAAGAGTTGAACATTTTCGTTTCA	
CGCGAAATATATGATGGACGCGTTGCGGGCGCTTGATGGAACAGACATTT	900
CAAATCAGCTTCACTGGGGCCATGCGGGCCGTTTCTGTTGCGCCCGCTTCA	
ACCGATTGATGCTTCAGCTCATTTTGCCGGTGAGAACATAT	992

FIG. 78

NSDISIIIESFIPLEKEGKLLVDVKRPGSIVLQARFFSEIVKKLPQQTVEI	
ETEDNFLTIIRSGHSEFRLNGLNADEYPRLPQIEEENVFQIPADLLKTVI	100
RQTVFAVSTSETRPILTGVDNWKVEHGELVCTATDSHRLAMRKVKIIESEN	
EVSYNVVI PGKSLNELSKIILDDGNHPVDIVMTANQVLFKAEHLLFFSRL	200
LDGNYPETARLIPTESKTTMIVNAKEFLQAI DRASLLAREGRNNVVKLTT	
LPGGMLEISSISPEIGKVTEQLQTESLEGEELNISFS AKYMMDALRALDG	300
TDIQISFTGAMRPFLLRPLHTDSMLQLILPVRTY	

FIG. 79

ATGATTAACCGCGTCATTTTGGTCGGCAGGTAAACGAGAGATCCGGAGTT	
GCGTTACACTCCAAGCGGAGTGGCTGTTGCCACGTTTACGCTCGCGGTCA	100
ACCGTCCGTTTACAAATCAGCAGGGCGAGCGGAAACGGATTTTATTCAA	
TGTGTCGTTTGGCGCCGCCAGGCGGAAAACGTCGCCAACTTTTGA AAAA	200
GGGGAGCTTGGCTGGTGTGATGGCCGACTGCAAACCCGCAGCTATGAAA	
ATCAAGAAGGTCGGCGTGTGTACGTGACGGAAGTGGTGGCTGATAGCGTC	300
CAATTTCTTGAGCCGAAAGGAACGAGCGAGCAGCGAGGGGCGACAGCAGG	
CGGCTACTATGGGGATCCATTCCCATTTCGGGCAAGATCAGAACCACCAAT	400
ATCCGAACGAAAAAGGGTTTGGCCGCATCGATGACGATCCTTTCGCCAAT	
GACGGCCAGCCGATCGATATTTCTGATGATGATTTGCCGTTT	492

FIG. 80

MINRVILVGRLTRDPELRYTPSGVAVATFTLAVNRPFTNQSYENQEGRRV	
YVTEVVADSVQFLEPKGTSEQRGATAGGYQGERETDFIQCVVWRRQAEN	100
VANFLKKGSLAGVDGRLQTRGDPFPFGQDQNHQYPNEKGFGRIDDDPFAN	
DGQPIDISDDDLPF	164

FIG. 81

ATGCTGGAACGCGTATGGGGAACATTGAAAAACGGCGTTTTTCTCCCCT
 TTATTTATTATACGGCAATGAGCCGTTTTTTATTAACGGAAACGTATGAGC 100
 GATTGGTGAACGCAGCGCTTGGCCCCGAGGAGCGGGAGTGGAACCTTGCT
 GTGTACGACTGCGAGGAAACGCCGATCGAGGCGGCGCTTGAGGAGGCCGA 200
 GACGGTGCCGTTTTTCGGCGAGCGGCGTGTCAATCTCATCAAGCATCCAT
 ATTTTTTTTACGTCTGAAAAAGAGAAGGAGATCGAACATGATTTGGCGAAG 300
 CTGGAGGCGTACTTGAAGGCGCCGTCGCCGTTTTTCGATCGTCGTCTTTTT
 CGCGCCGTACGAGAAGCTTGATGAGCGAAAAAAATTACGAAGCTCGCCA 400
 AAGAGCAAAGCGAAGTCGTCATCGCCGCCCGCTCGCCGAAGCGGAGCTG
 CGTGCCTGGGTGCGGCGCCGCATCGAGAGCCAAGGGGCGCAAGCAAGCGA 500
 CGAGGCGATTGATGTCCTGTTGCGGCGGGCCGGGACGCAGCTTTCCGCCT
 TGGCGAATGAAATCGATAAATTGGCCCTGTTTGCCGGATCGGGCGGAACC 600
 ATCGAGGCGGCGGCGGTTGAGCGGCTTGTCGCCCGCACGCCGGAAGAAAA
 CGTATTTGTGCTTGTCGAGCAAGTGCGCAAGCGCGACATTCCAGCAGCGT 700
 TGCAGACGTTTTATGATCTGCTTGAAAACAATGAAGAGCCGATCAAAATT
 TTGGCGTTGCTCGCCGCCCATTTCCGCTTGCTTTTCGCAAGTGAAATGGCT 800
 TGCCTCCTTAGGCTACGGACAGGCGCAAATTGCTGCGGCGCTCAAGGTGC
 ACCCGTTCCGCGTCAAGCTCGCTCTTGCTCAAGCGGCCCGCTTCGCTGAC 900
 GGAGAGCTTGCTGAGGCGATCAACGAGCTCGCTGACGCCGATTACGAAGT
 GAAAAGCGGGGCGGTCGATCGCCGTTGGCCGTTGAGCTGCTTCTGATGC 1000
 GCTGGGGCGCCCGCCCGGCGCAAGCGGGGCGCCACGGCCGGCGG

FIG. 82

MLERVWGNIEKRRFSPLYLLYGNEPFLLTETYERLVNAALGPEEREWNLA
 VYDCEETPIEAALAEAETVPFFGERRVILIKHPYFFTSEKEKEIEHDLAK 100
 LEAYLKAPSPFSIVVFFAPYEKLDERKKITKLAKQSEVVIAAPLAEAEEL
 RAWVRRRIESQGAQASDEAIDVLLRRAGTQLSALANEIDKLALFAGSGGT 200
 IEAAVERLVARTPEENVFVLVEQVAKRDI PAALQTFYDLLENNEEPIKI
 LALLAAHFRLLSQVKWLASLGYGQAQIAAALKVHPFRVKLALAQAARFAD 300
 GELAEAINELADADYEVKSGAVDRRLAVELLLMRWGARPAQAGRHR

FIG. 83

ATGCGATGGGAACAGCTAGCGAAACGCCAGCCGGTGGTGGCGAAAATGCT	
GCAAAGCGGCTTGAAAAAGGGCGGATTTCTCATGCGTACTTGTTTGAGG	100
GGCAGCGGGGGACGGGCAAAAAAGCGGCCAGTTTGTTGTTGGCGAAACGT	
TTGTTTTGTCTGTCCCCAATCGGAGTTTCCCCGTGTCTAGAGTGCCGCAA	200
CTGCCGGCGCATCGACTCCGGCAACCACCCTGACGTCCGGGTGATCGGCC	
CAGATGGAGGATCAATCAAAAAGGAACAAATCGAATGGCTGCAGCAAGAG	300
TTCTCGAAAACAGCGGTGAGTCGGATAAAAAAATGTACATCGTTGAGCA	
CGCCGATCAAATGACGACAAGCGCTGCCAACAGCCTTCTGAAATTTTGG	400
AAGAGCCGCATCCGGGGACGGTGGCGGTATTGCTGACTGAGCAATACCAC	
CGCCTGCTAGGGACGATCGTTTCCCGCTGTCAAGTGCTTTCGTTCCGGCC	500
GTTGCCGCCGGCAGAGCTCGCCAGGGACTTGTGAGGAGCACGTGCCGT	
TGCCGTTGGCGCTGTTGGCTGCCATTTGACAAACAGCTTCGAGGAAGCA	600
CTGGCGCTTGCCAAAGATAGTTGGTTTGCCGAGGCGGAACATTAGTGCT	
ACAATGGTATGAGATGCTGGGCAAGCCGAGCTGCAGCTTTTGTTTTTCA	700
TCCACGACCGCTTGTTTCCGCATTTTTTGAAAGCCATCAGCTTGACCTT	
GGACTTG	757

FIG. 84

MRWEQLAKRQPVVAKMLQSGLEKGRISHAYLFEGQRGTGKKAASLLAKR	
LFCLSPIGVSPCLECRNCRRIDSGNHPDVRVIGPDGGSIKKEQIEWLQQE	100
FSKTAVESDKMYIVEHADQMTTSAANSLLKFLEEPHPGTVAVLLTEQYH	
RLLGTIVSRCQVLSFRPLPPAELAQGLVEEHVPLPLALLAAHLTNSFEEA	200
LALAKDSWFAEARTLVLQWYEMLGKPELQLLFFIHDRLFPHFLESHQLDL	
GL	252

FIG. 85

GTGGCATACCAAGCGTTATATCGCGTGTTTCGGCCGCAGCGCTTTGCGGA	
CATGGTCGGCCAAGAACACGTGACCAAGACGTTGCAAAGCGCCCTGCTTC	100
AACATAAAATATCGCACGCTTACTTATTTTCCGGCCCCGCGCGGTACAGGA	
AAAACGAGCGCAGCGAAAATTTTCGCCAAGGCGGTCAACTGTGAACAGGC	200
GCCAGCGGCGGAGCCATGCAATGAGTGTCCAGCTTGCCTCGGCATTACGA	
ATGGAACGGTTCCTGATGTGCTGGAATTGACGCTGCTTCCAACAACCGC	300
GTCGATGAAATTCGTGATATCCGTGAGAAGGTGAAATTTGCGCCAACGTC	
GGCCCCGTACAAAGTGTATATCATCGACGAGGTGCATATGCTGTGATCG	400
GTGCGTTTAACGCGCTGTTGAAAACGTTGGAGGAGCCGCCGAAACACGTC	
ATTTTCATTTTGGCCACGACCGAGCCGCACAAAATTCCGGCGACGATCAT	500
TTCCCGCTGCCAACGGTTCGATTTTCGCCGCATCCCGCTTCAGGCGATCG	
TTTCACGGCTAAAGTACGTCGCAAGCGCCCAAGGTGTGAGGCGTCAGAT	600
GAGGCATTGTCCGCCATCGCCCGTGCTGCAGACGGGGGGATGCGCGATGC	
GCTCAGCTTGCTTGATCAAGCCATTTTCGTTTCAGCGACGGGAAACTTCGGC	700
TCGACGACGTGCTGGCGATGACCGGGGCTGCATCATTTGCCGCCTTATCG	
AGCTTCATCGAAGCCATCCACCGCAAAGATACAGCGGCGGTTCTTCAGCA	800
CTTGGAACGATGATGGCGCAAGGGAAAGATCCGCATCGTTTGGTTGAAG	
ACTTGATTTTGTACTATCGCGATTTATTGCTGTACAAAACCGCTCCCTAT	900
GTGGAGGGAGCGATTCAAATTGCTGTGCTTGACGAAGCGTTCACTTCACT	
GTGCGAAATGATTCGGGTTTCCAATTTATACGAGGCCATCGAGTTGCTGA	1000
ACAAAAGCCAGCAAGAGATGAAGTGGACAAACCACCGCGCCTTCTGTTG	
GAAGTGGCGCTTGTGAAACTTTGCCATCCATCAGCCGCCGCCCGCTCGCT	1100
GTCGGCTTCCGAGTTGGAACCGTTGATAAAGCGGATTGAAACGCTGGAGG	
CGGAATTGCGGCGCCTGAAGGAACAACCGCCTGCCCCTCCGTGACCGCC	1200
GCGCCGTTGAAAAAACTGTCCAAACCGATGAAAACGGGGGGATATAAAGC	
CCCGGTTGGCCGCATTTACGAGCTGTTGAAACAGGCGACGCATGAAGATT	1300
TAGCTTTGGTGAAAGGATGCTGGGCGGATGTGCTCGACACGTTGAAACGG	
CAGCATAAAGTGTGCGACGCTGCCTTGCTGCAAGAGAGCGAGCCGTTGC	1400
AGCGAGCGCCTCAGCGTTTGTATTAAAATTCAAATACGAAATCCACTGCA	
AAATGGCGACCGATCCCACAAGTTCGGTCAAAGAAAACGTCGAAGCGATT	1500
TTGTTTGAGCTGACAAACCGCCGCTTTGAAATGGTAGCCATTCCCGAGGG	
AGAATGGGGAAAAATAAGAGAAGAGTTCATCCGCAATAAGGACGCCATGG	1600
TGGAAAAAAGCGAAGAAGATCCGTTAATCGCCGAAGCGAAGCGGCTGTTT	
GGCGAAGAGCTGATCGAAATTAAAGAA	1677

FIG. 86

ATGGTGACAAAAGAGCAAAAAGAGCGGTTTCTCATCCTGCTTGAGCAGCT	
GAAGATGACGTCGGACGAATGGATGCCGCATTTTCGTGAGGCAGCCATTC	100
GCAAAGTCGTGATCGATAAAGAGGAGAAAAGCTGGCATTTTTATTTTCAG	
TTGACAACGTGCTGCCGGTTCATGTATACAAAACGTTTGCCGATCGGCT	200
GCAGACGGCGTTCCGCCATATCGCCGCCGTCCGCCATACGATGGAGGTCG	
AAGCGCCGCGCGTAACCTGAGGCGGATGTGCAGGCGTATTGGCCGCTTTGC	300
CTTGCCGAGCTGCAAGAAGGCATGTGCGCCGCTTGTCGATTGGCTCAGCCG	
GCAGACGCCTGAGCTGAAAGGAAACAAGCTGCTTGTCGTTGCCCGCCATG	400
AAGCGGAAGCGCTGGCGATCAAACGGCGGTTCCGCCAAAAAATCGCTGAT	
GTGTACGCTTCGTTTGGGTTTCCCCCCTTCAGCTTGACGTCAGCGTCGA	500
GCCGTCCAAGCAAGAAATGGAACAGTTTTTTGGCGCAAAAACAGCAAGAGG	
ACGAAGAGCGAGCGCTTGCTGTACTGACCGATTTAGCGAGGGAAGAAGAA	600
AAGGCCGCGTCTGCGCCGCCGTCCGGTCCGCTTGTCATCGGCTATCCGAT	
CCGCGACGAGGAGCCGGTGCGGCGGCTTGAAACGATCGTCGAAGAAGAGC	700
GGCGCGTCGTTGTGCAAGGCTATGTATTTGACGCCGAAGTGAGCGAATTA	
AAAAGCGGCCGCACGCTGTTGACCATGAAAATCACAGATTACACGAACTC	800
GATTTTAGTCAAAATGTTCTCGCGCGACAAAGAGGACGCCGAGCTTATGA	
GCGGCGTCAAAAAGGCATGTGGGTGAAAGTGCGCGGCAGCGTGCAAAAC	900
GATACGTTTCGTCCGTGATTTGGTTCATCATCGCCAACGATTTGAACGAAAT	
CGCCGCAAACGAACGGCAAGATACGGCGCCGGAAGGGGAAAAGAGGGTCG	1000
AGCTCCATTTGCATACCCCGATGAGCCAAATGGACGCGGTACCTCGGTG	
ACAAAACCTCATTGAGCAAGCGAAAAAATGGGGGCATCCGGCGATCGCCGT	1100
CACCGACCATGCCGTTGTTCACTCGTTTCCGGAGGCCTACAGCGCGGCGA	
AAAAACACGGCATGAAGGTCATTTACGGCCTTGAGGCGAACATCGTCGAC	1200
GATGGCGTGCCGATCGCCTACAATGAGACGCACCGCCGTCTTTCGGAGGA	
AACGTACGTCGTCTTTGACGTCGAGACGACGGGCCTGTGCGGCTGTGTACA	1300
ATACGATCATTGAGCTGGCGGCGGTGAAAGTGAAAGACGGCGAGATCATC	
GACCGATTTCATGTCGTTTGCCAACCTGGACATCCGTTGTGCGGTGACAAC	1400
GATGGAGCTGACTGGGATCACCGATGAGATGGTGAAAGACGCCCCGAAGC	
CGGACGAGGTGCTAGCCCGTTTTGTTGACTGGGCGGCGATGCGACGCTT	1500
GTTGCCACACAACGCCAGCTTTGACATCGGTTTTTTTAAACGCGGGCCTCGC	
TCGCATGGGGCGCGGCAAAATCGCGAATCCAGTCATCGATACGCTCGAGC	1600
TGGCCCGTTTTTTTATACCCGGATTTGAAAAACCATCGGCTCAATACATTG	
TGCAAAAAATTTGACATTGAATTGACGCAGCATACCGCGCCATCTACGA	1700
CGCGGAGGCGACCGGGCATTGCTTATGCGGCTGTTGAAGGAAGCGGAAG	
AGCGCGGCATACTGTTTCATGACGAATTAAACAGCCGCACGCACAGCGAA	1800
GCGTCCTATCGGCTTGCGCGCCCGTTCCATGTGACGCTGTTGGCGCAAAA	
CGAGACTGGATTGAAAAATTTGTTCAAGCTTGTGTTCATTGTCGCACATTC	1900
AATATTTTACCGTGTGCCGCGCATCCCGCGCTCCGTGCTCGTCAAGCAC	
CGCGACGGCCTGCTTGTCGGCTCGGGCTGCGACAAAGGAGAGCTGTTTGA	2000
CAACTTGATCCAAAAGGCGCCGGAAGAAGTCGAAGACATCGCCCGTTTTT	
ACGATTTTCTTGAAGTGATCCGCCGGACGTGTACAAGCCGCTCATCGAG	2100
ATGGATTATGTGAAAGACGAAGAGATGATCAAAAACATCATCCGCAGCAT	
CGTCGCCCTTGGTGAGAAGCTTGACATCCCGGTTGTCGCCACTGGCAACG	2200

FIG. 88A

TCCATTACTTGAACCCAGAAGATAAAATTTACCGGAAAATCTTAATCCAT
 TCGCAAGGCGGGGCGAATCCGCTCAACCGCCATGAACTGCCGGATGTATA 2300
 TTTCCGTACGACGAATGAAATGCTTGACTGCTTCTCGTTTTTAGGGCCGG
 AAAAAGCGAAGGAAATCGTCGTTGACAACACGCAAAAAATCGCTTCGTTA 2400
 ATCGGCGATGTCAAGCCGATCAAAGATGAGCTGTATACGCCGCGCATTGA
 AGGGGCGGACGAGGAAATCAGGGAAATGAGCTACCGGCGGGCGAAGGAAA 2500
 TTTACGGCGACCCGTTGCCGAACTTGTTGAAGAGCGGCTTGAGAAGGAG
 CTAAAAAGCATCATCGGCCATGGCTTTGCCGTCAATTTATTTGATCTCGCA 2600
 CAAGCTTGTAAGAAAATCGCTCGATGACGGCTACCTTGTCGGGTTCGCGCG
 GATCGGTTCGGCTCGTCGTTTGTGCGGACGATGACGGAAATCACCGAGGTC 2700
 AATCCGCTGCCGCCGCATTACGTTTGCCCGAACTGCAAGCATTTCGGAGTT
 CTTTAACGACGGTTCAGTCGGCTCAGGGTTTGATTTGCCGGATAAAAACT 2800
 GCGGCGATGTGGGACGAAATACAAGAAAGACGGGCACGACATCCCGTTT
 GAGACGTTTTCTCGGCTTTAAAGGCGACAAAGTGCCGGATATCGACTTGAA 2900
 CTTTTCCGGCGAATACCAGCCGCGCGCCCACTATACGAAAGTGCTGT
 TTGGCGAAGACAACGTCTACCGCGCCGGGACGATTGGCACGGTCGCTGAC 3000
 AAAACGGCGTACGGATTTGTCAAAGCGTATGCGAGCGACCATAACTTAGA
 GCTGCGCGGCGCGGAAATCGACGGCTCGCGGCTGGCTGCACCGGGGTGAA 3100
 GCGGACGACCGGGCAGCATCCGGGCGGCATCATCGTCGTCCCGGATTATA
 TGAAATTTACGATTTTACGCCGATTCAATATCCGGCCGATGACACGTCC 3200
 TCTGAATGGCGGACGACCCATTTCTGACTTCCATTTCGATCCACGACAATTT
 GTTGAAGCTCGATATTCTCGGGCACGACGATCCGACGGTCATTTCGCATGC 3300
 TGCAAGATTTAAGCGGCATCGATCCGAAAACGATCCCGACCGACGACCCG
 GATGTGATGGGCATTTTCAGCAGCACCGAGCCGCTTGCGGTTACGCCGGA 3400
 GCAAATCATGTGCAATGTCGGCACGATCGGCATTCCGGAGTTTGGCACGC
 GCTTCGTTTCGGCAAATGTTGGAAGAGACAAGGCCAAAAACGTTTTCCGAA 3500
 CTCGTGCAAATTTCCGGCTTGTCGCGACGGCACCGATGTGTGGCTCGGCAA
 CGCGCAAGAGCTCATTTAAAACGGCACGTGTACGTTATCGGAAGTCATCG 3600
 GCTGCCGCGACGACATTATGGTCTATTTGATTTACCGCGGGCTCGAGCCG
 TCGCTCGCTTTTAAAATCATGGAATCCGTGCGCAAAGGAAAAGGCTTAAC 3700
 GCCGGAGTTTGAAGCAGAAATGCGCAAACATGACGTGCCGGAGTGGTACA
 TCGATTTCATGCAAAAAAATCAAGTACATGTTCCCGAAAAGCGCACGCCGCC 3800
 GCCTACGTGTTAATGGCGGTGCGCATCGCCTACTTTAAGGTGCACCATCC
 GCTTTTGTATTACGCGTCGTACTTTACGGTGCGGGCGGAGGACTTTGACC 3900
 TTGACGCCATGATCAAAGGATCACCCGCCATTTCGCAAGCGGATTGAGGAA
 ATCAACGCCAAAGGCATTTCAGGCGACGGCGAAAGAAAAAGCTTGCTCAC 4000
 GGTTCTTGAGGTGGCCTTAGAGATGTGCGAGCGCGGCTTTTCCTTTAAAA
 ATATCGATTTGTACCGCTCGCAGGCGACGGAATTCGTCATTGACGGCAAT 4100
 TCTCTCATTCCGCCGTTCAACGCCATTCCGGGGCTTGGGACGAACGTGGC
 GCAGGCGATCGTGCGCGCCCGGAGGAAGGCGAGTTTTTGTGCAAGGAGG 4200
 ATTTGCAACAGCGCGGCAAATTTGTCGAAAACGCTGCTCGAGTATCTAGAA
 AGCCGCGGCTGCCTTGACTCGCTTCCAGACCATAACCAGCTGTGCTGTT 4300
 T

FIG. 88B

MVTKEQKERFLILLEQLKMTSDEWMPHFREAAIRKVVIDKEEKSWHFFYFQ	
FDNVLPVHVYKTFADRLQTAFRHIAAVRHTMEVEAPRVTEADVQAYWPLC	100
LAELQEGMSPLVDWLSRQTPELKGNKLLVVARHEAEALAIKRRFAKKIAD	
VYASFQFPPLQLDVSVEPSKQEMEQLAQKQEQEDEERALAVLTDLAREEE	200
KAASAPPSGPLVIGYPIDEEPVRRLLETIVEEERRVVVQGYVFDAEVSEL	
KSGRTLLTMKITDYTNSILVKMFSRDKEDAELMSGVKKGMWVKVRGSVQN	300
DTFVRDLVIIANDLNEIAANERQDTAPEGEKRVELHLHTPMSQMDAVTSV	
TKLIEQAKKWGHPIAVTDHAVVQSFPEAYSAAKKHGMKVIYGLEANIVD	400
DGVPIAYNETHRRLSEETYVVFVETTGLSAVYNTIIELAAVKVKDGEII	
DRFMSFANPGHPLSVTTMELTGITDEMVKDAPKPDEVLARFVDWAGDATL	500
VAHNASFDIGFLNAGLARMGRGKIANPVIDTLELARFLYPDLKNHRLNTL	
CKKFDIELTQHHRAIYDAEATGHLLMRLLEKEAEERGILFHDELNSRTHSE	600
ASYRLARPFHVTLTLLAQNETGLKNLFLKLVSLSHIQYFHRVPRIPRSVLVKH	
RDGLLVGSGCDKGELFDNLIQKAPEEVEDIARFYDFLEVHPPDVYKPLIE	700
MDYVKDEEMIKNIIRSIVALGEKLDIPVVATGNVHYLNPEDKIYRKILIH	
SQGGANPLNRHELDPVYFRTTNEMLDCFSFLGPEKAKEIVVDNTQKIASL	800
IGDVKPIKDELYTPRIEGADEEIREMSYRRAKEIYGDPLPKLVEERLEKE	
LKSIIGHGFAVIYLISHKLVKKSLDDGYLVGSRGSGSSFVATMTEITEV	900
NPLPPHYVCPNCKHSEFFNDGSGVSGFDLPDKNCPRCGTYKKDGHDIPIF	
ETFLGFKGDKVPDIDLNFSGEYQPRAHNYTKVLFGEDNVYRAGTIGTVAD	1000
KTAYGFVKAYASDHNLELRGAEIDLAAGCTGVKRTTGQHPGGIIVVPDYM	
EIYDFTPIQYPADDTSSSEWRTHFDHFSIHDNLLKLDILGHDDPTVIRML	1100
QDLSGIDPKTIPTDDPDVMGIFSSSTEPLGVTPEQIMCNVGTIGIPEFGTR	
FVRQMLEETRPKTFSELVQISGLSHGTDVWLGNALQELIQNGTCTLSEVIG	1200
CRDDIMVYLIYRGLEPSLAFKIMESVRKGKGLTPEFEAEMRKHDVPEWYI	
DSCKKIKYMFPAKAAAYVLMAVRIAYFKVHHPLLYASYFTVRAEDFDL	1300
DAMIKGSPAIRKRIEEINAKGIQATAKEKSLTVLEVALEMCEGFSFKN	
IDLYRSQATEFVIDGNSLIPPFNAIPGLGTNVAQAIVRAREEGEFLSKED	1400
LQQRGKLSKTLLEYLESRGCLDSLPHNQLSLF	

FIG. 89

COMBINED DECLARATION FOR PATENT
APPLICATION AND POWER OF ATTORNEY
(Includes Reference to PCT International Applications)

ATTORNEY'S DOCKET NUMBER

22221/1030 (RU-339)

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

- I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

**ENZYMES DERIVED FROM THERMOPHILIC ORGANISMS THAT FUNCTION AS A CHROMOSOMAL
REPLICASE, PREPARATION AND USE THEREOF**

the specification of which (check only one item below):

☒ is attached hereto.

☐ was filed as U.S. Patent Application Serial No. _____ on _____ and was amended on _____
(if applicable).

☐ was filed as PCT International Application No. _____ on _____ and was amended under PCT Article 19 on _____
(if applicable).

I hereby state that I have reviewed and understand the contents of the above-identified specifications, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the patentability of this application in accordance with Title 37, Code of Federal Regulations, § 1.56(a).

I hereby claim foreign priority benefits under Title 35, United States Code, § 119 of any foreign application(s) for patent or inventor's certificate or of any PCT international application(s) designating at least one country other than the United States listed below and have also identified below any foreign application(s) for patent or inventor's certificate or any PCT international application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) of which priority is claimed:

PRIOR APPLICATION(S) AND ANY PRIORITY CLAIMS UNDER 35 U.S.C. 119:

COUNTRY (If PCT, indicate "PCT")	APPLICATION NUMBER	DATE OF FILING (day, month, year)	PRIORITY CLAIMED UNDER 35 USC 119
United States	09/057,416	8-APRIL-1998	<input checked="" type="checkbox"/> YES <input type="checkbox"/> NO
United States	08/823,407	8-APRIL-1997	<input checked="" type="checkbox"/> YES <input type="checkbox"/> NO
United States	60/143,202	8-APRIL-1997	<input checked="" type="checkbox"/> YES <input type="checkbox"/> NO

I hereby claim the benefit under Title 35, United States Code, § 120 of any United States application(s) or PCT international application(s) designating the United States of America that is/are listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in that/those prior application(s) in the manner provided by the first paragraph of Title 35, United States Code, § 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, § 1.56(a) which occurred between the filing date of the prior application(s) and the national or PCT International filing date of this application:

PRIOR U.S. APPLICATIONS OR PCT INTERNATIONAL APPLICATIONS DESIGNATING THE U.S. FOR BENEFIT UNDER 35 U.S.C. 120:

U.S. APPLICATIONS		STATUS (Check One)			
U.S. APPLICATION NUMBER	U.S. FILING DATE	PATENTED	PENDING	ABANDONED	
PCT APPLICATIONS DESIGNATING THE U.S.					
PCT APPLICATION NO.	PCT FILING DATE	U.S. SERIAL NUMBERS ASSIGNED (if any)			

COMBINED DECLARATION FOR PATENT APPLICATION
AND POWER OF ATTORNEY (Continue)

ATTORNEY'S DOCKET NUMBER
22221/1030 (RU-339)

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. **Michael L. Goldman, Registration No. 30,727; Joseph M. Noto, Registration No. 32,163; Grant E. Pollack, Registration No. 34,097; Ann R. Pokalsky, Registration No. 34,697; Gunnar G. Leinberg, Registration No. 35,584; Edwin V. Merkel, Registration No. 40,087; Georgia Evans, Registration No. 44,597; Alice Y. Choi, Registration No. 45,758**

Send Correspondence to:

**Michael L. Goldman
NIXON PEABODY LLP
Clinton Square, P.O. Box 31051
Rochester, New York 14603**

Direct Telephone Calls to:
(name and telephone number)
**Michael L. Goldman
(716) 263-1304**

201	FULL NAME OF INVENTOR	FAMILY NAME O'Donnell	FIRST GIVEN NAME Michael	SECOND GIVEN NAME E.
	RESIDENCE & CITIZENSHIP	CITY Hastings-on-Hudson	STATE/FOREIGN COUNTRY New York	COUNTRY OF CITIZENSHIP United States
	POST OFFICE ADDRESS	P.O. ADDRESS 16 Maple Lane	CITY Hastings-on-Hudson	STATE & ZIP CODE/CTRY New York 10706/USA
202	FULL NAME OF INVENTOR	FAMILY NAME Yuzhakov	FIRST GIVEN NAME Alexander	SECOND GIVEN NAME
	RESIDENCE & CITIZENSHIP	CITY Belmont	STATE/FOREIGN COUNTRY Massachusetts	COUNTRY OF CITIZENSHIP Russia
	POST OFFICE ADDRESS	P.O. ADDRESS 24 Clyde Street	CITY Belmont	STATE & ZIP CODE/CTRY Massachusetts 02478/USA
203	FULL NAME OF INVENTOR	FAMILY NAME Yurieva	FIRST GIVEN NAME Olga	SECOND GIVEN NAME
	RESIDENCE & CITIZENSHIP	CITY New York	STATE/FOREIGN COUNTRY New York	COUNTRY OF CITIZENSHIP Russia
	POST OFFICE ADDRESS	P.O. ADDRESS 430 East 63rd Str., Apt. 3G	CITY New York	STATE & ZIP CODE/CTRY New York 10021/USA
204	FULL NAME OF INVENTOR	FAMILY NAME Jeruzalmi	FIRST GIVEN NAME David	SECOND GIVEN NAME
	RESIDENCE & CITIZENSHIP	CITY New York	STATE/FOREIGN COUNTRY New York	COUNTRY OF CITIZENSHIP United States
	POST OFFICE ADDRESS	P.O. ADDRESS 1161 York Avenue, #111	CITY New York	STATE & ZIP CODE/CTRY New York 10021/USA
205	FULL NAME OF INVENTOR	FAMILY NAME Bruck	FIRST GIVEN NAME Irina	SECOND GIVEN NAME
	RESIDENCE & CITIZENSHIP	CITY New York	STATE/FOREIGN COUNTRY New York	COUNTRY OF CITIZENSHIP United States
	POST OFFICE ADDRESS	P.O. ADDRESS 1161 York Avenue, Apt. 11M	CITY New York	STATE & ZIP CODE/CTRY New York 10021/USA
206	FULL NAME OF INVENTOR	FAMILY NAME Kuriyan	FIRST GIVEN NAME John	SECOND GIVEN NAME
	RESIDENCE & CITIZENSHIP	CITY New York	STATE/FOREIGN COUNTRY New York	COUNTRY OF CITIZENSHIP United States
	POST OFFICE ADDRESS	P.O. ADDRESS 430 East 63rd, Apt. 12E	CITY New York	STATE & ZIP CODE/CTRY New York 10021/USA

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statement may jeopardize the validity of the application or any patent issuing thereon.

SIGNATURE OF INVENTOR 201 UNSIGNED	SIGNATURE OF INVENTOR 202 UNSIGNED	SIGNATURE OF INVENTOR 203 UNSIGNED
DATE	DATE	DATE
SIGNATURE OF INVENTOR 204 UNSIGNED	SIGNATURE OF INVENTOR 205 UNSIGNED	SIGNATURE OF INVENTOR 206 UNSIGNED
DATE	DATE	DATE

Page 3 of 3

09716964-412100

SEQUENCE LISTING

<110> O'Donnell, Michael E.
Yuzhakov, Alexander
Yurieva, Olga
Jeruzalmi, David
Bruck, Irina
Kuriyan, John

<120> ENZYMES DERIVED FROM THERMOPHILIC ORGANISMS THAT
FUNCTION AS A CHROMOSOMAL REPLICASE, PREPARATION AND
USE THEREOF

<130> 22221/1030

<140>

<141>

<150> 60/143,202

<151> 1997-04-08

<150> 08/823,407

<151> 1997-04-08

<150> 09/057,416

<151> 1998-04-08

<160> 212

<170> PatentIn Ver. 2.1

<210> 1

<211> 2007

<212> DNA

<213> *Thermus thermophilus*

<400> 1

```
tccgggggtg gggttcccag gtagaccccg gcccctcccg tgagcccctt taccagggcc 60
gccacctcct ccaggggggc caaggcgtgc aaggagagga acgtccgcac cacgccctat 120
actagccttg tgagcgccct ctaccgccgc ttccgcccc tcacctcca ggaggtggtg 180
gggcaggagc acgtgaagga gcccctcctc aaggccatcc gggaggggag gctcgcccag 240
gcctacctct tctccgggcc caggggcgtg ggcaagacca ccacggcgag gctcctcgcc 300
atggcggttg ggtgccaggg ggaagacccc ccttgcgggg totgccccca ctgccaggcg 360
gtgcagaggg gcgcccaccc ggacgtggtg gacattgacg ccgccagcaa caactccgtg 420
gaggacgtgc gggagctgag ggaaaggatc cacctcgccc ccctctctgc ccccaggaag 480
gtcttcatcc tggacgaggc ccacatgctc tccaaaagcg ccttcaacgc cctcctcaag 540
accctggagg agccccgcgc ccacgtcctc ttcgtcttcg ccaccaccga gcccagagag 600
atgcccccca ccatactctc ccgcacccag cacttccgct tccgccgcct cacggaggag 660
```

gagatcgctt ttaagctccg gcgcacccctg gagggcggtg ggcgggaggc ggaggaggag 720
gccctcctcc tcctcgcccg cctggcggac ggggccctta gggacgcgga aagcctcctg 780
gagcgcttcc tcctcctgga agggcccccctc acccggaagg aggtggagcg cgccctaggc 840
tccccccag ggaccggggt ggccgagatc gccgcctccc tcgcgagggg gaaaacggcg 900
gaggccctgg gctcgcggcg gcgcctctac ggggaagggt acgccccgag gagcctggtc 960
tcgggccttt tggaggtgtt ccgggaaggc ctctacgcgc ccttcggcct cgcggaacc 1020
ccccctcccg ccccgcccca ggccctgatc gccgccatga ccgcccggga cgaggccatg 1080
gagcgccctg cccgcgcctc cgacgcctta agcctggagg tggccctcct ggaggcgga 1140
agggccctgg ccgcccaggc cctaccccag cccacgggcg ctccttcccc agaggtcggc 1200
cccaagccgg aaagccccc gaccccgga ccccaaggc ccgaggaggc gcccgacctg 1260
cgggagcggt ggcgggcctt cctcgaggcc ctcaggccca ccctacgggc cttcgtgcgg 1320
gaggcccgcc cggaggtccg ggaaggccag ctctgcctcg ctttccccga ggacaaggcc 1380
ttccactacc gcaaggcctc ggaacagaag gtgaggctcc tccccctggc ccaggcccat 1440
ttcgggggtg agggaggtcgt cctcgtcctg gagggagaaa aaaaaagcct gagcccaagg 1500
ccccgcccgg cccacctcc tgaagcgccc gcaccccg gcccctccga ggaggaggta 1560
gaggcgaggg aagcggcgga ggaggcccg gagggagcct tgaggcggtt ggtccgcctc 1620
ctggggggggc ggggtgctctg ggtgcggcg ccaggaccg gggaggcgcc ggaggaggaa 1680
cccctgagcc aagacgagat aggggggtact ggtatataat gggggcatga cgcgggaccac 1740
cgacctcgga caagagaccg tggacaacat cctcaagcgc ctccgccgta ttgaggggcca 1800
ggtgcggggg ctccagaaga tgggtggccga gggccgcccc tgcgacgagg tcctcaccga 1860
gatgaccgcc accaagaagg ccatggaggc ggcggccacc ctgatcctcc acgagttcct 1920
gaacgtctgc gccgcccagg tctccgagg caaggtgaac cccaagaagc ccgaggagat 1980
cgccaccatg ctgaagaact tcattcta 2007

<210> 2

<211> 529

<212> PRT

<213> *Thermus thermophilus*

<400> 2

Met	Ser	Ala	Leu	Tyr	Arg	Arg	Phe	Arg	Pro	Leu	Thr	Phe	Gln	Glu	Val
1				5					10					15	
Val	Gly	Gln	Glu	His	Val	Lys	Glu	Pro	Leu	Leu	Lys	Ala	Ile	Arg	Glu
			20					25					30		
Gly	Arg	Leu	Ala	Gln	Ala	Tyr	Leu	Phe	Ser	Gly	Pro	Arg	Gly	Val	Gly
		35					40					45			
Lys	Thr	Thr	Thr	Ala	Arg	Leu	Leu	Ala	Met	Ala	Val	Gly	Cys	Gln	Gly
	50					55					60				
Glu	Asp	Pro	Pro	Cys	Gly	Val	Cys	Pro	His	Cys	Gln	Ala	Val	Gln	Arg
65					70					75					80
Gly	Ala	His	Pro	Asp	Val	Val	Asp	Ile	Asp	Ala	Ala	Ser	Asn	Asn	Ser
				85				90						95	

Val Glu Asp Val Arg Glu Leu Arg Glu Arg Ile His Leu Ala Pro Leu
 100 105 110

Ser Ala Pro Arg Lys Val Phe Ile Leu Asp Glu Ala His Met Leu Ser
 115 120 125

Lys Ser Ala Phe Asn Ala Leu Leu Lys Thr Leu Glu Glu Pro Pro Pro
 130 135 140

His Val Leu Phe Val Phe Ala Thr Thr Glu Pro Glu Arg Met Pro Pro
 145 150 155 160

Thr Ile Leu Ser Arg Thr Gln His Phe Arg Phe Arg Arg Leu Thr Glu
 165 170 175

Glu Glu Ile Ala Phe Lys Leu Arg Arg Ile Leu Glu Ala Val Gly Arg
 180 185 190

Glu Ala Glu Glu Glu Ala Leu Leu Leu Leu Ala Arg Leu Ala Asp Gly
 195 200 205

Ala Leu Arg Asp Ala Glu Ser Leu Leu Glu Arg Phe Leu Leu Leu Glu
 210 215 220

Gly Pro Leu Thr Arg Lys Glu Val Glu Arg Ala Leu Gly Ser Pro Pro
 225 230 235 240

Gly Thr Gly Val Ala Glu Ile Ala Ala Ser Leu Ala Arg Gly Lys Thr
 245 250 255

Ala Glu Ala Leu Gly Leu Ala Arg Arg Leu Tyr Gly Glu Gly Tyr Ala
 260 265 270

Pro Arg Ser Leu Val Ser Gly Leu Leu Glu Val Phe Arg Glu Gly Leu
 275 280 285

Tyr Ala Ala Phe Gly Leu Ala Gly Thr Pro Leu Pro Ala Pro Pro Gln
 290 295 300

Ala Leu Ile Ala Ala Met Thr Ala Leu Asp Glu Ala Met Glu Arg Leu
 305 310 315 320

Ala Arg Arg Ser Asp Ala Leu Ser Leu Glu Val Ala Leu Leu Glu Ala
 325 330 335

Gly Arg Ala Leu Ala Ala Glu Ala Leu Pro Gln Pro Thr Gly Ala Pro
 340 345 350

Ser Pro Glu Val Gly Pro Lys Pro Glu Ser Pro Pro Thr Pro Glu Pro
 355 360 365

Pro Arg Pro Glu Glu Ala Pro Asp Leu Arg Glu Arg Trp Arg Ala Phe
 370 375 380

Leu Glu Ala Leu Arg Pro Thr Leu Arg Ala Phe Val Arg Glu Ala Arg
 385 390 395 400

Pro Glu Val Arg Glu Gly Gln Leu Cys Leu Ala Phe Pro Glu Asp Lys
 405 410 415

Ala Phe His Tyr Arg Lys Ala Ser Glu Gln Lys Val Arg Leu Leu Pro
 420 425 430

Leu Ala Gln Ala His Phe Gly Val Glu Glu Val Val Leu Val Leu Glu
 435 440 445

Gly Glu Lys Lys Ser Leu Ser Pro Arg Pro Arg Pro Ala Pro Pro Pro
 450 455 460

Glu Ala Pro Ala Pro Pro Gly Pro Pro Glu Glu Glu Val Glu Ala Glu
 465 470 475 480

Glu Ala Ala Glu Glu Ala Pro Glu Glu Ala Leu Arg Arg Val Val Arg
 485 490 495

Leu Leu Gly Gly Arg Val Leu Trp Val Arg Arg Pro Arg Thr Arg Glu
 500 505 510

Ala Pro Glu Glu Glu Pro Leu Ser Gln Asp Glu Ile Gly Gly Thr Gly
 515 520 525

Ile

<210> 3

<211> 1590

<212> DNA

<213> *Thermus thermophilus*

<400> 3

gtgagcgccc tctaccgccg cttccgcccc ctcaccttcc aggaggtggt ggggcaggag 60
 cacgtgaagg agccccctct caaggccatc cgggagggga ggctcgccca ggcctacctc 120
 ttctccgggc ccagggggtt gggcaagacc accacggcga ggctcctcgc catggcggtg 180
 ggggtgccagg gggaagaccc cccttgcggt gtctgcccc actgccaggc ggtgcagagg 240

```

ggcgcccacc cggacgtggt ggacattgac gccgccagca acaactccgt ggaggacgtg 300
cgggagctga gggaaaggat ccacctcgcc cccctctctg ccccaggaa ggtcttcac 360
ctggacgagg ccacatgct ctccaaaagc gccttcaacg cctcctcaa gacctggag 420
gagccccgcg ccacgtcct cttcgtcttc gccaccaccg agcccgagag gatgcccccc 480
accatcctct cccgcaccca gcacttcgcg ttccgcccgc tcacggagga ggagatcgcc 540
tttaagctcc ggcgcacccg ggaggccgtg gggcgggagg cggaggagga ggccctcctc 600
ctcctcgccc gcctggcgga cggggccctt agggacgcgg aaagcctcct ggagcgcttc 660
ctcctccttg aaggccccct cacccggaag gaggtggagc gcgccctagg ctccccccca 720
gggaccgggg tggccgagat cgcgcctcc ctcgcgaggg ggaaaacggc ggaggccctg 780
ggcctcgccc ggcgcctcta cggggaaggg tacgccccga ggagcctggt ctcgggcctt 840
ttggaggtgt tccgggaagg cctctacgcc gccttcggcc tcgcgggaac ccccttccc 900
gccccgcccc aggcctgat cgcgcctatg accgcctgg acgaggccat ggagcgccctc 960
gcccgccgct ccgacgcctt aagcctggag gtggccctcc tggaggcggg aaggggccctg 1020
gccgcccagg ccctaccca gccacgggc gctccttccc cagaggtcgg cccaagccg 1080
gaaagcccc cgaccccgga accccaagg cccgaggagg cgcgcgacct gcgggagcgg 1140
tggcgggect tcctcgaggc cctcaggccc accctacggg ccttcgtcgg ggaggcccg 1200
ccggaggtcc gggaaggcca gctctgcctc gctttccccg aggacaaggc cttccactac 1260
cgcaaggcct cggaacagaa ggtgaggctc ctccccctgg cccaggccca tttcgggggtg 1320
gaggaggtcg tcctcgtcct ggaggagaa aaaaaaagcc tgagcccaag gccccgccc 1380
gccccacctc ctgaagcgcc cgcacccccg ggccctcccg aggaggaggt agaggcggag 1440
gaagcggcgg aggaggcccc ggaggaggcc ttgaggcggg tgggtccgcct cctggggggg 1500
cgggtgctct ggggtgcggc gccaggacc cgggaggcgc cggaggagga acccctgagc 1560
caagacgaga tagggggtac tggtatataa 1590

```

<210> 4

<211> 464

<212> PRT

<213> *Thermus thermophilus*

<400> 4

```

Met Ser Ala Leu Tyr Arg Arg Phe Arg Pro Leu Thr Phe Gln Glu Val
  1              5              10              15

Val Gly Gln Glu His Val Lys Glu Pro Leu Leu Lys Ala Ile Arg Glu
      20              25              30

Gly Arg Leu Ala Gln Ala Tyr Leu Phe Ser Gly Pro Arg Gly Val Gly
      35              40              45

Lys Thr Thr Thr Ala Arg Leu Leu Ala Met Ala Val Gly Cys Gln Gly
      50              55              60

Glu Asp Pro Pro Cys Gly Val Cys Pro His Cys Gln Ala Val Gln Arg
      65              70              75              80

Gly Ala His Pro Asp Val Val Asp Ile Asp Ala Ala Ser Asn Asn Ser
      85              90              95

```

Val Glu Asp Val Arg Glu Leu Arg Glu Arg Ile His Leu Ala Pro Leu
 100 105 110
 Ser Ala Pro Arg Lys Val Phe Ile Leu Asp Glu Ala His Met Leu Ser
 115 120 125
 Lys Ser Ala Phe Asn Ala Leu Leu Lys Thr Leu Glu Glu Pro Pro Pro
 130 135 140
 His Val Leu Phe Val Phe Ala Thr Thr Glu Pro Glu Arg Met Pro Pro
 145 150 155 160
 Thr Ile Leu Ser Arg Thr Gln His Phe Arg Phe Arg Arg Leu Thr Glu
 165 170 175
 Glu Glu Ile Ala Phe Lys Leu Arg Arg Ile Leu Glu Ala Val Gly Arg
 180 185 190
 Glu Ala Glu Glu Glu Ala Leu Leu Leu Leu Ala Arg Leu Ala Asp Gly
 195 200 205
 Ala Leu Arg Asp Ala Glu Ser Leu Leu Glu Arg Phe Leu Leu Leu Glu
 210 215 220
 Gly Pro Leu Thr Arg Lys Glu Val Glu Arg Ala Leu Gly Ser Pro Pro
 225 230 235 240
 Gly Thr Gly Val Ala Glu Ile Ala Ala Ser Leu Ala Arg Gly Lys Thr
 245 250 255
 Ala Glu Ala Leu Gly Leu Ala Arg Arg Leu Tyr Gly Glu Gly Tyr Ala
 260 265 270
 Pro Arg Ser Leu Val Ser Gly Leu Leu Glu Val Phe Arg Glu Gly Leu
 275 280 285
 Tyr Ala Ala Phe Gly Leu Ala Gly Thr Pro Leu Pro Ala Pro Pro Gln
 290 295 300
 Ala Leu Ile Ala Ala Met Thr Ala Leu Asp Glu Ala Met Glu Arg Leu
 305 310 315 320
 Ala Arg Arg Ser Asp Ala Leu Ser Leu Glu Val Ala Leu Leu Glu Ala
 325 330 335
 Gly Arg Ala Leu Ala Ala Glu Ala Leu Pro Gln Pro Thr Gly Ala Pro
 340 345 350

05716964-112100

Ser Pro Glu Val Gly Pro Lys Pro Glu Ser Pro Pro Thr Pro Glu Pro
 355 360 365

Pro Arg Pro Glu Glu Ala Pro Asp Leu Arg Glu Arg Trp Arg Ala Phe
 370 375 380

Leu Glu Ala Leu Arg Pro Thr Leu Arg Ala Phe Val Arg Glu Ala Arg
 385 390 395 400

Pro Glu Val Arg Glu Gly Gln Leu Cys Leu Ala Phe Pro Glu Asp Lys
 405 410 415

Ala Phe His Tyr Arg Lys Ala Ser Glu Gln Lys Val Arg Leu Leu Pro
 420 425 430

Leu Ala Gln Ala His Phe Gly Val Glu Glu Val Val Leu Val Leu Glu
 435 440 445

Gly Glu Lys Lys Lys Pro Glu Pro Lys Ala Pro Pro Gly Pro Thr Ser
 450 455 460

<210> 5

<211> 454

<212> PRT

<213> Thermus thermophilus

<400> 5

Met Ser Ala Leu Tyr Arg Arg Phe Arg Pro Leu Thr Phe Gln Glu Val
 1 5 10 15

Val Gly Gln Glu His Val Lys Glu Pro Leu Leu Lys Ala Ile Arg Glu
 20 25 30

Gly Arg Leu Ala Gln Ala Tyr Leu Phe Ser Gly Pro Arg Gly Val Gly
 35 40 45

Lys Thr Thr Thr Ala Arg Leu Leu Ala Met Ala Val Gly Cys Gln Gly
 50 55 60

Glu Asp Pro Pro Cys Gly Val Cys Pro His Cys Gln Ala Val Gln Arg
 65 70 75 80

Gly Ala His Pro Asp Val Val Asp Ile Asp Ala Ala Ser Asn Asn Ser

09716964-112100

	85		90		95
Val Glu Asp Val Arg Glu Leu Arg Glu Arg Ile His Leu Ala Pro Leu	100		105		110
Ser Ala Pro Arg Lys Val Phe Ile Leu Asp Glu Ala His Met Leu Ser	115		120		125
Lys Ser Ala Phe Asn Ala Leu Leu Lys Thr Leu Glu Glu Pro Pro Pro	130		135		140
His Val Leu Phe Val Phe Ala Thr Thr Glu Pro Glu Arg Met Pro Pro	145		150		155
Thr Ile Leu Ser Arg Thr Gln His Phe Arg Phe Arg Arg Leu Thr Glu		165		170	175
Glu Glu Ile Ala Phe Lys Leu Arg Arg Ile Leu Glu Ala Val Gly Arg		180		185	190
Glu Ala Glu Glu Glu Ala Leu Leu Leu Leu Ala Arg Leu Ala Asp Gly		195		200	205
Ala Leu Arg Asp Ala Glu Ser Leu Leu Glu Arg Phe Leu Leu Leu Glu		210		215	220
Gly Pro Leu Thr Arg Lys Glu Val Glu Arg Ala Leu Gly Ser Pro Pro	225		230		235
Gly Thr Gly Val Ala Glu Ile Ala Ala Ser Leu Ala Arg Gly Lys Thr		245		250	255
Ala Glu Ala Leu Gly Leu Ala Arg Arg Leu Tyr Gly Glu Gly Tyr Ala		260		265	270
Pro Arg Ser Leu Val Ser Gly Leu Leu Glu Val Phe Arg Glu Gly Leu		275		280	285
Tyr Ala Ala Phe Gly Leu Ala Gly Thr Pro Leu Pro Ala Pro Pro Gln		290		295	300
Ala Leu Ile Ala Ala Met Thr Ala Leu Asp Glu Ala Met Glu Arg Leu	305		310		315
Ala Arg Arg Ser Asp Ala Leu Ser Leu Glu Val Ala Leu Leu Glu Ala		325		330	335
Gly Arg Ala Leu Ala Ala Glu Ala Leu Pro Gln Pro Thr Gly Ala Pro					

09716964.12100

340 345 350
Ser Pro Glu Val Gly Pro Lys Pro Glu Ser Pro Pro Thr Pro Glu Pro
355 360 365
Pro Arg Pro Glu Glu Ala Pro Asp Leu Arg Glu Arg Trp Arg Ala Phe
370 375 380
Leu Glu Ala Leu Arg Pro Thr Leu Arg Ala Phe Val Arg Glu Ala Arg
385 390 395 400
Pro Glu Val Arg Glu Gly Gln Leu Cys Leu Ala Phe Pro Glu Asp Lys
405 410 415
Ala Phe His Tyr Arg Lys Ala Ser Glu Gln Lys Val Arg Leu Leu Pro
420 425 430
Leu Ala Gln Ala His Phe Gly Val Glu Glu Val Val Leu Val Leu Glu
435 440 445
Gly Glu Lys Lys Lys Ala
450

<210> 6
<211> 32
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer

<400> 6
cgcaagcttc acgcstacct sttctccggs ac

32

<210> 7
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: peptide

<400> 7
His Ala Tyr Leu Phe Ser Gly Thr
1 5

<210> 8
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer

<400> 8
cgcggaattcg tgctcsggsg gctcctcsag sgtc

34

<210> 9
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: peptide

<400> 9
Lys Thr Leu Glu Glu Pro Pro Glu His
1 5

<210> 10
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer

<400> 10
gcgcggatcc ggagggagaa aaaaaaagcc tcagccca

38

<210> 11
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer

<400> 11
gcgcggatcc ggagggagag aagaaaagcc tcagccca

38

<210> 12
 <211> 28
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer

<400> 12
 gaattaaatt cgcgcttcgg gaggtggg 28

<210> 13
 <211> 27
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer

<400> 13
 gcgcgaattc gcgcttcggg aggtggg 27

<210> 14
 <211> 29
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer

<400> 14
 gcgcgaattc gggcgcttca ggaggtggg 29

<210> 15
 <211> 31
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer

<400> 15
 gtggtgcata tggtagcgcc cctctaccgc c 31

<210> 16
<211> 31
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer

<400> 16
gtggtggtcg acccaggagg gccacctcca g

31

<210> 17
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: peptide

<400> 17
Gly Xaa Xaa Gly Xaa Gly Lys Thr
1 5

<210> 18
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: peptide

<400> 18
Lys Pro Asp Pro Lys Ala Pro Pro Gly Pro Thr Ser
1 5 10

<210> 19
<211> 180
<212> PRT
<213> Escherichia coli

<400> 19
Met Ser Tyr Gln Val Leu Ala Arg Lys Trp Arg Pro Gln Thr Phe Ala
1 5 10 15

Asp Val Val Gly Gln Glu His Val Leu Thr Ala Leu Ala Asn Gly Leu
 20 25 30
 Ser Leu Gly Arg Ile His His Ala Tyr Leu Phe Ser Gly Thr Arg Gly
 35 40 45
 Val Gly Lys Thr Ser Ile Ala Arg Leu Leu Ala Lys Gly Leu Asn Cys
 50 55 60
 Glu Thr Gly Ile Thr Ala Thr Pro Cys Gly Val Cys Asp Asn Cys Arg
 65 70 75 80
 Glu Ile Glu Gln Gly Arg Phe Val Asp Leu Ile Glu Ile Asp Ala Ala
 85 90 95
 Ser Arg Thr Lys Val Glu Asp Thr Arg Asp Leu Leu Asp Asn Val Gln
 100 105 110
 Tyr Ala Pro Ala Arg Gly Arg Phe Lys Val Tyr Leu Ile Asp Glu Val
 115 120 125
 His Met Leu Ser Arg His Ser Phe Asn Ala Leu Leu Lys Thr Leu Glu
 130 135 140
 Glu Pro Pro Glu His Val Lys Phe Leu Leu Ala Thr Thr Asp Pro Gln
 145 150 155 160
 Lys Leu Pro Val Thr Ile Leu Ser Arg Cys Leu Gln Phe His Leu Lys
 165 170 175
 Ala Leu Asp Val
 180

<210> 20

<211> 180

<212> PRT

<213> Bacillus subtilis

<400> 20

Met Ser Tyr Gln Ala Leu Tyr Arg Val Phe Arg Pro Gln Arg Phe Glu
 1 5 10 15

Asp Val Val Gly Gln Glu His Ile Thr Lys Thr Leu Gln Asn Ala Leu
 20 25 30

Leu Gln Lys Lys Phe Ser His Ala Tyr Leu Phe Ser Gly Pro Arg Gly

09716964-112100

```

          35          40          45
Thr Gly Lys Thr Ser Ala Ala Lys Ile Phe Ala Lys Ala Val Asn Cys
  50          55          60

Glu His Ala Pro Val Asp Glu Pro Cys Asn Glu Cys Ala Ala Cys Lys
  65          70          75          80

Gly Ile Thr Asn Gly Ser Ile Ser Asp Val Ile Glu Ile Asp Ala Ala
          85          90          95

Ser Asn Asn Gly Val Asp Glu Ile Arg Asp Ile Arg Asp Lys Val Lys
          100          105          110

Phe Ala Pro Ser Ala Val Thr Tyr Lys Val Tyr Ile Ile Asp Glu Val
          115          120          125

His Met Leu Ser Ile Gly Ala Phe Asn Ala Leu Leu Lys Thr Leu Glu
          130          135          140

Glu Pro Pro Glu His Cys Ile Phe Ile Leu Ala Thr Thr Glu Pro His
          145          150          155          160

Lys Ile Pro Leu Thr Ile Ile Ser Arg Cys Gln Arg Phe Asp Phe Lys
          165          170          175

Arg Ile Thr Ser
          180

<210> 21
<211> 294
<212> PRT
<213> Escherichia coli

<400> 21
Met Ser Tyr Gln Val Leu Ala Arg Lys Trp Arg Pro Gln Thr Phe Ala
  1          5          10          15

Asp Val Val Gly Gln Glu His Val Leu Thr Ala Leu Ala Asn Gly Leu
          20          25          30

Ser Leu Gly Arg Ile His His Ala Tyr Leu Phe Ser Gly Thr Arg Gly
          35          40          45

Val Gly Lys Thr Ser Ile Ala Arg Leu Leu Ala Lys Gly Leu Asn Cys
          50          55          60
```

Glu Thr Gly Ile Thr Ala Thr Pro Cys Gly Val Cys Asp Asn Cys Arg
 65 70 75 80
 Glu Ile Glu Gln Gly Arg Phe Val Asp Leu Ile Glu Ile Asp Ala Ala
 85 90 95
 Ser Arg Thr Lys Val Glu Asp Thr Arg Asp Leu Leu Asp Asn Val Gln
 100 105 110
 Tyr Ala Pro Ala Arg Gly Arg Phe Lys Val Tyr Leu Ile Asp Glu Val
 115 120 125
 His Met Leu Ser Arg His Ser Phe Asn Ala Leu Leu Lys Thr Leu Glu
 130 135 140
 Glu Pro Pro Glu His Val Lys Phe Leu Leu Ala Thr Thr Asp Pro Gln
 145 150 155 160
 Lys Leu Pro Val Thr Ile Leu Ser Arg Cys Leu Gln Phe His Leu Lys
 165 170 175
 Ala Leu Asp Val Glu Gln Ile Arg His Gln Leu Glu His Ile Leu Asn
 180 185 190
 Glu Glu His Ile Ala His Glu Pro Arg Ala Leu Gln Leu Leu Ala Arg
 195 200 205
 Ala Ala Glu Gly Ser Leu Arg Asp Ala Leu Ser Leu Thr Asp Gln Ala
 210 215 220
 Ile Ala Ser Gly Asp Gly Gln Val Ser Thr Gln Ala Val Ser Ala Met
 225 230 235 240
 Leu Gly Thr Leu Asp Asp Asp Gln Ala Leu Ser Leu Val Glu Ala Met
 245 250 255
 Val Glu Ala Asn Gly Glu Arg Val Met Ala Leu Ile Asn Glu Ala Ala
 260 265 270
 Ala Arg Gly Ile Glu Trp Glu Ala Leu Leu Val Glu Met Leu Gly Leu
 275 280 285
 Leu His Arg Ile Ala Met
 290

<210> 22

<211> 294

<212> PRT

<213> Haemophilus influenzae

<400> 22

Met Ser Tyr Gln Val Leu Ala Arg Lys Trp Arg Pro Lys Thr Phe Ala
1 5 10 15

Asp Val Val Gly Gln Glu His Ile Ile Thr Ala Leu Ala Asn Gly Leu
20 25 30

Lys Asp Asn Arg Leu His His Ala Tyr Leu Phe Ser Gly Thr Arg Gly
35 40 45

Val Gly Lys Thr Ser Ile Ala Arg Leu Phe Ala Lys Gly Leu Asn Cys
50 55 60

Val His Gly Val Thr Ala Thr Pro Cys Gly Glu Cys Glu Asn Cys Lys
65 70 75 80

Ala Ile Glu Gln Gly Asn Phe Ile Asp Leu Ile Glu Ile Asp Ala Ala
85 90 95

Ser Arg Thr Lys Val Glu Asp Thr Arg Glu Leu Leu Asp Asn Val Gln
100 105 110

Tyr Lys Pro Val Val Gly Arg Phe Lys Val Tyr Leu Ile Asp Glu Val
115 120 125

His Met Leu Ser Arg His Ser Phe Asn Ala Leu Leu Lys Thr Leu Glu
130 135 140

Glu Pro Pro Glu Tyr Val Lys Phe Leu Leu Ala Thr Thr Asp Pro Gln
145 150 155 160

Lys Leu Pro Val Thr Ile Leu Ser Arg Cys Leu Gln Phe His Leu Lys
165 170 175

Ala Leu Asp Glu Thr Gln Ile Ser Gln His Leu Ala His Ile Leu Thr
180 185 190

Gln Glu Asn Ile Pro Phe Glu Asp Pro Ala Leu Val Lys Leu Ala Lys
195 200 205

Ala Ala Gln Gly Ser Ile Arg Asp Ser Leu Ser Leu Thr Asp Gln Ala
210 215 220

Ile Ala Met Gly Asp Arg Gln Val Thr Asn Asn Val Val Ser Asn Met
225 230 235 240

Leu Gly Leu Leu Asp Asp Asn Tyr Ser Val Asp Ile Leu Tyr Ala Leu
 245 250 255

His Gln Gly Asn Gly Glu Leu Leu Met Arg Thr Leu Gln Arg Val Ala
 260 265 270

Asp Ala Ala Gly Asp Trp Asp Lys Leu Leu Gly Glu Cys Ala Glu Lys
 275 280 285

Leu His Gln Ile Ala Leu
 290

<210> 23

<211> 294

<212> PRT

<213> Bacillus subtilis

<400> 23

Met Ser Tyr Gln Ala Leu Tyr Arg Val Phe Arg Pro Gln Arg Phe Glu
 1 5 10 15

Asp Val Val Gly Gln Glu His Ile Thr Lys Thr Leu Gln Asn Ala Leu
 20 25 30

Leu Gln Lys Lys Phe Ser His Ala Tyr Leu Phe Ser Gly Pro Arg Gly
 35 40 45

Thr Gly Lys Thr Ser Ala Ala Lys Ile Phe Ala Lys Ala Val Asn Cys
 50 55 60

Glu His Ala Pro Val Asp Glu Pro Cys Asn Glu Cys Ala Ala Cys Lys
 65 70 75 80

Gly Ile Thr Asn Gly Ser Ile Ser Asp Val Ile Glu Ile Asp Ala Ala
 85 90 95

Ser Asn Asn Gly Val Asp Glu Ile Arg Asp Ile Arg Asp Lys Val Lys
 100 105 110

Phe Ala Pro Ser Ala Val Thr Tyr Lys Val Tyr Ile Ile Asp Glu Val
 115 120 125

His Met Leu Ser Ile Gly Ala Phe Asn Ala Leu Leu Lys Thr Leu Glu
 130 135 140

Glu Pro Pro Glu His Cys Ile Phe Ile Leu Ala Thr Thr Glu Pro His

145 150 155 160

Lys Ile Pro Leu Thr Ile Ile Ser Arg Cys Gln Arg Phe Asp Phe Lys
 165 170 175

Arg Ile Thr Ser Gln Ala Ile Val Gly Arg Met Asn Lys Ile Val Asp
 180 185 190

Ala Glu Gln Leu Gln Val Glu Glu Gly Ser Leu Glu Ile Ile Ala Ser
 195 200 205

Ala Ala His Gly Gly Met Arg Asp Ala Leu Ser Leu Leu Asp Gln Ala
 210 215 220

Ile Ser Phe Ser Gly Asp Ile Leu Lys Val Glu Asp Ala Leu Leu Ile
225 230 235 240

Thr Gly Ala Val Ser Gln Leu Tyr Ile Gly Lys Leu Ala Lys Ser Leu
 245 250 255

His Asp Lys Asn Val Ser Asp Ala Leu Glu Thr Leu Asn Glu Leu Leu
 260 265 270

Gln Gln Gly Lys Asp Pro Ala Lys Leu Ile Glu Asp Met Ile Phe Tyr
 275 280 285

Phe Arg Asp Met Leu Leu
 290

<210> 24
<211> 300
<212> PRT
<213> *Caulobacter crescentus*

<400> 24
Asp Ala Tyr Thr Val Leu Ala Arg Lys Tyr Arg Pro Arg Thr Phe Glu
1 5 10 15

Asp Leu Ile Gly Gln Glu Ala Met Val Arg Thr Leu Ala Asn Ala Phe
 20 25 30

Ser Thr Gly Arg Ile Ala His Ala Phe Met Leu Thr Gly Val Arg Gly
 35 40 45

Val Gly Lys Thr Thr Thr Ala Arg Leu Leu Ala Arg Ala Leu Asn Tyr
 50 55 60

Glu Thr Asp Thr Val Lys Gly Pro Ser Val Asp Leu Thr Thr Glu Gly
 65 70 75 80
 Tyr His Cys Arg Ser Ile Ile Glu Gly Arg His Met Asp Val Leu Glu
 85 90 95
 Leu Asp Ala Ala Ser Arg Thr Lys Val Asp Glu Met Arg Glu Leu Leu
 100 105 110
 Asp Gly Val Arg Tyr Ala Pro Val Glu Ala Arg Tyr Lys Val Tyr Ile
 115 120 125
 Ile Asp Glu Val His Met Leu Ser Thr Ala Ala Phe Asn Ala Leu Leu
 130 135 140
 Lys Thr Leu Glu Glu Pro Pro Pro His Ala Lys Phe Ile Phe Ala Thr
 145 150 155 160
 Thr Glu Ile Arg Lys Val Pro Val Thr Ile Leu Ser Arg Cys Gln Arg
 165 170 175
 Phe Asp Leu Arg Arg Val Glu Pro Asp Val Leu Val Lys His Phe Asp
 180 185 190
 Arg Ile Ser Ala Lys Glu Gly Ala Arg Ile Glu Met Asp Ala Leu Ala
 195 200 205
 Leu Ile Ala Arg Ala Ala Glu Gly Ser Val Arg Asp Gly Leu Ser Leu
 210 215 220
 Leu Asp Gln Ala Ile Val Gln Thr Glu Arg Gly Gln Thr Val Thr Ser
 225 230 235 240
 Thr Val Val Arg Asp Met Leu Gly Leu Ala Asp Arg Ser Gln Thr Ile
 245 250 255
 Ala Leu Tyr Glu His Val Met Ala Gly Lys Thr Lys Asp Ala Leu Glu
 260 265 270
 Gly Phe Arg Ala Leu Trp Gly Phe Gly Ala Asp Pro Ala Val Val Met
 275 280 285
 Leu Asp Val Leu Asp His Cys His Ala Ser Ala Val
 290 295 300

<210> 25
 <211> 260

<212> PRT

<213> Mycoplasma genitalium

<400> 25

Met His Gln Val Phe Tyr Gln Lys Tyr Arg Pro Ile Asn Phe Lys Gln
1 5 10 15

Thr Leu Gly Gln Glu Ser Ile Arg Lys Ile Leu Val Asn Ala Ile Asn
20 25 30

Arg Asp Lys Leu Pro Asn Gly Tyr Ile Phe Ser Gly Glu Arg Gly Thr
35 40 45

Gly Lys Thr Thr Phe Ala Lys Ile Ile Ala Lys Ala Ile Asn Cys Leu
50 55 60

Asn Trp Asp Gln Ile Asp Val Cys Asn Ser Cys Asp Val Cys Lys Ser
65 70 75 80

Ile Asn Thr Asn Ser Ala Ile Asp Ile Val Glu Ile Asp Ala Ala Ser
85 90 95

Lys Asn Gly Ile Asn Asp Ile Arg Glu Leu Val Glu Asn Val Phe Asn
100 105 110

His Pro Phe Thr Phe Lys Lys Lys Val Tyr Ile Leu Asp Glu Ala His
115 120 125

Met Leu Thr Thr Gln Ser Trp Gly Gly Leu Leu Lys Thr Leu Glu Glu
130 135 140

Ser Pro Pro Tyr Val Leu Phe Ile Phe Thr Thr Thr Glu Phe Asn Lys
145 150 155 160

Ile Pro Leu Thr Ile Leu Ser Arg Cys Gln Ser Phe Phe Phe Lys Lys
165 170 175

Ile Thr Ser Asp Leu Ile Leu Glu Arg Leu Asn Asp Ile Ala Lys Lys
180 185 190

Glu Lys Ile Lys Ile Glu Lys Asp Ala Leu Ile Lys Ile Ala Asp Leu
195 200 205

Ser Gln Gly Ser Leu Arg Asp Gly Leu Ser Leu Leu Asp Gln Leu Ala
210 215 220

Ile Ser Leu Ile Val Lys Lys Leu Val Leu Leu Met Leu Lys Lys His
225 230 235 240

180	185	190
Glu Ala Glu Glu Glu Ala Leu Leu Leu Leu Ala Arg Leu Ala Asp Gly		
195	200	205
Ala Leu Arg Asp Ala Glu Ser Leu Leu Glu Arg Phe Leu Leu Leu Glu		
210	215	220
Gly Pro Leu Thr Arg Lys Glu Val Glu Arg Ala Leu Gly Ser Pro Pro		
225	230	235 240
Gly Thr Gly Val Ala Glu Ile Ala Ala Ser Leu Ala Arg Gly Lys Thr		
245	250	255
Ala Glu Ala Leu Gly Leu Ala Arg Arg Leu Tyr Gly Glu Gly Tyr Ala		
260	265	270
Pro Arg Ser Leu Val Ser Gly Leu Leu Glu Val Phe Arg Glu Gly Leu		
275	280	285

Tyr

<210> 27
 <211> 94
 <212> DNA
 <213> Thermus thermophilus

<400> 27
 gccggaggga gaaaaaaaaa gccgagccca agggcccgcc cggccccacc ccgaagcgcc 60
 cgcacccccg ggccccccga ggaggaggag aggc 94

<210> 28
 <211> 11
 <212> PRT
 <213> Thermus thermophilus

<400> 28
 Val Leu Glu Gly Glu Lys Lys Ser Leu Ser Pro
 1 5 10

<210> 29
 <211> 23
 <212> DNA
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 29

cacgcntacc tnttctccgg nac

23

<210> 30

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 30

gtgctcnggn ggctcctcnt cngtc

25

<210> 31

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 31

gtgggatccg tggttctgga tctcgatgaa gaa

33

<210> 32

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 32

gtgggatcca cggststcs gacgagaag

29

<210> 33

<211> 34

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 33

gcgggatcct caacgaggac ctctccatct tcaa

34

<210> 34

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 34

gcgggatcct tgtcgtsag sgtsagsgcg tcgta

35

<210> 35

<211> 39

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 35

gggaaggacc agcgcgtact cccctgctc ctagggtg

39

<210> 36

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 36

gtgtggatcc ttcttcttsc ccatsgc

27

<210> 37

<211> 27

<212> DNA

<213> Artificial Sequence

[illegible]

	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
0	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100

[illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible][illegible]

<220>

<223> Description of Artificial Sequence: primer

<400> 44

cggggatcca cctcaatcac ctcggtgg

27

<210> 45

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 45

cggggatccg ccaccttgcg gctccgggtg

30

<210> 46

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 46

gcgctctaga cgagttccca aagcgtgcgg t

31

<210> 47

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 47

cgcgtctaga tcacctgtat ccaga

25

<210> 48

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 48

gcggcgcata tgggtggtggt cctggacctg gag

33

<210> 49

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 49

cgcgctctaga tcacctgtat ccaga

25

<210> 50

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 50

gtsctsgtsa agacscactt

20

<210> 51

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 51

sagsagsgcg ttgaasgtgt g

21

<210> 52

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 56

catcctgaag atgaacgcca gca

23

<210> 57

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 57

agggttatcca caggggtcat gtgca

25

<210> 58

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 58

gtgtgtcata tgaacataac ggttcctcaa

29

<210> 59

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 59

gcgcgaattc tcccttgtgg aaggcttag

29

<210> 60

<211> 13

<212> PRT

<213> Thermus thermophilus

<400> 60

Arg Val Glu Leu Asp Tyr Asp Ala Leu Thr Leu Asp Asp
1 5 10

<210> 61

<211> 14

<212> PRT

<213> *Thermus thermophilus*

<400> 61

Phe Phe Ile Glu Ile Gln Asn His Gly Leu Ser Glu Gln Lys
1 5 10

<210> 62

<211> 8

<212> PRT

<213> *Thermus thermophilus*

<400> 62

Phe Phe Ile Glu Ile Gln Asn His
1 5

<210> 63

<211> 8

<212> PRT

<213> *Thermus thermophilus*

<400> 63

Tyr Asp Ala Leu Thr Leu Asp Asp
1 5

<210> 64

<211> 6

<212> PRT

<213> *Thermus thermophilus*

<400> 64

Ala Met Gly Lys Lys Lys
1 5

<210> 65

<211> 9

<212> PRT

<213> Thermus thermophilus

<400> 65

Phe Asn Lys Ser His Ser Ala Ala Tyr

1

5

<210> 66

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: peptide

<400> 66

Val Val Xaa Asp Xaa Glu Thr Thr Gly

1

5

<210> 67

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: peptide

<400> 67

His Asn Ala Xaa Phe Asp Xaa Gly Phe

1

5

<210> 68

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: peptide

<400> 68

Val Val Xaa Asp Xaa Glu Thr Thr Gly

1

5

<210> 69

<211> 7

<212> PRT

<213> *Thermus thermophilus*

<400> 69

Val Leu Val Lys Thr His Leu

1

5

<210> 70

<211> 6

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: peptide

<400> 70

His Arg Ala Leu Tyr Asp

1

5

<210> 71

<211> 7

<212> PRT

<213> *Thermus thermophilus*

<400> 71

His Thr Phe Asn Ala Leu Leu

1

5

<210> 72

<211> 34

<212> PRT

<213> *Escherichia coli*

<400> 72

Asp Arg Tyr Phe Leu Glu Leu Ile Arg Thr Gly Arg Pro Asp Glu Glu

1

5

10

15

Ser Tyr Leu His Ala Ala Val Glu Leu Ala Glu Ala Arg Gly Leu Pro

20

25

30

Val Val

<210> 73
 <211> 34
 <212> PRT
 <213> *Vibrio cholerae*

<400> 73
 Asp His Phe Tyr Leu Glu Leu Ile Arg Thr Gly Arg Ala Asp Glu Glu
 1 5 10 15
 Ser Tyr Leu His Phe Ala Leu Asp Val Ala Glu Gln Tyr Asp Leu Pro
 20 25 30
 Val Val

<210> 74
 <211> 34
 <212> PRT
 <213> *Haemophilus influenzae*

<400> 74
 Asp His Phe Tyr Leu Ala Leu Ser Arg Thr Gly Arg Pro Asn Glu Glu
 1 5 10 15
 Arg Tyr Ile Gln Ala Ala Leu Lys Leu Ala Glu Arg Cys Asp Leu Pro
 20 25 30
 Leu Val

<210> 75
 <211> 34
 <212> PRT
 <213> *Rickettsia prowazekii*

<400> 75
 Asp Arg Phe Tyr Phe Glu Ile Met Arg His Asp Leu Pro Glu Glu Gln
 1 5 10 15
 Phe Ile Glu Asn Ser Tyr Ile Gln Ile Ala Ser Glu Leu Ser Ile Pro
 20 25 30
 Ile Val

<210> 76
 <211> 34
 <212> PRT
 <213> *Helicobacter pylori*

<400> 76
 Asp Asp Phe Tyr Leu Glu Ile Met Arg His Gly Ile Leu Asp Gln Arg
 1 5 10 15
 Phe Ile Asp Glu Gln Val Ile Lys Met Ser Leu Glu Thr Gly Leu Lys
 20 25 30
 Ile Ile

<210> 77
 <211> 34
 <212> PRT
 <213> *Synechocystis* sp.

<400> 77
 Asp Asp Tyr Tyr Leu Glu Ile Gln Asp His Gly Ser Val Glu Asp Arg
 1 5 10 15
 Leu Val Asn Ile Asn Leu Val Lys Ile Ala Gln Glu Leu Asp Ile Lys
 20 25 30
 Ile Val

<210> 78
 <211> 34
 <212> PRT
 <213> *Mycobacterium tuberculosis*

<400> 78
 Asp Asn Tyr Phe Leu Glu Leu Met Asp His Gly Leu Thr Ile Glu Arg
 1 5 10 15
 Arg Val Arg Asp Gly Leu Leu Glu Ile Gly Arg Ala Leu Asn Ile Pro
 20 25 30
 Pro Leu

<210> 79
 <211> 46
 <212> PRT
 <213> Escherichia coli

<400> 79
 Asn Lys Arg Arg Ala Lys Asn Gly Glu Pro Pro Leu Asp Ile Ala Ala
 1 5 10 15
 Ile Pro Leu Asp Asp Lys Lys Ser Phe Asp Met Leu Gln Arg Ser Glu
 20 25 30
 Thr Thr Ala Val Phe Gln Leu Glu Ser Arg Gly Met Lys Asp
 35 40 45

<210> 80
 <211> 46
 <212> PRT
 <213> Vibrio cholerae

<400> 80
 Asn Pro Arg Leu Lys Lys Ala Gly Lys Pro Pro Val Arg Ile Glu Ala
 1 5 10 15
 Ile Pro Leu Asp Asp Ala Arg Ser Phe Arg Asn Leu Gln Asp Ala Lys
 20 25 30
 Thr Thr Ala Val Phe Gln Leu Glu Ser Arg Gly Met Lys Glu
 35 40 45

<210> 81
 <211> 46
 <212> PRT
 <213> Haemophilus influenzae

<400> 81
 Asn Val Arg Met Val Arg Glu Gly Lys Pro Arg Val Asp Ile Ala Ala
 1 5 10 15
 Ile Pro Leu Asp Asp Pro Glu Ser Phe Glu Leu Leu Lys Arg Ser Glu
 20 25 30
 Thr Thr Ala Val Phe Gln Leu Glu Ser Arg Gly Met Lys Asp
 35 40 45

<210> 85
 <211> 46
 <212> PRT
 <213> Mycobacterium tuberculosis

<400> 85
 Ile Asp Asn Val Arg Ala Asn Arg Gly Ile Asp Leu Asp Leu Glu Ser
 1 5 10 15
 Val Pro Leu Asp Asp Lys Ala Thr Tyr Glu Leu Leu Gly Arg Gly Asp
 20 25 30
 Thr Leu Gly Val Phe Gln Leu Asp Gly Gly Pro Met Arg Asp
 35 40 45

<210> 86
 <211> 3729
 <212> DNA
 <213> Thermus thermophilus

<400> 86
 atgggcccggg agctccgctt cgcccacctc caccagcaca cccagttctc cctcctggac 60
 ggggcccgcga agctttccga cctcctcaag tgggtcaagg agaagacccc cgaggacccc 120
 gccttggcca tgaccgacca cggcaacctc ttcggggccg tggagttcta caagaaggcc 180
 accgaaatgg gcatcaagcc catcctgggc tacgaggcct acgtggcggc ggaaagccgc 240
 tttgaccgca agcggggaaa gggcctagac gggggctact ttcacctcac cctcctcgcc 300
 aaggacttca cgggggtacca gaacctggtg cgcctggcga gccgggctta cctggagggg 360
 ttttacgaaa agccccgat tgaccgggag atcctgcgcg agcacgccga gggcctcatc 420
 gccctctcgg ggtgcctcgg ggcggagatc ccccagttca tcctccagga ccgtctggac 480
 ctggccgagg cccggctcaa cgagtacctc tccatcttca aggaccgctt ctcatcgag 540
 atccagaacc acggcctccc cgagcagaaa aaggtcaacg aggtcctcaa ggagttcgcc 600
 cgaaagtacg gcctggggat ggtggccacc aacgacggcc attacgtgag gaaggaggac 660
 gcccgcgccc acgaggtcct cctcgccatc cagtccaaga gcacctgga cgaccccggg 720
 cgctggcgct tcccctgcga cgagttctac gtgaagaccc ccgaggagat gcggggccatg 780
 ttccccgagg aggagtgggg ggacgagccc tttgacaaca ccgtggagat cgcccgcgatg 840
 tgcaacgtgg agctgcccac cggggacaag atggtctacc gaatcccccg cttccccctc 900
 cccgagggggc ggaccgaggc ccagtacctc atggagctca cttcaaggg gctcctccgc 960
 cgctacccgg accggatcac cgagggcttc taccgggagg tcttcgcct tttggggaag 1020
 cttccccccc acggggacgg ggaggccttg gccgaggcct tggcccagg gtgacggcg 1080
 gcttgggaga ggctcatgaa gagcctcccc cctttggccg ggtcaagga gtggacggcg 1140
 gaggccattt tccaccgggc cctttacgag ctttcctgta tagagcgcac ggggtttccc 1200
 ggctacttcc tcatcgtcca ggactacatc aactgggccc ggagaaacgg cgtctccgtg 1260
 gggcccggca gggggagcgc cgccgggagc ctggtggcct acgccgtggg gatcaccaac 1320
 attgaccccc tccgcttcgg cctcctcttt gagcgcttcc tgaaccggga gagggctctc 1380
 atgcccgcga ttgacacgga cttctccgac cgggagcggg accgggtgat ccagtacgtg 1440
 cgggagcgct acggcgagga caaggtggcc cagatcgcca ccctgggaag cctcgccctc 1500
 aaggccgccc tcaaggacgt ggcccgggtc tacggcatcc cccacaagaa ggccgaggaa 1560

```

ttggccaagc tcatcccggg gcagttcggg aagcccaagc ccctgcagga ggccatccag 1620
gtggtgccgg agcttagggc ggagatggag aaggacccca aggtgcggga ggtcctcgag 1680
gtggccatgc gcctggaggg cctgaaccgc cagcctccg tccacgccgc cggggtgggtg 1740
atcgccgccc agccccctac ggacctcgtc cccctcatgc gcgaccagga agggcgcccc 1800
gtcaccaggt acgacatggg ggcgggtggg gccttggggc ttttgaagat ggactttttg 1860
ggcctccgca ccctcacctt cctggacgag gtcaagcgca tcgtcaaggc gtcccagggg 1920
gtggagctgg actacgatgc cctccccctg gacgacccca agaccttcgc cctcctctcc 1980
cggggggaga ccaagggggg cttccagctg gagtcggggg ggatgaccgc cagctccgc 2040
ggcctcaagc cgcggcgctt tgaggacctg atcgccatcc tctccctcta ccgccccggg 2100
cccatggagc acatccccac ctacatccgc cgccaccacg ggctggagcc cgtgagctac 2160
agcgagtttc cccacgccga gaagtacctt aagcccatcc tggacgagac ctacggcatc 2220
cccgctctacc aggagcagat catgcagatc gcctcggccg tggcggggta ctccctgggc 2280
gaggcggacc tcctgcggcg gtccatgggc aagaagaagg tggaggagat gaagtccac 2340
cgggagcgct tcgtccaggg ggccaaggaa aggggcgtgc ccgaggagga ggccaaccgc 2400
ctctttgaca tgctggaggg cttcgccaac tacggcttca acaaatccca cgctgccgcc 2460
tacagcctcc tctcctacca gaccgcctac gtgaaggccc actaccctgt ggagttcatg 2520
gccgccctcc tctcctgga gcggcacgac tccgacaagg tggccgagta catccgcgac 2580
gcccgggcca tgggcataga ggtccttccc ccggacgtca accgctccgg gtttgacttc 2640
ctggtccagg gccggcagat ccttttcggc ctctccgcgg tgaagaacgt gggcgaggcg 2700
gcggcgagg ccattctccg ggagcgggag cggggcgggc cctaccggag cctcggcgac 2760
ttcctcaagc ggctggacga gaaggtgtc aacaagcgga ccctggagtc cctcatcaag 2820
gcgggcgccc tggacggctt cggggaaaag gcgcggctcc tcgcctccct ggaagggtc 2880
ctcaagtggg cggccgagaa ccgggagaag gcccgcctcg gcacgatggg cctcttcagc 2940
gaagtggagg agccgccttt ggccgaggcc gccccctgg acgagatcac ccggctccgc 3000
tacgagaagg aggcctggg gatctacgtc tccggccacc ccattctgg gtaccccggg 3060
ctccgggaga cggccacctg caccctggag gagcttccc acctggccc ggacctgccg 3120
ccccggtcta gggctctcct tgccgggatg gtggaggagg tgggtgcgca gcccacaaag 3180
agcggcgagg tgatggccc cttcgtcctc tccgacgaga cgggggcgct tgaggcggtg 3240
gcattcggcc gggcctacga ccaggtctcc ccgaggctca aggaggacac ccccgctgc 3300
gtcctcgccg aggtggagcg ggaggagggg ggcgtgcggg tgctggccca ggccgtttg 3360
acctacgagg agctggagca ggtcccccg gccctcgagg tggaggtgga ggcctccctc 3420
ctggacgacc ggggggtggc ccacctgaaa agcctcctgg acgagcacgc ggggacctc 3480
ccccgtacg tccgggtcca gggcgcttc ggcgaggccc tcctcgccct gagggaggtg 3540
cgggtggggg aggaggctgt aggcggccgc gtggttcgg gcctacctcc tgcccgaccg 3600
ggaggctcct ctccagggcg gccaggcggg ggaggcccag gaggcggtgc ccttctaggg 3660
ggtgggcccgt gagacctagc gccatcgttc tcgccggggg caaggaggcc tgggcccgc 3720
cccttttgg 3729

```

<210> 87

<211> 1245

<212> PRT

<213> *Thermus thermophilus*

<400> 87

Met Gly Arg Glu Leu Arg Phe Ala His Leu His Gln His Thr Gln Phe

1

5

10

15

Ser Leu Leu Asp Gly Ala Pro Lys Leu Ser Asp Leu Leu Lys Trp Val
 20 25 30
 Glu Glu Thr Thr Pro Glu Asp Pro Ala Leu Ala Met Thr Asp His Gly
 35 40 45
 Asn Leu Phe Gly Ala Val Glu Phe Tyr Lys Lys Ala Thr Glu Met Gly
 50 55 60
 Ile Lys Pro Ile Leu Gly Tyr Glu Ala Tyr Val Ala Ala Glu Ser Arg
 65 70 75 80
 Phe Asp Arg Lys Arg Gly Lys Gly Leu Asp Gly Gly Tyr Phe His Leu
 85 90 95
 Thr Leu Leu Ala Lys Asp Phe Thr Gly Tyr Gln Asn Leu Val Arg Leu
 100 105 110
 Ala Ser Arg Ala Tyr Leu Glu Gly Phe Tyr Glu Lys Pro Arg Ile Asp
 115 120 125
 Arg Glu Ile Leu Arg Glu His Ala Glu Gly Leu Ile Ala Leu Ser Gly
 130 135 140
 Cys Leu Gly Ala Glu Ile Pro Gln Phe Ile Leu Gln Asp Arg Leu Asp
 145 150 155 160
 Leu Ala Glu Ala Arg Leu Asn Glu Tyr Leu Ser Ile Phe Lys Asp Arg
 165 170 175
 Phe Phe Ile Glu Ile Gln Asn His Gly Leu Pro Glu Gln Lys Lys Val
 180 185 190
 Asn Glu Val Leu Lys Glu Phe Ala Arg Lys Tyr Gly Leu Gly Met Val
 195 200 205
 Ala Thr Asn Asp Gly His Tyr Val Arg Lys Glu Asp Ala Arg Ala His
 210 215 220
 Glu Val Leu Leu Ala Ile Gln Ser Lys Ser Thr Leu Asp Asp Pro Gly
 225 230 235 240
 Ala Leu Ala Leu Pro Cys Glu Glu Phe Tyr Val Lys Thr Pro Glu Glu
 245 250 255
 Met Arg Ala Met Phe Pro Glu Glu Glu Val Gly Gly Arg Ser Pro Leu
 260 265 270

Thr Thr Pro Trp Arg Ser Pro His Val Gln Arg Gly Ala Ala Ile Gly
 275 280 285

Thr Arg Trp Ser Thr Arg Ile Pro Arg Phe Pro Leu Pro Glu Gly Arg
 290 295 300

Thr Glu Ala Gln Tyr Leu Met Glu Leu Thr Phe Lys Gly Leu Leu Arg
 305 310 315 320

Arg Tyr Pro Asp Arg Ile Thr Glu Gly Phe Tyr Arg Glu Val Phe Arg
 325 330 335

Leu Ser Gly Lys Leu Pro Pro His Gly Asp Gly Glu Ala Leu Ala Glu
 340 345 350

Ala Leu Ala Gln Val Glu Arg Glu Ala Trp Glu Arg Leu Met Lys Ser
 355 360 365

Leu Pro Pro Leu Ala Gly Val Lys Glu Trp Thr Ala Glu Ala Ile Phe
 370 375 380

His Arg Ala Leu Tyr Glu Leu Ser Ala Ile Glu Arg Met Gly Phe Pro
 385 390 395 400

Gly Leu Leu Pro His Arg Pro Gly Leu His Gln Leu Gly Pro Glu Lys
 405 410 415

Gly Val Ser Val Gly Pro Gly Arg Gly Gly Ala Ala Gly Ser Leu Val
 420 425 430

Ala Tyr Ala Val Gly Ile Thr Asn Ile Asp Pro Leu Arg Phe Gly Leu
 435 440 445

Leu Phe Glu Arg Phe Leu Asn Pro Glu Arg Val Ser Met Pro Asp Ile
 450 455 460

Asp Thr Asp Phe Ser Asp Arg Glu Arg Asp Arg Val Ile Gln Tyr Val
 465 470 475 480

Arg Glu Arg Tyr Gly Glu Asp Lys Val Ala Gln Ile Gly Thr Leu Gly
 485 490 495

Ser Leu Ala Ser Lys Ala Ala Leu Lys Glu Val Ala Arg Val Tyr Gly
 500 505 510

Ile Pro Arg Lys Lys Ala Glu Glu Leu Ala Lys Leu Ile Pro Val Gln
 515 520 525

Val Gln Gly Ala Lys Glu Arg Gly Val Pro Glu Glu Glu Ala Asn Arg
 785 790 795 800
 Leu Phe Asp Met Leu Glu Ala Phe Ala Asn Tyr Gly Phe Asn Lys Ser
 805 810 815
 His Ala Ala Ala Tyr Ser Leu Leu Ser Tyr Gln Thr Ala Tyr Val Lys
 820 825 830
 Ala His Tyr Pro Val Glu Phe Met Ala Ala Leu Leu Ser Val Glu Arg
 835 840 845
 His Asp Ser Asp Lys Val Ala Glu Tyr Ile Arg Asp Ala Arg Ala Met
 850 855 860
 Gly Ile Glu Val Leu Pro Pro Asp Val Asn Arg Ser Gly Phe Asp Phe
 865 870 875 880
 Leu Val Gln Gly Arg Gln Ile Leu Phe Gly Leu Ser Ala Val Lys Asn
 885 890 895
 Val Gly Glu Ala Ala Ala Glu Ala Ile Leu Arg Glu Arg Glu Arg Gly
 900 905 910
 Gly Pro Tyr Arg Ser Leu Gly Asp Phe Leu Lys Arg Leu Asp Glu Lys
 915 920 925
 Val Leu Asn Lys Arg Thr Leu Glu Ser Leu Ile Lys Ala Gly Ala Leu
 930 935 940
 Asp Gly Phe Gly Glu Arg Ala Arg Leu Leu Ala Ser Leu Glu Gly Leu
 945 950 955 960
 Leu Lys Trp Ala Ala Glu Asn Arg Glu Lys Ala Arg Ser Gly Met Met
 965 970 975
 Gly Leu Phe Ser Glu Val Glu Glu Pro Pro Leu Ala Glu Ala Ala Pro
 980 985 990
 Leu Asp Glu Ile Thr Arg Leu Arg Tyr Glu Lys Glu Ala Leu Gly Ile
 995 1000 1005
 Tyr Val Ser Gly His Pro Ile Leu Arg Tyr Pro Gly Leu Arg Glu Thr
 1010 1015 1020
 Ala Thr Cys Thr Leu Glu Glu Leu Pro His Leu Ala Arg Asp Leu Pro
 1025 1030 1035 1040

Pro Arg Ser Arg Val Leu Leu Ala Gly Met Val Glu Glu Val Val Arg
 1045 1050 1055

Lys Pro Thr Lys Ser Gly Gly Met Met Ala Arg Phe Val Leu Ser Asp
 1060 1065 1070

Glu Thr Gly Ala Leu Glu Ala Val Ala Phe Gly Arg Ala Tyr Asp Gln
 1075 1080 1085

Val Ser Pro Arg Leu Lys Glu Asp Thr Pro Val Leu Val Leu Ala Glu
 1090 1095 1100

Val Glu Arg Glu Glu Gly Gly Val Arg Val Leu Ala Gln Ala Val Trp
 1105 1110 1115 1120

Thr Tyr Gln Glu Leu Glu Gln Val Pro Arg Ala Leu Glu Val Glu Val
 1125 1130 1135

Glu Ala Ser Leu Pro Asp Asp Arg Gly Val Ala His Leu Lys Ser Leu
 1140 1145 1150

Leu Asp Glu His Ala Gly Thr Leu Pro Leu Tyr Val Arg Val Gln Gly
 1155 1160 1165

Ala Phe Gly Glu Ala Leu Leu Ala Leu Arg Glu Val Arg Val Gly Glu
 1170 1175 1180

Glu Ala Leu Gly Ala Leu Glu Ala Ala Gly Phe Pro Ala Tyr Leu Leu
 1185 1190 1195 1200

Pro Asn Arg Glu Val Ser Pro Arg Leu Thr Gly Ser Gly Gly Pro Arg
 1205 1210 1215

Gly Arg Ala Leu Ser Thr Gly Leu Ala Leu Lys Thr Tyr Pro Ile Ala
 1220 1225 1230

Leu Pro Gly Gly Asn Glu Ala Leu Ala Arg Pro Leu Leu
 1235 1240 1245

<210> 88

<211> 198

<212> PRT

<213> Thermus thermophilus

<400> 88

Val Glu Arg Val Val Arg Thr Leu Leu Asp Gly Arg Phe Leu Leu Glu
 1 5 10 15

Glu Gly Val Gly Leu Trp Glu Trp Arg Tyr Pro Phe Pro Leu Glu Gly
 20 25 30

Glu Ala Val Val Val Leu Asp Leu Glu Thr Thr Gly Leu Ala Gly Leu
 35 40 45

Asp Glu Val Ile Glu Val Gly Leu Leu Arg Leu Glu Gly Gly Arg Arg
 50 55 60

Leu Pro Phe Gln Ser Leu Val Arg Pro Leu Pro Pro Ala Glu Ala Arg
 65 70 75 80

Ser Trp Asn Leu Thr Gly Ile Pro Arg Glu Ala Leu Glu Glu Ala Pro
 85 90 95

Ser Leu Glu Glu Val Leu Glu Lys Ala Tyr Pro Leu Arg Gly Asp Ala
 100 105 110

Thr Leu Val Ile His Asn Ala Ala Phe Asp Leu Gly Phe Leu Arg Pro
 115 120 125

Ala Leu Glu Gly Leu Gly Tyr Arg Leu Glu Asn Pro Val Val Asp Ser
 130 135 140

Leu Arg Leu Ala Arg Arg Gly Leu Pro Gly Leu Arg Arg Tyr Gly Leu
 145 150 155 160

Asp Ala Leu Ser Glu Val Leu Glu Leu Pro Arg Arg Thr Cys His Arg
 165 170 175

Ala Leu Glu Asp Val Glu Arg Thr Leu Ala Val Val His Glu Val Tyr
 180 185 190

Tyr Met Leu Thr Ser Gly
 195

<210> 89

<211> 182

<212> PRT

<213> *Deinococcus radiodurans*

<400> 89

Pro Trp Pro Gln Asp Val Val Val Phe Asp Leu Glu Thr Thr Gly Phe
 1 5 10 15

Ser Pro Ala Ser Ala Ala Ile Val Glu Ile Gly Ala Val Arg Ile Val

Asp Thr Ile Ile Glu Leu Ala Ala Val Lys Val Lys Gly Gly Glu Ile
50 55 60

Ile Asp Lys Phe Glu Ala Phe Ala Asn Pro His Arg Pro Leu Ser Ala
65 70 75 80

Thr Ile Ile Glu Leu Thr Gly Ile Thr Asp Asp Met Leu Gln Asp Ala
85 90 95

Pro Asp Val Val Asp Val Ile Arg Asp Phe Arg Glu Trp Ile Gly Asp
100 105 110

Asp Ile Leu Val Ala His Asn Ala Ser Phe Asp Met Gly Phe Leu Asn
115 120 125

Val Ala Tyr Lys Lys Leu Leu Glu Val Glu Lys Ala Lys Asn Pro Val
130 135 140

Ile Asp Thr Leu Glu Leu Gly Arg Phe Leu Tyr Pro Glu Phe Lys Asn
145 150 155 160

His Arg Leu Asn Thr Leu Cys Lys Lys Phe Asp Ile Glu Leu Thr Gln
165 170 175

His His Arg Ala Ile Tyr Asp Thr Glu Ala Thr Ala Tyr Leu Leu Leu
180 185 190

Lys Met Leu Lys Asp Ala Ala Glu Lys
195 200

<210> 91

<211> 188

<212> PRT

<213> Haemophilus influenzae

<400> 91

Met Ile Asn Pro Asn Arg Gln Ile Val Leu Asp Thr Glu Thr Thr Gly
1 5 10 15

Met Asn Gln Leu Gly Ala His Tyr Glu Gly His Cys Ile Ile Glu Ile
20 25 30

Gly Ala Val Glu Leu Ile Asn Arg Arg Tyr Thr Gly Asn Asn Xaa His
35 40 45

Ile Tyr Ile Lys Pro Asp Arg Pro Xaa Asp Pro Asp Ala Ile Lys Val
50 55 60

His Gly Ile Thr Asp Glu Met Leu Ala Asp Lys Pro Glu Phe Lys Glu
65 70 75 80

Val Ala Gln Asp Phe Leu Asp Tyr Ile Asn Gly Ala Glu Leu Leu Ile
85 90 95

His Asn Ala Pro Phe Asp Val Gly Phe Met Asp Tyr Glu Phe Arg Lys
100 105 110

Leu Asn Leu Asn Val Lys Thr Asp Asp Ile Cys Leu Val Thr Asp Thr
115 120 125

Leu Gln Met Ala Arg Gln Met Tyr Pro Gly Lys Arg Asn Asn Leu Asp
130 135 140

Ala Leu Cys Asp Arg Leu Gly Ile Asp Asn Ser Lys Arg Thr Leu His
145 150 155 160

Gly Ala Leu Leu Asp Ala Glu Ile Leu Ala Asp Val Tyr Leu Met Met
165 170 175

Thr Gly Gly Gln Thr Asn Leu Phe Asp Glu Glu Glu
180 185

<210> 92

<211> 189

<212> PRT

<213> Escherichia coli

<400> 92

Met Ser Thr Ala Ile Thr Arg Gln Ile Val Leu Asp Thr Glu Thr Thr
1 5 10 15

Gly Met Asn Gln Ile Gly Ala His Ser Glu Gly His Lys Ile Ile Glu
20 25 30

Ile Gly Ala Val Glu Val Val Asn Arg Arg Leu Thr Gly Asn Asn Phe
35 40 45

His Val Tyr Leu Lys Asp Arg Leu Val Asp Pro Glu Ala Phe Gly Val
50 55 60

His Gly Ile Ala Val Asp Phe Leu Leu Asp Lys Pro Thr Phe Ala Glu
65 70 75 80

Val Ala Val Glu Phe Met Asp Tyr Ile Arg Gly Ala Glu Leu Val Ile

85	90	95
His Asn Ala Ala Phe Asp Ile Gly Phe Met Asp Tyr Glu Phe Ser Leu		
100	105	110
Leu Lys Arg Asp Ile Ala Lys Thr Asn Thr Phe Cys Lys Val Thr Asp		
115	120	125
Ser Leu Ala Val Ala Arg Lys Met Phe Pro Gly Lys Arg Asn Ser Leu		
130	135	140
Asp Ala Leu Cys Ala Arg Tyr Glu Ile Asp Asn Ser Lys Arg Thr Leu		
145	150	155
His Gly Ala Leu Leu Asp Ala Gln Ile Leu Ala Glu Val Tyr Leu Ala		
165	170	175
Met Thr Gly Gly Gln Thr Ser Met Ala Phe Ala Met Glu		
180	185	
<210> 93		
<211> 201		
<212> PRT		
<213> Helicobacter pylori		
<400> 93		
Asn Leu Glu Tyr Leu Lys Ala Cys Gly Leu Asn Phe Ile Glu Thr Ser		
1	5	10
Glu Asn Leu Ile Thr Leu Lys Asn Leu Lys Thr Pro Leu Lys Asp Glu		
20	25	30
Val Phe Ser Phe Ile Asp Leu Glu Thr Thr Gly Ser Cys Pro Ile Lys		
35	40	45
His Glu Ile Leu Glu Ile Gly Ala Val Gln Val Lys Gly Gly Glu Ile		
50	55	60
Ile Asn Arg Phe Glu Thr Leu Val Lys Val Lys Ser Val Pro Asp Tyr		
65	70	75
Ile Ala Glu Leu Thr Gly Ile Thr Tyr Glu Asp Thr Leu Asn Ala Pro		
85	90	95
Ser Ala His Glu Ala Leu Gln Glu Leu Arg Leu Phe Leu Gly Asn Ser		
100	105	110

Val Phe Val Ala His Asn Ala Asn Phe Asp Tyr Asn Phe Leu Gly Arg
115 120 125

Tyr Phe Val Glu Lys Leu His Cys Pro Leu Leu Asn Leu Lys Leu Cys
130 135 140

Thr Leu Asp Leu Ser Lys Arg Ala Ile Leu Ser Met Arg Tyr Ser Leu
145 150 155 160

Ser Phe Leu Lys Glu Leu Leu Gly Phe Gly Ile Glu Val Ser His Arg
165 170 175

Ala Tyr Ala Asp Ala Leu Ala Ser Tyr Lys Leu Phe Glu Ile Cys Leu
180 185 190

Leu Asn Leu Pro Ser Tyr Ile Lys Thr
195 200

<210> 94
<211> 630
<212> DNA
<213> Thermus thermophilus

<400> 94
atggtggagc ggggtggcg gacccttctg gacgggaggt tcctcctgga ggagggggtg 60
gggctttggg agtggcgcta cccctttccc ctggaggggg aggcggtggt ggtcctggac 120
ctggagacca cggggcttgc cggcctggac gaggtgattg aggtgggcct cctccgcctg 180
gaggggggga ggcgcctccc cttccagagc ctcgtccggc ccctcccgcc cgccgaagcc 240
cgttcgtgga acctcaccgg catcccccg gaggccctgg aggaggcccc ctccctggag 300
gaggttctgg agaaggccta cccctccgc ggcgacgcca ccttggtgat ccacaacgcc 360
gcctttgacc tgggcttctt ccgcccggcc ttggagggcc tgggctaccg cctggaaaac 420
cccgtggtgg actccctgcg cttggccaga cggggcttac caggccttag gcgctacggc 480
ctggacgccc tctccgaggt cctggagctt cccgaagga cctgccaccg ggccctcgag 540
gacgtggagc gcaccctcgc cgtggtgcac gaggtatact atatgcttac gtccggccgt 600
ccccgcacgc tttgggaact cgggaggtag 630

<210> 95
<211> 210
<212> PRT
<213> Thermus thermophilus

<400> 95
Met Val Glu Arg Val Val Arg Thr Leu Leu Asp Gly Arg Phe Leu Leu
1 5 10 15

Glu Glu Gly Val Gly Leu Trp Glu Trp Arg Tyr Pro Phe Pro Leu Glu

00716964-12100

20 25 30

Gly Glu Ala Val Val Val Leu Asp Leu Glu Thr Thr Gly Leu Ala Gly
35 40 45

Leu Asp Glu Val Ile Glu Val Gly Leu Leu Arg Leu Glu Gly Gly Arg
50 55 60

Arg Leu Pro Phe Gln Ser Leu Val Arg Pro Leu Pro Pro Ala Glu Ala
65 70 75 80

Arg Ser Trp Asn Leu Thr Gly Ile Pro Arg Glu Ala Leu Glu Glu Ala
85 90 95

Pro Ser Leu Glu Glu Val Leu Glu Lys Ala Tyr Pro Leu Arg Gly Asp
100 105 110

Ala Thr Leu Val Ile His Asn Ala Ala Phe Asp Leu Gly Phe Leu Arg
115 120 125

Pro Ala Leu Glu Gly Leu Gly Tyr Arg Leu Glu Asn Pro Val Val Asp
130 135 140

Ser Leu Arg Leu Ala Arg Arg Gly Leu Pro Gly Leu Arg Arg Tyr Gly
145 150 155 160

Leu Asp Ala Leu Ser Glu Val Leu Glu Leu Pro Arg Arg Thr Cys His
165 170 175

Arg Ala Leu Glu Asp Val Glu Arg Thr Leu Ala Val Val His Glu Val
180 185 190

Tyr Tyr Met Leu Thr Ser Gly Arg Pro Arg Thr Leu Trp Glu Leu Gly
195 200 205

Arg Glx
210

<210> 96
<211> 461
<212> PRT
<213> *Pseudomonas marcesans*

<400> 96
Met Leu Glu Ala Ser Trp Glu Lys Val Gln Ser Ser Leu Lys Gln Asn
1 5 10 15

Leu Ser Lys Pro Ser Tyr Glu Thr Trp Ile Arg Pro Thr Glu Phe Ser
 20 25 30
 Gly Phe Lys Asn Gly Glu Leu Thr Leu Ile Ala Pro Asn Ser Phe Ser
 35 40 45
 Ser Ala Trp Leu Lys Asn Asn Tyr Ser Gln Thr Ile Gln Glu Thr Ala
 50 55 60
 Glu Glu Ile Phe Gly Glu Pro Val Thr Val His Val Lys Val Lys Ala
 65 70 75 80
 Asn Ala Glu Ser Ser Asp Glu His Tyr Ser Ser Ala Pro Ile Thr Pro
 85 90 95
 Pro Leu Glu Ala Ser Pro Gly Ser Val Asp Ser Ser Gly Ser Ser Leu
 100 105 110
 Arg Leu Ser Lys Lys Thr Leu Pro Leu Leu Asn Leu Arg Tyr Val Phe
 115 120 125
 Asn Arg Phe Val Val Gly Pro Asn Ser Arg Met Ala His Ala Ala Ala
 130 135 140
 Met Ala Val Ala Glu Ser Pro Gly Arg Glu Phe Asn Pro Leu Phe Ile
 145 150 155 160
 Cys Gly Gly Val Gly Leu Gly Lys Thr His Leu Met Gln Ala Ile Gly
 165 170 175
 His Tyr Arg Leu Glu Ile Asp Pro Gly Ala Lys Val Ser Tyr Val Ser
 180 185 190
 Thr Glu Thr Phe Thr Asn Asp Leu Ile Leu Ala Ile Arg Gln Asp Arg
 195 200 205
 Met Gln Ala Phe Arg Asp Arg Tyr Arg Ala Ala Asp Leu Ile Leu Val
 210 215 220
 Asp Asp Ile Gln Phe Ile Glu Gly Lys Glu Tyr Thr Gln Glu Glu Phe
 225 230 235 240
 Phe His Thr Phe Asn Ala Leu His Asp Ala Gly Ser Gln Ile Val Leu
 245 250 255
 Ala Ser Asp Arg Pro Pro Ser Gln Ile Pro Arg Leu Gln Glu Arg Leu
 260 265 270

Met Ser Arg Phe Ser Met Gly Leu Ile Ala Asp Val Gln Ala Pro Asp
275 280 285

Leu Glu Thr Arg Met Ala Ile Leu Gln Lys Lys Ala Glu His Glu Arg
290 295 300

Val Gly Leu Pro Arg Asp Leu Ile Gln Phe Ile Ala Gly Arg Phe Thr
305 310 315 320

Ser Asn Ile Arg Glu Leu Glu Gly Ala Leu Thr Arg Ala Ile Ala Phe
325 330 335

Ala Ser Ile Thr Gly Leu Pro Met Thr Val Asp Ser Ile Ala Pro Met
340 345 350

Leu Asp Pro Asn Gly Gln Gly Val Glu Val Thr Pro Lys Gln Val Leu
355 360 365

Asp Lys Val Ala Glu Val Phe Lys Val Thr Pro Asp Glu Met Arg Ser
370 375 380

Ala Ser Arg Arg Arg Pro Val Ser Gln Ala Arg Gln Val Gly Met Tyr
385 390 395 400

Leu Met Arg Gln Gly Thr Asn Leu Ser Leu Pro Arg Ile Gly Asp Thr
405 410 415

Phe Gly Gly Lys Asp His Thr Thr Val Met Tyr Ala Ile Glu Gln Val
420 425 430

Glu Lys Lys Leu Ser Ser Asp Pro Gln Ile Ala Ser Gln Val Gln Lys
435 440 445

Ile Arg Asp Leu Leu Gln Ile Asp Ser Arg Arg Lys Arg
450 455 460

<210> 97

<211> 447

<212> PRT

<213> Synechocystis sp.

<400> 97

Met Val Ser Cys Glu Asn Leu Trp Gln Gln Ala Leu Ala Ile Leu Ala
1 5 10 15

Thr Gln Leu Thr Lys Pro Ala Phe Asp Thr Trp Ile Lys Ala Ser Val
20 25 30

Leu Ile Ser Leu Gly Asp Gly Val Ala Thr Ile Gln Val Glu Asn Gly
 35 40 45
 Phe Val Leu Asn His Leu Gln Lys Ser Tyr Gly Pro Leu Leu Met Glu
 50 55 60
 Val Leu Thr Asp Leu Thr Gly Gln Glu Ile Thr Val Lys Leu Ile Thr
 65 70 75 80
 Asp Gly Leu Glu Pro His Ser Leu Ile Gly Gln Glu Ser Ser Leu Pro
 85 90 95
 Met Glu Thr Thr Pro Lys Asn Ala Thr Ala Leu Asn Gly Lys Tyr Thr
 100 105 110
 Phe Ser Arg Phe Val Val Gly Pro Thr Asn Arg Met Ala His Ala Ala
 115 120 125
 Ser Leu Ala Val Ala Glu Ser Pro Gly Arg Glu Phe Asn Pro Leu Phe
 130 135 140
 Leu Cys Gly Gly Val Gly Leu Gly Lys Thr His Leu Met Gln Ala Ile
 145 150 155 160
 Ala His Tyr Arg Leu Glu Met Tyr Pro Asn Ala Lys Val Tyr Tyr Val
 165 170 175
 Ser Thr Glu Arg Phe Thr Asn Asp Leu Ile Thr Ala Ile Arg Gln Asp
 180 185 190
 Asn Met Glu Asp Phe Arg Ser Tyr Tyr Arg Ser Ala Asp Phe Leu Leu
 195 200 205
 Ile Asp Asp Ile Gln Phe Ile Lys Gly Lys Glu Tyr Thr Gln Glu Glu
 210 215 220
 Phe Phe His Thr Phe Asn Ser Leu His Glu Ala Gly Lys Gln Val Val
 225 230 235 240
 Val Ala Ser Asp Arg Ala Pro Gln Arg Ile Pro Gly Leu Gln Asp Arg
 245 250 255
 Leu Ile Ser Arg Phe Ser Met Gly Leu Ile Ala Asp Ile Gln Val Pro
 260 265 270
 Asp Leu Glu Thr Arg Met Ala Ile Leu Gln Lys Lys Ala Glu Tyr Asp
 275 280 285

00116964.12100

50	55	60
Thr Ile Tyr Glu Leu Thr Gly Glu Glu Leu Ser Ile Lys Phe Val Ile		
65	70	75 80
Pro Gln Asn Gln Asp Val Glu Asp Phe Met Pro Lys Pro Gln Val Lys		
	85	90 95
Lys Ala Val Lys Glu Asp Thr Ser Asp Phe Pro Gln Asn Met Leu Asn		
	100	105 110
Pro Lys Tyr Thr Phe Asp Thr Phe Val Ile Gly Ser Gly Asn Arg Phe		
	115	120 125
Ala His Ala Ala Ser Leu Ala Val Ala Glu Ala Pro Ala Lys Ala Tyr		
	130	135 140
Asn Pro Leu Phe Ile Tyr Gly Gly Val Gly Leu Gly Lys Thr His Leu		
	145	150 155 160
Met His Ala Ile Gly His Tyr Val Ile Asp His Asn Pro Ser Ala Lys		
	165	170 175
Val Val Tyr Leu Ser Ser Glu Lys Phe Thr Asn Glu Phe Ile Asn Ser		
	180	185 190
Ile Arg Asp Asn Lys Ala Val Asp Phe Arg Asn Arg Tyr Arg Asn Val		
	195	200 205
Asp Val Leu Leu Ile Asp Asp Ile Gln Phe Leu Ala Gly Lys Glu Gln		
	210	215 220
Thr Gln Glu Glu Phe Phe His Thr Phe Asn Thr Leu His Glu Glu Ser		
	225	230 235 240
Lys Gln Ile Val Ile Ser Ser Asp Arg Pro Pro Lys Glu Ile Pro Thr		
	245	250 255
Leu Glu Asp Arg Leu Arg Ser Arg Phe Glu Trp Gly Leu Ile Thr Asp		
	260	265 270
Ile Thr Pro Pro Asp Leu Glu Thr Arg Ile Ala Ile Leu Arg Lys Lys		
	275	280 285
Ala Lys Ala Glu Gly Leu Asp Ile Pro Asn Glu Val Met Leu Tyr Ile		
	290	295 300
Ala Asn Gln Ile Asp Ser Asn Ile Arg Glu Leu Glu Gly Ala Leu Ile		

305 310 315 320

Arg Val Val Ala Tyr Ser Ser Leu Ile Asn Lys Asp Ile Asn Ala Asp
 325 330 335

Leu Ala Ala Glu Ala Leu Lys Asp Ile Ile Pro Ser Ser Lys Pro Lys
 340 345 350

Val Ile Thr Ile Lys Glu Ile Gln Arg Val Val Gly Gln Gln Phe Asn
 355 360 365

Ile Lys Leu Glu Asp Phe Lys Ala Lys Lys Arg Thr Lys Ser Val Ala
 370 375 380

Phe Pro Arg Gln Ile Ala Met Tyr Leu Ser Arg Glu Met Thr Asp Ser
385 390 395 400

Ser Leu Pro Lys Ile Gly Glu Glu Phe Gly Gly Arg Asp His Thr Thr
 405 410 415

Val Ile His Ala His Glu Lys Ile Ser Lys Leu Leu Ala Asp Asp Glu
 420 425 430

Gln Leu Gln Gln His Val Lys Glu Ile Lys Glu Gln Leu Lys
 435 440 445

<210> 99
<211> 507
<212> PRT
<213> Mycobacterium tuberculosis

<400> 99
Met Thr Asp Asp Pro Gly Ser Gly Phe Thr Thr Val Trp Asn Ala Val
1 5 10 15

Val Ser Glu Leu Asn Gly Asp Pro Lys Val Asp Asp Gly Pro Ser Ser
 20 25 30

Asp Ala Asn Leu Ser Ala Pro Leu Thr Pro Gln Gln Arg Ala Trp Leu
 35 40 45

Asn Leu Val Gln Pro Leu Thr Ile Val Glu Gly Phe Ala Leu Leu Ser
50 55 60

Val Pro Ser Ser Phe Val Gln Asn Glu Ile Glu Arg His Leu Arg Ala
65 70 75 80

Pro Ile Thr Asp Ala Leu Ser Arg Arg Leu Gly His Gln Ile Gln Leu
 85 90 95

Gly Val Arg Ile Ala Pro Pro Ala Thr Asp Glu Ala Asp Asp Thr Thr
 100 105 110

Val Pro Pro Ser Glu Asn Pro Ala Thr Thr Ser Pro Asp Thr Thr Thr
 115 120 125

Asp Asn Asp Glu Ile Asp Asp Ser Ala Ala Ala Arg Gly Asp Asn Gln
 130 135 140

His Ser Trp Pro Ser Tyr Phe Thr Glu Arg Pro His Asn Thr Asp Ser
 145 150 155 160

Ala Thr Ala Gly Val Thr Ser Leu Asn Arg Arg Tyr Thr Phe Asp Thr
 165 170 175

Phe Val Ile Gly Ala Ser Asn Arg Phe Ala His Ala Ala Ala Leu Ala
 180 185 190

Ile Ala Glu Ala Pro Ala Arg Ala Tyr Asn Pro Leu Phe Ile Trp Gly
 195 200 205

Glu Ser Gly Leu Gly Lys Thr His Leu Leu His Ala Ala Gly Asn Tyr
 210 215 220

Ala Gln Arg Leu Phe Pro Gly Met Arg Val Lys Tyr Val Ser Thr Glu
 225 230 235 240

Glu Phe Thr Asn Asp Phe Ile Asn Ser Leu Arg Asp Asp Arg Lys Val
 245 250 255

Ala Phe Lys Arg Ser Tyr Arg Asp Val Asp Val Leu Leu Val Asp Asp
 260 265 270

Ile Gln Phe Ile Glu Gly Lys Glu Gly Ile Gln Glu Glu Phe Phe His
 275 280 285

Thr Phe Asn Thr Leu His Asn Ala Asn Lys Gln Ile Val Ile Ser Ser
 290 295 300

Asp Arg Pro Pro Lys Gln Leu Ala Thr Leu Glu Asp Arg Leu Arg Thr
 305 310 315 320

Arg Phe Glu Trp Gly Leu Ile Thr Asp Val Gln Pro Pro Glu Leu Glu
 325 330 335

Ala Leu Asp Trp Ile Arg Arg His Tyr Ala Gly Leu Ile Gln Glu Gly
 50 55 60

Pro Arg Leu Leu Gly Ala Gln Ala Pro Arg Phe Glu Leu Arg Val Val
 65 70 75 80

Pro Gly Val Val Val Gln Glu Asp Ile Phe Gln Pro Pro Pro Ser Pro
 85 90 95

Pro Ala Gln Ala Gln Pro Glu Asp Thr Phe Lys Thr Ser Trp Trp Gly
 100 105 110

Pro Thr Thr Pro Trp Pro His Gly Gly Ala Val Ala Val Ala Glu Ser
 115 120 125

Pro Gly Arg Ala Tyr Asn Pro Leu Phe Ile Tyr Gly Gly Arg Gly Leu
 130 135 140

Gly Lys Thr Tyr Leu Met His Ala Val Gly Pro Leu Arg Ala Lys Arg
 145 150 155 160

Phe Pro His Met Arg Leu Glu Tyr Val Ser Thr Glu Thr Phe Thr Asn
 165 170 175

Glu Leu Ile Asn Arg Pro Ser Ala Arg Asp Arg Met Thr Glu Phe Arg
 180 185 190

Glu Arg Tyr Arg Ser Val Asp Leu Leu Leu Val Asp Asp Val Gln Phe
 195 200 205

Ile Ala Gly Lys Glu Arg Thr Gln Glu Glu Phe Phe His Thr Phe Asn
 210 215 220

Ala Leu Tyr Glu Ala His Lys Gln Ile Ile Leu Ser Ser Asp Arg Pro
 225 230 235 240

Pro Lys Asp Ile Leu Thr Leu Glu Ala Arg Leu Arg Ser Arg Phe Glu
 245 250 255

Trp Gly Leu Ile Thr Asp Asn Pro Ala Pro Asp Leu Glu Thr Arg Ile
 260 265 270

Ala Ile Leu Lys Met Asn Ala Ser Ser Gly Pro Glu Asp Pro Glu Asp
 275 280 285

Ala Leu Glu Tyr Ile Ala Arg Gln Val Thr Ser Asn Ile Arg Glu Trp
 290 295 300

Glu Gly Ala Leu Met Arg Ala Ser Pro Phe Ala Ser Leu Asn Gly Val
305 310 315 320

Glu Leu Thr Arg Ala Val Ala Ala Lys Ala Leu Arg His Leu Arg Pro
325 330 335

Arg Glu Leu Glu Ala Asp Pro Leu Glu Ile Ile Arg Lys Ala Ala Gly
340 345 350

Pro Val Arg Pro Glu Thr Pro Gly Gly Ala His Gly Glu Arg Arg Lys
355 360 365

Lys Glu Val Val Leu Pro Arg Gln Leu Ala Met Tyr Leu Val Arg Glu
370 375 380

Leu Thr Pro Ala Ser Leu Pro Glu Ile Gly Gln Leu Phe Gly Gly Arg
385 390 395 400

Asp His Thr Thr Val Arg Tyr Ala Ile Gln Lys Val Gln Glu Leu Ala
405 410 415

Gly Lys Pro Asp Arg Glu Val Gln Gly Leu Leu Arg Thr Leu Arg Glu
420 425 430

Ala Cys Thr Asp Pro Val Asp Asn Leu Trp Ile Thr Cys Gly
435 440 445

<210> 101

<211> 467

<212> PRT

<213> Escherichia coli

<400> 101

Met Ser Leu Ser Leu Trp Gln Gln Cys Leu Ala Arg Leu Gln Asp Glu
1 5 10 15

Leu Pro Ala Thr Glu Phe Ser Met Trp Ile Arg Pro Leu Gln Ala Glu
20 25 30

Leu Ser Asp Asn Thr Leu Ala Leu Tyr Ala Pro Asn Arg Phe Val Leu
35 40 45

Asp Trp Val Arg Asp Lys Tyr Leu Asn Asn Ile Asn Gly Leu Leu Thr
50 55 60

Ser Phe Cys Gly Ala Asp Ala Pro Gln Leu Arg Phe Glu Val Gly Thr

00716964-112100

65		70		75		80									
Lys	Pro	Val	Thr	Gln	Thr	Pro	Gln	Ala	Ala	Val	Thr	Ser	Asn	Val	Ala
				85					90					95	
Ala	Pro	Ala	Gln	Val	Ala	Gln	Thr	Gln	Pro	Gln	Arg	Ala	Ala	Pro	Ser
		100						105					110		
Thr	Arg	Ser	Gly	Trp	Asp	Asn	Val	Pro	Ala	Pro	Ala	Glu	Pro	Thr	Tyr
		115					120					125			
Arg	Ser	Asn	Val	Asn	Val	Lys	His	Thr	Phe	Asp	Asn	Phe	Val	Glu	Gly
	130					135					140				
Lys	Ser	Asn	Gln	Leu	Ala	Arg	Ala	Ala	Ala	Arg	Gln	Val	Ala	Asp	Asn
145				150						155					160
Pro	Gly	Gly	Ala	Tyr	Asn	Pro	Leu	Phe	Leu	Tyr	Gly	Gly	Thr	Gly	Leu
				165					170					175	
Gly	Lys	Thr	His	Leu	Leu	His	Ala	Val	Gly	Asn	Gly	Ile	Met	Ala	Arg
			180					185					190		
Lys	Pro	Asn	Ala	Lys	Val	Val	Tyr	Met	His	Ser	Glu	Arg	Phe	Val	Gln
		195					200					205			
Asp	Met	Val	Lys	Ala	Leu	Gln	Asn	Asn	Ala	Ile	Glu	Glu	Phe	Lys	Arg
	210					215					220				
Tyr	Tyr	Arg	Ser	Val	Asp	Ala	Leu	Leu	Ile	Asp	Asp	Ile	Gln	Phe	Phe
225				230						235				240	
Ala	Asn	Lys	Glu	Arg	Ser	Gln	Glu	Glu	Phe	Phe	His	Thr	Phe	Asn	Ala
				245					250					255	
Leu	Leu	Glu	Gly	Asn	Gln	Gln	Ile	Ile	Leu	Thr	Ser	Asp	Arg	Tyr	Pro
		260					265						270		
Lys	Glu	Ile	Asn	Gly	Val	Glu	Asp	Arg	Leu	Lys	Ser	Arg	Phe	Gly	Trp
		275					280					285			
Gly	Leu	Thr	Val	Ala	Ile	Glu	Pro	Pro	Glu	Leu	Glu	Thr	Arg	Val	Ala
	290					295					300				
Ile	Leu	Met	Lys	Lys	Ala	Asp	Glu	Asn	Asp	Ile	Arg	Leu	Pro	Gly	Glu
305					310					315				320	
Val	Ala	Phe	Phe	Ile	Ala	Lys	Arg	Leu	Arg	Ser	Asn	Val	Arg	Glu	Leu

	325		330		335
Glu Gly Ala Leu Asn Arg Val Ile Ala Asn Ala Asn Phe Thr Gly Arg					
	340		345		350
Ala Ile Thr Ile Asp Phe Val Arg Glu Ala Leu Arg Asp Leu Leu Ala					
	355		360		365
Leu Gln Glu Lys Leu Val Thr Ile Asp Asn Ile Gln Lys Thr Val Ala					
	370		375		380
Glu Tyr Tyr Lys Ile Lys Val Ala Asp Leu Leu Ser Lys Arg Arg Ser					
	385		390		395
Arg Ser Val Ala Arg Pro Arg Gln Met Ala Met Ala Leu Ala Lys Glu					
	405		410		415
Leu Thr Asn His Ser Leu Pro Glu Ile Gly Asp Ala Phe Gly Gly Arg					
	420		425		430
Asp His Thr Thr Val Leu His Ala Cys Arg Lys Ile Glu Gln Leu Arg					
	435		440		445
Glu Glu Ser His Asp Ile Lys Glu Asp Phe Ser Asn Leu Ile Arg Thr					
	450		455		460
Leu Ser Ser					
465					
<210> 102					
<211> 440					
<212> PRT					
<213> Thermatoga maritima					
<400> 102					
Met Lys Glu Arg Ile Leu Gln Glu Ile Lys Thr Arg Val Asn Arg Lys					
1 5 10 15					
Ser Trp Glu Leu Trp Phe Ser Ser Phe Asp Val Lys Ser Ile Glu Gly					
20 25 30					
Asn Lys Val Val Phe Ser Val Gly Asn Leu Phe Ile Lys Glu Trp Leu					
35 40 45					
Glu Lys Lys Tyr Tyr Ser Val Leu Ser Lys Ala Val Lys Val Val Leu					
50 55 60					

Gly Asn Asp Ala Thr Phe Glu Ile Thr Tyr Glu Ala Phe Glu Pro His
 65 70 75 80
 Ser Ser Tyr Ser Glu Pro Leu Val Lys Lys Arg Ala Val Leu Leu Thr
 85 90 95
 Pro Leu Asn Pro Asp Tyr Thr Phe Glu Asn Phe Val Val Gly Pro Gly
 100 105 110
 Asn Ser Phe Ala Tyr His Ala Ala Leu Glu Val Ala Lys His Pro Gly
 115 120 125
 Arg Tyr Asn Pro Leu Phe Ile Tyr Gly Gly Val Gly Leu Gly Lys Thr
 130 135 140
 His Leu Leu Gln Ser Ile Gly Asn Tyr Val Val Gln Asn Glu Pro Asp
 145 150 155 160
 Leu Arg Val Met Tyr Ile Thr Ser Glu Lys Phe Leu Asn Asp Leu Val
 165 170 175
 Asp Ser Met Lys Glu Gly Lys Leu Asn Glu Phe Arg Glu Lys Tyr Arg
 180 185 190
 Lys Lys Val Asp Ile Leu Leu Ile Asp Asp Val Gln Phe Leu Ile Gly
 195 200 205
 Lys Thr Gly Val Gln Thr Glu Leu Phe His Thr Phe Asn Glu Leu His
 210 215 220
 Asp Ser Gly Lys Gln Ile Val Ile Cys Ser Asp Arg Glu Pro Gln Lys
 225 230 235 240
 Leu Ser Glu Phe Gln Asp Arg Leu Val Ser Arg Phe Gln Met Gly Leu
 245 250 255
 Val Ala Lys Leu Glu Pro Pro Asp Glu Glu Thr Arg Lys Ser Ile Ala
 260 265 270
 Arg Lys Met Leu Glu Ile Glu His Gly Glu Leu Pro Glu Glu Val Leu
 275 280 285
 Asn Phe Val Ala Glu Asn Val Asp Asp Asn Leu Arg Arg Leu Arg Gly
 290 295 300
 Ala Ile Ile Lys Leu Leu Val Tyr Lys Glu Thr Thr Gly Lys Glu Val
 305 310 315 320

Asp Leu Lys Glu Ala Ile Leu Leu Leu Lys Asp Phe Ile Lys Pro Asn
 325 330 335

Arg Val Lys Ala Met Asp Pro Ile Asp Glu Leu Ile Glu Ile Val Ala
 340 345 350

Lys Val Thr Gly Val Pro Arg Glu Glu Ile Leu Ser Asn Ser Arg Asn
 355 360 365

Val Lys Ala Leu Thr Ala Arg Arg Ile Gly Met Tyr Val Ala Lys Asn
 370 375 380

Tyr Leu Lys Ser Ser Leu Arg Thr Ile Ala Glu Lys Phe Asn Arg Ser
 385 390 395 400

His Pro Val Val Val Asp Ser Val Lys Lys Val Lys Asp Ser Leu Leu
 405 410 415

Lys Gly Asn Lys Gln Leu Lys Ala Leu Ile Asp Glu Val Ile Gly Glu
 420 425 430

Ile Ser Arg Arg Ala Leu Ser Gly
 435 440

<210> 103

<211> 457

<212> PRT

<213> Helicobacter pylori

<400> 103

Met Asp Thr Asn Asn Asn Ile Glu Lys Glu Ile Leu Ala Leu Val Lys
 1 5 10 15

Gln Asn Pro Lys Val Ser Leu Ile Glu Tyr Glu Asn Tyr Phe Ser Gln
 20 25 30

Leu Lys Tyr Asn Pro Asn Ala Ser Lys Ser Asp Ile Ala Phe Phe Tyr
 35 40 45

Ala Pro Asn Gln Val Leu Cys Thr Thr Ile Thr Ala Lys Tyr Gly Ala
 50 55 60

Leu Leu Lys Glu Ile Leu Ser Gln Asn Lys Val Gly Met His Leu Ala
 65 70 75 80

His Ser Val Asp Val Arg Ile Glu Val Ala Pro Lys Ile Gln Ile Asn
 85 90 95

Ala Gln Ser Asn Ile Asn Tyr Lys Ala Ile Lys Thr Ser Val Lys Asp
 100 105 110
 Ser Tyr Thr Phe Glu Asn Phe Val Val Gly Ser Cys Asn Asn Thr Val
 115 120 125
 Tyr Glu Ile Ala Lys Lys Val Ala Gln Ser Asp Thr Pro Pro Tyr Asn
 130 135 140
 Pro Val Leu Phe Tyr Gly Gly Thr Gly Leu Gly Lys Thr His Ile Leu
 145 150 155 160
 Asn Ala Ile Gly Asn His Ala Leu Glu Lys His Lys Lys Val Val Leu
 165 170 175
 Val Thr Ser Glu Asp Phe Leu Thr Asp Phe Leu Lys His Leu Asp Asn
 180 185 190
 Lys Thr Met Asp Ser Phe Lys Ala Lys Tyr Arg His Cys Asp Phe Phe
 195 200 205
 Leu Leu Asp Asp Ala Gln Phe Leu Gln Gly Lys Pro Lys Leu Glu Glu
 210 215 220
 Glu Phe Phe His Thr Phe Asn Glu Leu His Ala Asn Ser Lys Gln Ile
 225 230 235 240
 Val Leu Ile Ser Asp Arg Ser Pro Lys Asn Ile Ala Gly Leu Glu Asp
 245 250 255
 Arg Leu Lys Ser Arg Phe Glu Trp Gly Ile Thr Ala Lys Val Met Pro
 260 265 270
 Pro Asp Leu Glu Thr Lys Leu Ser Ile Val Lys Gln Lys Cys Gln Leu
 275 280 285
 Asn Gln Ile Thr Leu Pro Glu Glu Val Met Glu Tyr Ile Ala Gln His
 290 295 300
 Ile Ser Asp Asn Ile Arg Gln Met Glu Gly Ala Ile Ile Lys Ile Ser
 305 310 315 320
 Val Asn Ala Asn Leu Met Asn Ala Ser Ile Asp Leu Asn Leu Ala Lys
 325 330 335
 Thr Val Leu Glu Asp Leu Gln Lys Asp His Ala Glu Gly Ser Ser Leu
 340 345 350

Glu Asn Ile Leu Leu Ala Val Ala Gln Ser Leu Asn Leu Lys Ser Ser
355 360 365

Glu Ile Lys Val Ser Ser Arg Gln Lys Asn Val Ala Leu Ala Arg Lys
370 375 380

Leu Val Val Tyr Phe Ala Arg Leu Tyr Thr Pro Asn Pro Thr Leu Ser
385 390 395 400

Leu Ala Gln Phe Leu Asp Leu Lys Asp His Ser Ser Ile Ser Lys Met
405 410 415

Tyr Ser Gly Val Lys Lys Met Leu Glu Glu Glu Lys Ser Pro Phe Val
420 425 430

Leu Ser Leu Arg Glu Glu Ile Lys Asn Arg Leu Asn Glu Leu Asn Asp
435 440 445

Lys Lys Thr Ala Phe Asn Ser Ser Glu
450 455

<210> 104

<211> 1305

<212> DNA

<213> Thermus thermophilus

<400> 104

gtgtgcacag aggccgtctg gcaacacggt ctggagcaca tccgccgcag catcaccgag 60
gtggagttcc acacctggtt tgaaaggatc cgccccttgg ggatccggga cggggtgctg 120
gagctcgccg tgcacacctc ctttgccctg gactggatcc ggcgccacta cgccggcctc 180
atccaggagg gccctcggtt cctcggggcc caggcgcccc ggtttgagct ccgggtggtg 240
cccggggtcg tagtcagga ggacatcttc cagccccgcg cgagcccccc ggcccaagct 300
caaccggaag atacctttaa aacttcgttg tggggcccaa caactccatg gccccacggc 360
ggcgccgtgg ccgtggccga gtcccccggc cgggcctaca acccctctt catctacggg 420
ggccgtggcc tgggaaagac ctacctgatg cagccgtgg gccactccg tgcgaagcgc 480
ttcccccaca tgagattaga gtacgtttcc acggaactt tcaccaacga gctcatcaac 540
cggccatccg cgagggaccg gatgacggag ttccgggagc ggtaccgctc cgtggacctc 600
ctgctggtgg acgacgtcca gttcatcgcc ggaaaggagc gcaccagga ggagtttttc 660
cacaccttca acgcccttta cgaggccac aagcagatca tcctctctc cgaccggccg 720
cccaaggaca tcctcaccct ggaggcgcg ctcgggagcc gctttgagtg gggcctgatc 780
accgacaatc cagccccga cctggaaacc cggatcgcca tcctgaagat gaacgccagc 840
agcgggcctg aggatccga ggacgccctg gagtacatcg cccggcaggt cacctccaac 900
atccgggagt gggaagggc cctcatgcgg gcacgcctt tcgcctccct caacggcggt 960
gagctgacct gcgccgtggc ggccaaggct ctccgacatc ttgcggccag ggagctggag 1020
gcggaacctt tggagatcat ccgcaaagcg gcgggaccag ttgggcctga aaccccgga 1080
ggagctcacg gggagcgccg caagaaggag gtggtcctcc cccggcagct cgccatgtac 1140

ctggtgcggg agctacccc ggctccctg cccgagatcg accagctcaa cgacgaccgg 1200
gaccacacca cggtcctcta cgccatccag aaggtccagg agctcgcgga aagcgaccgg 1260
gaggtgcagg gcctcctccg caccctccgg gaggcgtgca catga 1305

<210> 105

<211> 434

<212> PRT

<213> Thermus thermophilus

<400> 105

Val Ser His Glu Ala Val Trp Gln His Val Leu Glu His Ile Arg Arg
1 5 10 15

Ser Ile Thr Glu Val Glu Phe His Thr Trp Phe Glu Arg Ile Arg Pro
20 25 30

Leu Gly Ile Arg Asp Gly Val Leu Glu Leu Ala Val Pro Thr Ser Phe
35 40 45

Ala Leu Asp Trp Ile Arg Arg His Tyr Ala Gly Leu Ile Gln Glu Gly
50 55 60

Pro Arg Leu Leu Gly Ala Gln Ala Pro Arg Phe Glu Leu Arg Val Val
65 70 75 80

Pro Gly Val Val Val Gln Glu Asp Ile Phe Gln Pro Pro Pro Ser Pro
85 90 95

Pro Ala Gln Ala Gln Pro Glu Asp Thr Phe Lys Thr Ser Trp Trp Gly
100 105 110

Pro Thr Thr Pro Trp Pro His Gly Gly Ala Val Ala Val Ala Glu Ser
115 120 125

Pro Gly Arg Ala Tyr Asn Pro Leu Phe Ile Tyr Gly Gly Arg Gly Leu
130 135 140

Gly Lys Thr Tyr Leu Met His Ala Val Gly Pro Leu Arg Ala Lys Arg
145 150 155 160

Phe Pro His Met Arg Leu Glu Tyr Val Ser Thr Glu Thr Phe Thr Asn
165 170 175

Glu Leu Ile Asn Arg Pro Ser Ala Arg Asp Arg Met Thr Glu Phe Arg
180 185 190

Glu Arg Tyr Arg Ser Val Asp Leu Leu Leu Val Asp Asp Val Gln Phe

0016964.12100

195	200	205
Ile Ala Gly Lys Glu Arg Thr Gln Glu Glu Phe Phe His Thr Phe Asn		
210	215	220
Ala Leu Tyr Glu Ala His Lys Gln Ile Ile Leu Ser Ser Asp Arg Pro		
225	230	235 240
Pro Lys Asp Ile Leu Thr Leu Glu Ala Arg Leu Arg Ser Arg Phe Glu		
	245	250 255
Trp Gly Leu Ile Thr Asp Asn Pro Ala Pro Asp Leu Glu Thr Arg Ile		
	260	265 270
Ala Ile Leu Lys Met Asn Ala Ser Ser Gly Pro Glu Asp Pro Glu Asp		
	275	280 285
Ala Leu Glu Tyr Ile Ala Arg Gln Val Thr Ser Asn Ile Arg Glu Trp		
	290	295 300
Glu Gly Ala Leu Met Arg Ala Ser Pro Phe Ala Ser Leu Asn Gly Val		
305	310	315 320
Glu Leu Thr Arg Ala Val Ala Ala Lys Ala Leu Arg His Leu Arg Pro		
	325	330 335
Arg Glu Leu Glu Ala Asp Pro Leu Glu Ile Ile Arg Lys Ala Ala Gly		
	340	345 350
Pro Val Arg Pro Glu Thr Pro Gly Gly Ala His Gly Glu Arg Arg Lys		
	355	360 365
Lys Glu Val Val Leu Pro Arg Gln Leu Ala Met Tyr Leu Val Arg Glu		
	370	375 380
Leu Thr Pro Ala Ser Leu Pro Glu Ile Asp Gln Leu Asn Asp Asp Arg		
385	390	395 400
Asp His Thr Thr Val Leu Tyr Ala Ile Gln Lys Val Gln Glu Leu Ala		
	405	410 415
Glu Ser Asp Arg Glu Val Gln Gly Leu Leu Arg Thr Leu Arg Glu Ala		
	420	425 430
Cys Thr		

<210> 106
 <211> 1128
 <212> DNA
 <213> *Thermus thermophilus*

<400> 106
 atgaacataa cggttcccaa aaaactcctc tcggaccagc tttccctcct ggagcgcac 60
 gtccccctcta gaagcgccaa cccccctctac acctacctgg ggctttacgc cgaggaaggg 120
 gccttgatcc tcttcgggac caacggggag gtggacctcg aggtccgcct ccccgccgag 180
 gccc aaagcc ttccccgggt gctcgtcccc gccagccct tcttcagct ggtgcggagc 240
 cttcctgggg acctcgtggc cctcggcctc gcctcggagc cggg ccaggg ggggcagctg 300
 gagctctcct ccgggcgttt ccgcaccccg ctcagcctgg cccctgccga gggctacccc 360
 gagcttctgg tgcccagagg ggaggacaag ggggccttcc ccctccggac gcggatgccc 420
 tccggggagc tcgtcaaggc cttgacccac gtgcgctacg ccgcgagcaa cgaggagtac 480
 cgggccatct tccgcggggg gcagctggag ttctcccccc agggcttccg ggcggtggcc 540
 tccgacgggt accgcctcgc cctctacgac ctgcccctgc cccaagggtt ccaggccaag 600
 gccgtggtcc ccgcccggag cgtggacgag atggtgcggg tcctgaaggg ggcggacggg 660
 gccgaggccg tcctcgccct gggcgagggg gtgttgggcc tggccctcga gggcggaagc 720
 ggggtccgga tggccctccg cctcatggaa ggggagttcc ccgactacca gagggtcatc 780
 ccccaggagt tcgccctcaa ggtccaggtg gagggggagg ccctcaggga ggcggtgcgc 840
 cgggtgagcg tcctctccga ccggcagaac caccgggtgg acctcctttt ggaggaaggc 900
 cggatcctcc tctccgccga gggggactac ggcaaggggc aggaggaggt gcccgccag 960
 gtggaggggc cggacatggc cgtggcctac aacgcccgt acctcctcga ggccctcgcc 1020
 cccgtggggg accgggccc cctgggcac tccgggccc cgagcccag cctcatctgg 1080
 ggggacgggg aggggtaccg ggcggtggtg gtgcccctca ggggtctag 1128

<210> 107
 <211> 376
 <212> PRT
 <213> *Thermus thermophilus*

<400> 107
 Met Asn Ile Thr Val Pro Lys Lys Leu Leu Ser Asp Gln Leu Ser Leu
 1 5 10 15
 Leu Glu Arg Ile Val Pro Ser Arg Ser Ala Asn Pro Leu Tyr Thr Tyr
 20 25 30
 Leu Gly Leu Tyr Ala Glu Glu Gly Ala Leu Ile Leu Phe Gly Thr Asn
 35 40 45
 Gly Glu Val Asp Leu Glu Val Arg Leu Pro Ala Glu Ala Gln Ser Leu
 50 55 60
 Pro Arg Val Leu Val Pro Ala Gln Pro Phe Phe Gln Leu Val Arg Ser
 65 70 75 80

Leu Pro Gly Asp Leu Val Ala Leu Gly Leu Ala Ser Glu Pro Gly Gln
 85 90 95

Gly Gly Gln Leu Glu Leu Ser Ser Gly Arg Phe Arg Thr Arg Leu Ser
 100 105 110

Leu Ala Pro Ala Glu Gly Tyr Pro Glu Leu Leu Val Pro Glu Gly Glu
 115 120 125

Asp Lys Gly Ala Phe Pro Leu Arg Thr Arg Met Pro Ser Gly Glu Leu
 130 135 140

Val Lys Ala Leu Thr His Val Arg Tyr Ala Ala Ser Asn Glu Glu Tyr
 145 150 155 160

Arg Ala Ile Phe Arg Gly Val Gln Leu Glu Phe Ser Pro Gln Gly Phe
 165 170 175

Arg Ala Val Ala Ser Asp Gly Tyr Arg Leu Ala Leu Tyr Asp Leu Pro
 180 185 190

Leu Pro Gln Gly Phe Gln Ala Lys Ala Val Val Pro Ala Arg Ser Val
 195 200 205

Asp Glu Met Val Arg Val Leu Lys Gly Ala Asp Gly Ala Glu Ala Val
 210 215 220

Leu Ala Leu Gly Glu Gly Val Leu Ala Leu Ala Leu Glu Gly Gly Ser
 225 230 235 240

Gly Val Arg Met Ala Leu Arg Leu Met Glu Gly Glu Phe Pro Asp Tyr
 245 250 255

Gln Arg Val Ile Pro Gln Glu Phe Ala Leu Lys Val Gln Val Glu Gly
 260 265 270

Glu Ala Leu Arg Glu Ala Val Arg Arg Val Ser Val Leu Ser Asp Arg
 275 280 285

Gln Asn His Arg Val Asp Leu Leu Leu Glu Glu Gly Arg Ile Leu Leu
 290 295 300

Ser Ala Glu Gly Asp Tyr Gly Lys Gly Gln Glu Glu Val Pro Ala Gln
 305 310 315 320

Val Glu Gly Pro Asp Met Ala Val Ala Tyr Asn Ala Arg Tyr Leu Leu
 325 330 335

00746964.43400

Glu Ala Leu Ala Pro Val Gly Asp Arg Ala His Leu Gly Ile Ser Gly
 340 345 350

Pro Thr Ser Pro Ser Leu Ile Trp Gly Asp Gly Glu Gly Tyr Arg Ala
 355 360 365

Val Val Val Pro Leu Arg Val Glx
 370 375

<210> 108

<211> 376

<212> PRT

<213> Thermus thermophilus

<400> 108

Met Asn Ile Thr Val Pro Lys Lys Leu Leu Ser Asp Gln Leu Ser Leu
 1 5 10 15

Leu Glu Arg Ile Val Pro Ser Arg Ser Ala Asn Pro Leu Tyr Thr Tyr
 20 25 30

Leu Gly Leu Tyr Ala Glu Glu Gly Ala Leu Ile Leu Phe Gly Thr Asn
 35 40 45

Gly Glu Val Asp Leu Glu Val Arg Leu Pro Ala Glu Ala Gln Ser Leu
 50 55 60

Pro Arg Val Leu Val Pro Ala Gln Pro Phe Phe Gln Leu Val Arg Ser
 65 70 75 80

Leu Pro Gly Asp Leu Val Ala Leu Gly Leu Ala Ser Glu Pro Gly Gln
 85 90 95

Gly Gly Gln Leu Glu Leu Ser Ser Gly Arg Phe Arg Thr Arg Leu Ser
 100 105 110

Leu Ala Pro Ala Glu Gly Tyr Pro Glu Leu Leu Val Pro Glu Gly Glu
 115 120 125

Asp Lys Gly Ala Phe Pro Leu Arg Thr Arg Met Pro Ser Gly Glu Leu
 130 135 140

Val Lys Ala Leu Thr His Val Arg Tyr Ala Ala Ser Asn Glu Glu Tyr
 145 150 155 160

Arg Ala Ile Phe Arg Gly Val Gln Leu Glu Phe Ser Pro Gln Gly Phe
 165 170 175

Arg Ala Val Ala Ser Asp Gly Tyr Arg Leu Ala Leu Tyr Asp Leu Pro
 180 185 190

Leu Pro Gln Gly Phe Gln Ala Lys Ala Val Val Pro Ala Arg Ser Val
 195 200 205

Asp Glu Met Val Arg Val Leu Lys Gly Ala Asp Gly Ala Glu Ala Val
 210 215 220

Leu Ala Leu Gly Glu Gly Val Leu Ala Leu Ala Leu Glu Gly Gly Ser
 225 230 235 240

Gly Val Arg Met Ala Leu Arg Leu Met Glu Gly Glu Phe Pro Asp Tyr
 245 250 255

Gln Arg Val Ile Pro Gln Glu Phe Ala Leu Lys Val Gln Val Glu Gly
 260 265 270

Glu Ala Leu Arg Glu Ala Val Arg Arg Val Ser Val Leu Ser Asp Arg
 275 280 285

Gln Asn His Arg Val Asp Leu Leu Leu Glu Glu Gly Arg Ile Leu Leu
 290 295 300

Ser Ala Glu Gly Asp Tyr Gly Lys Gly Gln Glu Glu Val Pro Ala Gln
 305 310 315 320

Val Glu Gly Pro Asp Met Ala Val Ala Tyr Asn Ala Arg Tyr Leu Leu
 325 330 335

Glu Ala Leu Ala Pro Val Gly Asp Arg Ala His Leu Gly Ile Ser Gly
 340 345 350

Pro Thr Ser Pro Ser Leu Ile Trp Gly Asp Gly Glu Gly Tyr Arg Ala
 355 360 365

Val Val Val Pro Leu Arg Val Glx
 370 375

<210> 109

<211> 367

<212> PRT

<213> Escherichia coli

<400> 109

Met Lys Phe Thr Val Glu Arg Glu His Leu Leu Lys Pro Leu Gln Gln

00746964.12100

1	5	10	15
Val Ser Gly Pro Leu Gly Gly Arg Pro Thr Leu Pro Ile Leu Gly Asn			
20	25	30	
Leu Leu Leu Gln Val Ala Asp Gly Thr Leu Ser Leu Thr Gly Thr Asp			
35	40	45	
Leu Glu Met Glu Met Val Ala Arg Val Ala Leu Val Gln Pro His Glu			
50	55	60	
Pro Gly Ala Thr Thr Val Pro Ala Arg Lys Phe Phe Asp Ile Cys Arg			
65	70	75	80
Gly Leu Pro Glu Gly Ala Glu Ile Ala Val Gln Leu Glu Gly Glu Arg			
85	90	95	
Met Leu Val Arg Ser Gly Arg Ser Arg Phe Ser Leu Ser Thr Leu Pro			
100	105	110	
Ala Ala Asp Phe Pro Asn Leu Asp Asp Trp Gln Ser Glu Val Glu Phe			
115	120	125	
Thr Leu Pro Gln Ala Thr Met Lys Arg Leu Ile Glu Ala Thr Gln Phe			
130	135	140	
Ser Met Ala His Gln Asp Val Arg Tyr Tyr Leu Asn Gly Met Leu Phe			
145	150	155	160
Glu Thr Glu Gly Glu Glu Leu Arg Thr Val Ala Thr Asp Gly His Arg			
165	170	175	
Leu Ala Val Cys Ser Met Pro Ile Gly Gln Ser Leu Pro Ser His Ser			
180	185	190	
Val Ile Val Pro Arg Lys Gly Val Ile Glu Leu Met Arg Met Leu Asp			
195	200	205	
Gly Gly Asp Asn Pro Leu Arg Val Gln Ile Gly Ser Asn Asn Ile Arg			
210	215	220	
Ala His Val Gly Asp Phe Ile Phe Thr Ser Lys Leu Val Asp Gly Arg			
225	230	235	240
Phe Pro Asp Tyr Arg Arg Val Leu Pro Lys Asn Pro Asp Lys His Leu			
245	250	255	
Glu Ala Gly Cys Asp Leu Leu Lys Gln Ala Phe Ala Arg Ala Ala Ile			

001211 49697260

260	265	270
Leu Ser Asn Glu Lys Phe Arg Gly Val Arg Leu Tyr Val Ser Glu Asn		
275	280	285
Gln Leu Lys Ile Thr Ala Asn Asn Pro Glu Gln Glu Glu Ala Glu Glu		
290	295	300
Ile Leu Asp Val Thr Tyr Ser Gly Ala Glu Met Glu Ile Gly Phe Asn		
305	310	315
Val Ser Tyr Val Leu Asp Val Leu Asn Ala Leu Lys Cys Glu Asn Val		
325	330	335
Arg Met Met Leu Thr Asp Ser Val Ser Ser Val Gln Ile Glu Asp Ala		
340	345	350
Ala Ser Gln Ser Ala Ala Tyr Val Val Met Pro Met Arg Leu Glx		
355	360	365
<210> 110		
<211> 367		
<212> PRT		
<213> Proteus mirabilis		
<400> 110		
Met Lys Phe Ile Ile Glu Arg Glu Gln Leu Leu Lys Pro Leu Gln Gln		
1	5	10
Val Ser Gly Pro Leu Gly Gly Arg Pro Thr Leu Pro Ile Leu Gly Asn		
20	25	30
Leu Leu Leu Lys Val Thr Glu Asn Thr Leu Ser Leu Thr Gly Thr Asp		
35	40	45
Leu Glu Met Glu Met Met Ala Arg Val Ser Leu Ser Gln Ser His Glu		
50	55	60
Ile Gly Ala Thr Thr Val Pro Ala Arg Lys Phe Phe Asp Ile Trp Arg		
65	70	75
Gly Leu Pro Glu Gly Ala Glu Ile Ser Val Glu Leu Asp Gly Asp Arg		
85	90	95
Leu Leu Val Arg Ser Gly Arg Ser Arg Phe Ser Leu Ser Thr Leu Pro		
100	105	110

Ala Ser Asp Phe Pro Asn Leu Asp Asp Trp Gln Ser Glu Val Glu Phe
 115 120 125

Thr Leu Pro Gln Ala Thr Leu Lys Arg Leu Ile Glu Ser Thr Gln Phe
 130 135 140

Ser Met Ala His Gln Asp Val Arg Tyr Tyr Leu Asn Gly Met Leu Phe
 145 150 155 160

Glu Thr Glu Asn Thr Glu Leu Arg Thr Val Ala Thr Asp Gly His Arg
 165 170 175

Leu Ala Val Cys Ala Met Asp Ile Gly Gln Ser Leu Pro Gly His Ser
 180 185 190

Val Ile Val Pro Arg Lys Gly Val Ile Glu Leu Met Arg Leu Leu Asp
 195 200 205

Gly Ser Gly Glu Ser Leu Leu Gln Leu Gln Ile Gly Ser Asn Asn Leu
 210 215 220

Arg Ala His Val Gly Asp Phe Ile Phe Thr Ser Lys Leu Val Asp Gly
 225 230 235 240

Arg Phe Pro Asp Tyr Arg Arg Val Leu Pro Lys Asn Pro Thr Lys Thr
 245 250 255

Val Ile Ala Gly Cys Asp Ile Leu Lys Gln Ala Phe Ser Arg Ala Ala
 260 265 270

Ile Leu Ser Asn Glu Lys Phe Arg Gly Val Arg Ile Asn Leu Thr Asn
 275 280 285

Gly Gln Leu Lys Ile Thr Ala Asn Asn Pro Glu Gln Glu Glu Ala Glu
 290 295 300

Glu Ile Val Asp Val Gln Tyr Gln Gly Glu Glu Met Glu Ile Gly Phe
 305 310 315 320

Asn Val Ser Tyr Leu Leu Asp Val Leu Asn Thr Leu Lys Cys Glu Glu
 325 330 335

Val Lys Leu Leu Leu Thr Asp Ala Val Ser Ser Val Gln Val Glu Asn
 340 345 350

Val Ala Ser Ala Ala Ala Tyr Val Val Met Pro Met Arg Leu
 355 360 365

<210> 111
 <211> 366
 <212> PRT
 <213> Haemophilus influenzae

<400> 111
 Met Gln Phe Ser Ile Ser Arg Glu Asn Leu Leu Lys Pro Leu Gln Gln
 1 5 10 15
 Val Cys Gly Val Leu Ser Asn Arg Pro Asn Ile Pro Val Leu Asn Asn
 20 25 30
 Val Leu Leu Gln Ile Glu Asp Tyr Arg Leu Thr Ile Thr Gly Thr Asp
 35 40 45
 Leu Glu Val Glu Leu Ser Ser Gln Thr Gln Leu Ser Ser Ser Ser Glu
 50 55 60
 Asn Gly Thr Phe Thr Ile Pro Ala Lys Lys Phe Leu Asp Ile Cys Arg
 65 70 75 80
 Thr Leu Ser Asp Asp Ser Glu Ile Thr Val Thr Phe Glu Gln Asp Arg
 85 90 95
 Ala Leu Val Gln Ser Gly Arg Ser Arg Phe Thr Leu Ala Thr Gln Pro
 100 105 110
 Ala Glu Glu Tyr Pro Asn Leu Thr Asp Trp Gln Ser Glu Val Asp Phe
 115 120 125
 Glu Leu Pro Gln Asn Thr Leu Arg Arg Leu Ile Glu Ala Thr Gln Phe
 130 135 140
 Ser Met Ala Asn Gln Asp Ala Arg Tyr Phe Leu Asn Gly Met Lys Phe
 145 150 155 160
 Glu Thr Glu Gly Asn Leu Leu Arg Thr Val Ala Thr Asp Gly His Arg
 165 170 175
 Leu Ala Val Cys Thr Ile Ser Leu Glu Gln Glu Leu Gln Asn His Ser
 180 185 190
 Val Ile Leu Pro Arg Lys Gly Val Leu Glu Leu Val Arg Leu Leu Glu
 195 200 205
 Thr Asn Asp Glu Pro Ala Arg Leu Gln Ile Gly Thr Asn Asn Leu Arg
 210 215 220

00716964.112100

Val His Leu Lys Asn Thr Val Phe Thr Ser Lys Leu Ile Asp Gly Arg
225 230 235 240

Phe Pro Asp Tyr Arg Arg Val Leu Pro Arg Asn Ala Thr Lys Ile Val
245 250 255

Glu Gly Asn Trp Glu Met Leu Lys Gln Ala Phe Ala Arg Ala Ser Ile
260 265 270

Leu Ser Asn Glu Arg Ala Arg Ser Val Arg Leu Ser Leu Lys Glu Asn
275 280 285

Gln Leu Lys Ile Thr Ala Ser Asn Thr Glu His Glu Glu Ala Glu Glu
290 295 300

Ile Val Asp Val Asn Tyr Asn Gly Glu Glu Leu Glu Val Gly Phe Asn
305 310 315 320

Val Thr Tyr Ile Leu Asp Val Leu Asn Ala Leu Lys Cys Asn Gln Val
325 330 335

Arg Met Cys Leu Thr Asp Ala Phe Ser Ser Cys Leu Ile Glu Asn Cys
340 345 350

Glu Asp Ser Ser Cys Glu Tyr Val Ile Met Pro Met Arg Leu
355 360 365

<210> 112

<211> 367

<212> PRT

<213> Pseudomonas putida

<400> 112

Met His Phe Thr Ile Gln Arg Glu Ala Leu Leu Lys Pro Leu Gln Leu
1 5 10 15

Val Ala Gly Val Val Glu Arg Arg Gln Thr Leu Pro Val Leu Ser Asn
20 25 30

Val Leu Leu Val Val Gln Gly Gln Gln Leu Ser Leu Thr Gly Thr Asp
35 40 45

Leu Glu Val Glu Leu Val Gly Arg Val Gln Leu Glu Glu Pro Ala Glu
50 55 60

Pro Gly Glu Ile Thr Val Pro Ala Arg Lys Leu Met Asp Ile Cys Lys

00716964.12100

65		70		75		80									
Ser	Leu	Pro	Asn	Asp	Ala	Leu	Ile	Asp	Ile	Lys	Val	Asp	Glu	Gln	Lys
			85						90					95	
Leu	Leu	Val	Lys	Ala	Gly	Arg	Ser	Arg	Phe	Thr	Leu	Ser	Thr	Leu	Pro
		100						105					110		
Ala	Asn	Asp	Phe	Pro	Thr	Val	Glu	Gly	Pro	Gly	Ser	Leu	Thr	Cys	
		115						120				125			
Asn	Leu	Glu	Gln	Ser	Lys	Leu	Arg	Arg	Leu	Ile	Glu	Arg	Thr	Ser	Phe
	130						135					140			
Ala	Met	Ala	Gln	Gln	Asp	Val	Arg	Tyr	Tyr	Leu	Asn	Gly	Met	Leu	Leu
145						150				155					160
Glu	Val	Ser	Arg	Asn	Thr	Leu	Arg	Ala	Val	Ser	Thr	Asp	Gly	His	Arg
				165					170					175	
Leu	Ala	Leu	Cys	Ser	Met	Ser	Ala	Pro	Ile	Glu	Gln	Glu	Asp	Arg	His
			180					185						190	
Gln	Val	Ile	Val	Pro	Arg	Lys	Gly	Ile	Leu	Glu	Leu	Ala	Arg	Leu	Leu
		195						200					205		
Thr	Asp	Pro	Glu	Gly	Met	Val	Ser	Ile	Val	Leu	Gly	Gln	His	His	Ile
	210						215					220			
Arg	Ala	Thr	Thr	Gly	Glu	Phe	Thr	Phe	Thr	Ser	Lys	Leu	Val	Asp	Gly
225					230					235					240
Lys	Phe	Pro	Asp	Tyr	Glu	Arg	Val	Leu	Pro	Lys	Gly	Gly	Asp	Lys	Leu
				245					250					255	
Val	Val	Gly	Asp	Arg	Gln	Ala	Leu	Arg	Glu	Ala	Phe	Ser	Arg	Thr	Ala
			260					265					270		
Ile	Leu	Ser	Asn	Glu	Lys	Tyr	Arg	Gly	Ile	Arg	Leu	Gln	Leu	Ala	Ala
		275					280					285			
Gly	Gln	Leu	Lys	Ile	Gln	Ala	Asn	Asn	Pro	Glu	Gln	Glu	Glu	Ala	Glu
	290					295					300				
Glu	Glu	Ile	Ser	Val	Asp	Tyr	Glu	Gly	Ser	Ser	Leu	Glu	Ile	Gly	Phe
305					310					315					320
Asn	Val	Ser	Tyr	Leu	Leu	Asp	Val	Leu	Gly	Val	Met	Thr	Thr	Glu	Gln

325	330	335
Val Arg Leu Ile Leu Ser Asp Ser Asn Ser Ser Ala Leu Leu Gln Glu		
340	345	350
Ala Gly Asn Asp Asp Ser Ser Tyr Val Val Met Pro Met Arg Leu		
355	360	365
<210> 113		
<211> 366		
<212> PRT		
<213> Buchnera aphidicola		
<400> 113		
Met Lys Phe Thr Ile Gln Asn Asp Ile Leu Thr Lys Asn Leu Lys Lys		
1	5	10 15
Ile Thr Arg Val Leu Val Lys Asn Ile Ser Phe Pro Ile Leu Glu Asn		
20	25	30
Ile Leu Ile Gln Val Glu Asp Gly Thr Leu Ser Leu Thr Thr Thr Asn		
35	40	45
Leu Glu Ile Glu Leu Ile Ser Lys Ile Glu Ile Ile Thr Lys Tyr Ile		
50	55	60
Pro Gly Lys Thr Thr Ile Ser Gly Arg Lys Ile Leu Asn Ile Cys Arg		
65	70	75 80
Thr Leu Ser Glu Lys Ser Lys Ile Lys Met Gln Leu Lys Asn Lys Lys		
85	90	95
Met Tyr Ile Ser Ser Glu Asn Ser Asn Tyr Ile Leu Ser Thr Leu Ser		
100	105	110
Ala Asp Thr Phe Pro Asn His Gln Asn Phe Asp Tyr Ile Ser Lys Phe		
115	120	125
Asp Ile Ser Ser Asn Ile Leu Lys Glu Met Ile Glu Lys Thr Glu Phe		
130	135	140
Ser Met Gly Lys Gln Asp Val Arg Tyr Tyr Leu Asn Gly Met Leu Leu		
145	150	155 160
Glu Lys Lys Asp Lys Phe Leu Arg Ser Val Ala Thr Asp Gly Tyr Arg		
165	170	175

Leu Ala Ile Ser Tyr Thr Gln Leu Lys Lys Asp Ile Asn Phe Phe Ser
 180 185 190
 Ile Ile Ile Pro Asn Lys Ala Val Met Glu Leu Leu Lys Leu Leu Asn
 195 200 205
 Thr Gln Pro Gln Leu Leu Asn Ile Leu Ile Gly Ser Asn Ser Ile Arg
 210 215 220
 Ile Tyr Thr Lys Asn Leu Ile Phe Thr Thr Gln Leu Ile Glu Gly Glu
 225 230 235 240
 Tyr Pro Asp Tyr Lys Ser Val Leu Phe Lys Glu Lys Lys Asn Pro Ile
 245 250 255
 Ile Thr Asn Ser Ile Leu Leu Lys Lys Ser Leu Leu Arg Val Ala Ile
 260 265 270
 Leu Ala His Glu Lys Phe Cys Gly Ile Glu Ile Lys Ile Glu Asn Gly
 275 280 285
 Lys Phe Lys Val Leu Ser Asp Asn Gln Glu Glu Glu Thr Ala Glu Asp
 290 295 300
 Leu Phe Glu Ile Asp Tyr Phe Gly Glu Lys Ile Glu Ile Ser Ile Asn
 305 310 315 320
 Val Tyr Tyr Leu Leu Asp Val Ile Asn Asn Ile Lys Ser Glu Asn Ile
 325 330 335
 Ala Leu Phe Leu Asn Lys Ser Lys Ser Ser Ile Gln Ile Glu Ala Glu
 340 345 350
 Asn Asn Ser Ser Asn Ala Tyr Val Val Met Leu Leu Lys Arg
 355 360 365

<210> 114

<211> 39

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 114

gtgtggatcc tcgtccccct catgcgcgac caggaaggg

39

<210> 115
 <211> 27
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer

<400> 115
 gtgtggatcc gtggtgacct tagccac

27

<210> 116
 <211> 30
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer

<400> 116
 ttcgtgtccg aggaccttgt ggtccacaac

30

<210> 117
 <211> 3514
 <212> DNA
 <213> Aquifex aeolicus

<400> 117
 atgagtaagg atttcgtcca ccttcacctg cacacccagt tctcactcct ggacggggct 60
 ataaagatag acgagctcgt gaaaaaggca aaggagtatg gatacaaagc tgtcggaatg 120
 tcagaccacg gaaacctctt cggttcgtat aaattctaca aagccctgaa ggcggaagga 180
 attaagccca taatcggcat ggaagcctac tttaccacgg gttcgagggt tgacagaaag 240
 actaaaacga gcgaggacaa cataaccgac aagtacaacc accacctcat acttatagca 300
 aaggacgaaa aggtctaaag aacttaatga agctctcaac cctcgccctac aaagaagggt 360
 ttactacaa acccagaatt gattacgaac tccttgaaaa gtacggggag ggcctaatag 420
 cccttaccgc atgcctgaaa ggtgttccca cctactacgc ttctataaac gaagtgaaaa 480
 aggcgaggga atgggttaaag aagttcaagg atatattcgg agatgacctt tatttagaac 540
 ttcaagcgaa caacattcca gaacaggaag tggcaaacag gaacttaata gagatagcca 600
 aaaagtacga tgtgaaactc atagcgacgc aggacgcccc ctacctcaat cccgaagaca 660
 ggtacgcccc cacggttctt atggcacttc aaatgaaaaa gaccattcac gaactgagtt 720
 cgggaaactt caagtgttca aacgaagacc ttcactttgc tccacccgag tacatgtgga 780
 aaaagtttga aggttaagttc gaaggctggg aaaaggcact cctgaacact ctcgaggtaa 840
 tggaaaagac agcggacagc tttgagatat ttgaaaactc cacctacctc cttcccaagt 900
 acgacgttcc gcccgacaaa acccttgagg aatacctcag agaactcgcg tacaagggtt 960
 taagacagag gatagaaagg ggacaagcta aggatactaa agagtactgg gagaggctcg 1020

```

agtacgaact ggaagttata aacaaaatgg gctttgcggg atactttcttg atagttcagg 1080
acttcataaa ctgggctaag aaaaacgaca tacctgttgg acccggaagg ggaagtgcgtg 1140
gaggttcocct cgtcgcatac gccatcgga taacggacgt tgaccctata aagcacggat 1200
tcctttttga gaggttctta aaccccgaaa gggtttccat gccggatata gacgtggatt 1260
tctgtcagga caacagggaa aaggtcatag agtacgtaag gaacaagtac ggacacgaca 1320
acgtagctca gataatcacc tacaacgtaa tgaaggcgaa gcaaactctg agagacgtcg 1380
caagggccat gggactcccc tactccaccg cggacaaact cgcaaaactc attcctcagg 1440
gggacgttca gggaaactgg ctcagtctgg aagagatgta caaaacgcct gtggaggaac 1500
tccttcagaa gtacggagaa cacagaacgg acatagagga caacgtaaag aagttcagac 1560
agatatgcga agaaagtccg gagataaaac agctcgttga gacggccctg aagcttgaag 1620
gtctcacgag acacacctcc ctccacgccg cgggagtggt tatagcacca aagcccttga 1680
gcgagctcgt tccccctac tacgataaag agggcggaagt cgcaaccag tacgacatgg 1740
ttcagctcga agaactcggc ctctgaaga tggacttctt cggactcaaa accctcacag 1800
aactgaaact catgaaagaa ctcataaagg aaagacacgg agtggaata aacttccttg 1860
aacttcccc tgcgacccg aaagtttaca aactccttca ggaaggaaaa accacgggag 1920
tgttccagct cgaaagcagg ggaatgaaag aactcctgaa gaaactaaag cccgacagct 1980
ttgacgacat cgttgcggtc ctgcactct acagaccgg acctctaaag agcggactcg 2040
ttgacacata cattaagaga aagcacggaa aagaaccgt tgagtacccc tcccggagc 2100
ttgaaccgt ccttaaggaa acctacggag taatcgttta tcaggaacag gtgatgaaga 2160
tgtctcagat actttccggc tttactccc gagaggcgga taccctcaga aaggcgatag 2220
gtaagaagaa agcggattta atggctcaga tgaaagacaa gttcatacag ggagcggtg 2280
aaaggggata ccctgaagaa aagataagga agctctggga agacatagag aagttcgctt 2340
cctactcctt caacaagtct cactcggtag cttacgggta catctcctac tggaccgcct 2400
acgttaaagc ccactatccc gcggagttct tcgcggtaaa actcacaact gaaaagaacg 2460
acaacaagtt cctcaacctc ataaaagacg cttaaactctt cggatttgag atacttcccc 2520
ccgacataaa caagagtgat gtaggattta cgatagaagg tgaaaacagg ataaggttcg 2580
ggcttgcgag gataaaggga gtgggagagg aaactgctaa gataatcggt gaagctagaa 2640
agaagtataa gcagttcaaa gggcttgcg acttcataaa caaaaccaag aacaggaaga 2700
taaacaagaa agtcgtggaa gcactcgtaa aggcaggggc ttttgacttt actaagaaaa 2760
agaggaaaga actactcgt aaagtggcaa actctgaaaa agcattaatg gctacacaaa 2820
actccctttt cgggtgcacc aaagaagaag tggaagaact cgaccctta aagcttgaaa 2880
aggaagttct cggttttttac atttcagggc accccttga caactacgaa aagctcctca 2940
agaaccgcta cacacccatt gaagatttag aagagtggga caaggaaagc gaagcgggtc 3000
ttacaggagt tatcacggaa ctcaaagtaa aaaagacgaa aaacggagat tacatggcgg 3060
tcttcaacct cgttgacaag acgggactaa tagagtgtgt cgtcttccc ggagtttacg 3120
aagaggcaaa ggaactgata gaagaggaca gagtagtggt agtcaaagg tttctggacg 3180
aggaccttga aacggaaaat gtcaagttcg tggtgaaaga ggttttctcc cctgaggagt 3240
tcgcaaagga gatgaggaat accctttata tattcttaaa aagagagcaa gccctaaacg 3300
gcgttgccga aaaactaaag ggaattattg aaaacaacag gacggaggac ggatacaact 3360
tggttctcac ggttgatctg ggagactact togttgattt agcactccca caagatatga 3420
aactaaaggc tgacagaaag gttgtagagg agatagaaaa actgggagtg aaggtcataa 3480
tttagtaaat aacccttact tccgagtagt cccc 3514

```

<210> 118

<211> 1161

<212> PRT

<213> Aquifex aeolicus

<400> 118

Met Ser Lys Asp Phe Val His Leu His Leu His Thr Gln Phe Ser Leu
1 5 10 15

Leu Asp Gly Ala Ile Lys Ile Asp Glu Leu Val Lys Lys Ala Lys Glu
20 25 30

Tyr Gly Tyr Lys Ala Val Gly Met Ser Asp His Gly Asn Leu Phe Gly
35 40 45

Ser Tyr Lys Phe Tyr Lys Ala Leu Lys Ala Glu Gly Ile Lys Pro Ile
50 55 60

Ile Gly Met Glu Ala Tyr Phe Thr Thr Gly Ser Arg Phe Asp Arg Lys
65 70 75 80

Thr Lys Thr Ser Glu Asp Asn Ile Thr Asp Lys Tyr Asn His His Leu
85 90 95

Ile Leu Ile Ala Lys Asp Asp Lys Gly Leu Lys Asn Leu Met Lys Leu
100 105 110

Ser Thr Leu Ala Tyr Lys Glu Gly Phe Tyr Tyr Lys Pro Arg Ile Asp
115 120 125

Tyr Glu Leu Leu Glu Lys Tyr Gly Glu Gly Leu Ile Ala Leu Thr Ala
130 135 140

Cys Leu Lys Gly Val Pro Thr Tyr Tyr Ala Ser Ile Asn Glu Val Lys
145 150 155 160

Lys Ala Glu Glu Trp Val Lys Lys Phe Lys Asp Ile Phe Gly Asp Asp
165 170 175

Leu Tyr Leu Glu Leu Gln Ala Asn Asn Ile Pro Glu Gln Glu Val Ala
180 185 190

Asn Arg Asn Leu Ile Glu Ile Ala Lys Lys Tyr Asp Val Lys Leu Ile
195 200 205

Ala Thr Gln Asp Ala His Tyr Leu Asn Pro Glu Asp Arg Tyr Ala His
210 215 220

Thr Val Leu Met Ala Leu Gln Met Lys Lys Thr Ile His Glu Leu Ser
225 230 235 240

Ser Gly Asn Phe Lys Cys Ser Asn Glu Asp Leu His Phe Ala Pro Pro

001211-49697260

245	250	255
Glu Tyr Met Trp Lys Lys Phe Glu Gly Lys Phe Glu Gly Trp Glu Lys		
260	265	270
Ala Leu Leu Asn Thr Leu Glu Val Met Glu Lys Thr Ala Asp Ser Phe		
275	280	285
Glu Ile Phe Glu Asn Ser Thr Tyr Leu Leu Pro Lys Tyr Asp Val Pro		
290	295	300
Pro Asp Lys Thr Leu Glu Glu Tyr Leu Arg Glu Leu Ala Tyr Lys Gly		
305	310	315
Leu Arg Gln Arg Ile Glu Arg Gly Gln Ala Lys Asp Thr Lys Glu Tyr		
325	330	335
Trp Glu Arg Leu Glu Tyr Glu Leu Glu Val Ile Asn Lys Met Gly Phe		
340	345	350
Ala Gly Tyr Phe Leu Ile Val Gln Asp Phe Ile Asn Trp Ala Lys Lys		
355	360	365
Asn Asp Ile Pro Val Gly Pro Gly Arg Gly Ser Ala Gly Gly Ser Leu		
370	375	380
Val Ala Tyr Ala Ile Gly Ile Thr Asp Val Asp Pro Ile Lys His Gly		
385	390	395
Phe Leu Phe Glu Arg Phe Leu Asn Pro Glu Arg Val Ser Met Pro Asp		
405	410	415
Ile Asp Val Asp Phe Cys Gln Asp Asn Arg Glu Lys Val Ile Glu Tyr		
420	425	430
Val Arg Asn Lys Tyr Gly His Asp Asn Val Ala Gln Ile Ile Thr Tyr		
435	440	445
Asn Val Met Lys Ala Lys Gln Thr Leu Arg Asp Val Ala Arg Ala Met		
450	455	460
Gly Leu Pro Tyr Ser Thr Ala Asp Lys Leu Ala Lys Leu Ile Pro Gln		
465	470	475
Gly Asp Val Gln Gly Thr Trp Leu Ser Leu Glu Glu Met Tyr Lys Thr		
485	490	495
Pro Val Glu Glu Leu Leu Gln Lys Tyr Gly Glu His Arg Thr Asp Ile		

500	505	510
Glu Asp Asn Val Lys Lys Phe Arg Gln Ile Cys Glu Glu Ser Pro Glu		
515	520	525
Ile Lys Gln Leu Val Glu Thr Ala Leu Lys Leu Glu Gly Leu Thr Arg		
530	535	540
His Thr Ser Leu His Ala Ala Gly Val Val Ile Ala Pro Lys Pro Leu		
545	550	555
Ser Glu Leu Val Pro Leu Tyr Tyr Asp Lys Glu Gly Glu Val Ala Thr		
565	570	575
Gln Tyr Asp Met Val Gln Leu Glu Glu Leu Gly Leu Leu Lys Met Asp		
580	585	590
Phe Leu Gly Leu Lys Thr Leu Thr Glu Leu Lys Leu Met Lys Glu Leu		
595	600	605
Ile Lys Glu Arg His Gly Val Asp Ile Asn Phe Leu Glu Leu Pro Leu		
610	615	620
Asp Asp Pro Lys Val Tyr Lys Leu Leu Gln Glu Gly Lys Thr Thr Gly		
625	630	635
Val Phe Gln Leu Glu Ser Arg Gly Met Lys Glu Leu Leu Lys Lys Leu		
645	650	655
Lys Pro Asp Ser Phe Asp Asp Ile Val Ala Val Leu Ala Leu Tyr Arg		
660	665	670
Pro Gly Pro Leu Lys Ser Gly Leu Val Asp Thr Tyr Ile Lys Arg Lys		
675	680	685
His Gly Lys Glu Pro Val Glu Tyr Pro Phe Pro Glu Leu Glu Pro Val		
690	695	700
Leu Lys Glu Thr Tyr Gly Val Ile Val Tyr Gln Glu Gln Val Met Lys		
705	710	715
Met Ser Gln Ile Leu Ser Gly Phe Thr Pro Gly Glu Ala Asp Thr Leu		
725	730	735
Arg Lys Ala Ile Gly Lys Lys Lys Ala Asp Leu Met Ala Gln Met Lys		
740	745	750
Asp Lys Phe Ile Gln Gly Ala Val Glu Arg Gly Tyr Pro Glu Glu Lys		

001221-49697250

755		760		765
Ile Arg Lys Leu Trp Glu Asp Ile Glu Lys Phe Ala Ser Tyr Ser Phe				
770		775		780
Asn Lys Ser His Ser Val Ala Tyr Gly Tyr Ile Ser Tyr Trp Thr Ala				
785		790		795 800
Tyr Val Lys Ala His Tyr Pro Ala Glu Phe Phe Ala Val Lys Leu Thr				
	805		810	815
Thr Glu Lys Asn Asp Asn Lys Phe Leu Asn Leu Ile Lys Asp Ala Lys				
	820		825	830
Leu Phe Gly Phe Glu Ile Leu Pro Pro Asp Ile Asn Lys Ser Asp Val				
	835		840	845
Gly Phe Thr Ile Glu Gly Glu Asn Arg Ile Arg Phe Gly Leu Ala Arg				
	850		855	860
Ile Lys Gly Val Gly Glu Glu Thr Ala Lys Ile Ile Val Glu Ala Arg				
	865		870	875 880
Lys Lys Tyr Lys Gln Phe Lys Gly Leu Ala Asp Phe Ile Asn Lys Thr				
	885		890	895
Lys Asn Arg Lys Ile Asn Lys Lys Val Val Glu Ala Leu Val Lys Ala				
	900		905	910
Gly Ala Phe Asp Phe Thr Lys Lys Lys Arg Lys Glu Leu Leu Ala Lys				
	915		920	925
Val Ala Asn Ser Glu Lys Ala Leu Met Ala Thr Gln Asn Ser Leu Phe				
	930		935	940
Gly Ala Pro Lys Glu Glu Val Glu Glu Leu Asp Pro Leu Lys Leu Glu				
	945		950	955 960
Lys Glu Val Leu Gly Phe Tyr Ile Ser Gly His Pro Leu Asp Asn Tyr				
	965		970	975
Glu Lys Leu Leu Lys Asn Arg Tyr Thr Pro Ile Glu Asp Leu Glu Glu				
	980		985	990
Trp Asp Lys Glu Ser Glu Ala Val Leu Thr Gly Val Ile Thr Glu Leu				
	995		1000	1005
Lys Val Lys Lys Thr Lys Asn Gly Asp Tyr Met Ala Val Phe Asn Leu				

1010 1015 1020

Val Asp Lys Thr Gly Leu Ile Glu Cys Val Val Phe Pro Gly Val Tyr
1025 1030 1035 1040

Glu Glu Ala Lys Glu Leu Ile Glu Glu Asp Arg Val Val Val Val Lys
1045 1050 1055

Gly Phe Leu Asp Glu Asp Leu Glu Thr Glu Asn Val Lys Phe Val Val
1060 1065 1070

Lys Glu Val Phe Ser Pro Glu Glu Phe Ala Lys Glu Met Arg Asn Thr
1075 1080 1085

Leu Tyr Ile Phe Leu Lys Arg Glu Gln Ala Leu Asn Gly Val Ala Glu
1090 1095 1100

Lys Leu Lys Gly Ile Ile Glu Asn Asn Arg Thr Glu Asp Gly Tyr Asn
1105 1110 1115 1120

Leu Val Leu Thr Val Asp Leu Gly Asp Tyr Phe Val Asp Leu Ala Leu
1125 1130 1135

Pro Gln Asp Met Lys Leu Lys Ala Asp Arg Lys Val Val Glu Glu Ile
1140 1145 1150

Glu Lys Leu Gly Val Lys Val Ile Ile
1155 1160

<210> 119
<211> 2408
<212> DNA
<213> Aquifex aeolicus

<400> 119

atgaactacg ttcccttcgc gagaaagtac agaccgaaat tcttcaggga agtaatagga 60
caggaagctc ccgtaaggat actcaaaaac gctataaaaa acgacagagt ggctcacgcc 120
tacctctttg ccggaccgag gggggttggg aagacgacta ttgcaagaat tctcgcaaaa 180
gctttgaact gtaaaaatcc ctccaaagggt gagccctgcg gtgagtgcga aaactgcagg 240
gagatagaca ggggtgtgtt ccctgactta attgaaatgg atgccgcctc aaacaggggt 300
atagacgacg taagggcatt aaaagaagcg gtcaattaca aacctataaa aggaaagtac 360
aaggtttaca taatagacga agctcacatg ctcacgaaag aagctttcaa cgctctctta 420
aaaaccctcg aagagcccc tcccagaact gttttcgctc tttgtaccac ggagtacgac 480
aaaattcttc ccacgatact ctcaagggtg cagaggataa tcttctcaaa ggtaagaaag 540
gaaaaagtaa tagagtatct aaaaaagata tgtgaaaagg aagggattga gtgcgaagag 600
ggagcccttg aggttctggc tcatgcctct gaagggtgca tgagggatgc agcctctctc 660
ctggaccagg cgagcgttta cggggaaggc agggtaacaa aagaagtagt ggagaacttc 720


```

ctcgggaattc tcagtcagga aagcggttagg agttttctga aattgcttct gaactcagaa 780
gtggacgaag ctataaagtt cctcagagaa ctctcagaaa agggctacaa cctgaccaag 840
ttttgggaga tgttagaaga ggaagtgaga aacgcaatth tagtaaagag cctgaaaaat 900
cccgaaagcg tggttcagaa ctggcaggat tacgaagact tcaaagacta ccctctggaa 960
gccctcctct acgttgagaa cctgataaac aggggttaaag ttgaagcgag aacgagagaa 1020
cccttaagag cctttgaact cgcggtaata aagagcctta tagtcaaaga cataattccc 1080
gtatcccagc tcggaagtgt ggtaaaggaa accaaaaagg aagaaaagaa agttgaagta 1140
aaagaagagc caaaagtaaa agaagaaaaa ccaaaggagc aggaagagga caggttccag 1200
aaagtthttaa acgctgtgga cggcaaaatc cttaaaagaa tacttgaagg ggcaaaaagg 1260
gaagaaagag acggaaaaat cgtcctaaag atagaagcct cttatctgag aaccatgaaa 1320
aaggaaattg actcactaaa ggagacttht cctthtttag agtttgaacc cgtggaggat 1380
aaaaaaaaac ctcagaagtc cagcgggacg aggctgtht aaaggtaaag gagctcttca 1440
atgcaaaaat actcaaagta cgaagtaaaa gctaaggta taaaggtag aatgccgtg 1500
gaagagatag ggctgttht cgcactaata gacggcttgc ccaggtagc actcacgag 1560
acgaaggaag agggaaaggg agaagtht cgttttagcga ctcttataa agtcaaggaa 1620
ttgatggaag ctatggaggg tatgaaaaa cacataaagg atttagaat cctcgagag 1680
acggatgagg atttaactth ttaaagtatg ggtgtatctg agcaaaggth taagctaaaa 1740
acaaacctga aacctgcagg ggaccagccg aaagccataa aaaaactcct tgaaaacct 1800
aggaaaggcg taaaagaaca aacacttht cgtggtcacgg gaagcggaaa gactthtact 1860
ctagcaaacg taatagcgaa gtacaacaaa ccaactcttg tggtagttca caacaaaatt 1920
ctcggggcac agctatacag ggagthttaa gaactattcc ctgaaaacgc tgtagagtac 1980
thtgtctctt actacgacta ttaccaacct gaagcctaca ttcccgaaaa agatttatac 2040
atagaaaagg acgcgagtat aaacgaaagc tggaacgtht cagacactcc gccacgat 2100
ccgttctaga aaggagggac gttatagtag ttgcttcagt ttcttgcata tacggactcg 2160
ggaaacctga gcactacgaa aacctgagga taaaactcca aagggggaata agactgaact 2220
tgagtaagct cctgaggaaa ctggttgagc taggatatca gagaaatgac thtgccataa 2280
agagggtac cttctcggtt aggggagacg tggttgagat agtcccttct cacacggaag 2340
attacctcgt gagggtagag thtctgggacg acgaagttga aagaatagtc ctcatggagc 2400
ctctgaac 2408

```

<210> 120

<211> 473

<212> PRT

<213> Aquifex aeolicus

<400> 120

Met Asn Tyr Val Pro Phe Ala Arg Lys Tyr Arg Pro Lys Phe Phe Arg

1 5 10 15

Glu Val Ile Gly Gln Glu Ala Pro Val Arg Ile Leu Lys Asn Ala Ile

20 25 30

Lys Asn Asp Arg Val Ala His Ala Tyr Leu Phe Ala Gly Pro Arg Gly

35 40 45

Val Gly Lys Thr Thr Ile Ala Arg Ile Leu Ala Lys Ala Leu Asn Cys

50 55 60

Lys	Asn	Pro	Ser	Lys	Gly	Glu	Pro	Cys	Gly	Glu	Cys	Glu	Asn	Cys	Arg		
65					70					75					80		
Glu	Ile	Asp	Arg	Gly	Val	Phe	Pro	Asp	Leu	Ile	Glu	Met	Asp	Ala	Ala		
				85					90					95			
Ser	Asn	Arg	Gly	Ile	Asp	Asp	Val	Arg	Ala	Leu	Lys	Glu	Ala	Val	Asn		
			100					105					110				
Tyr	Lys	Pro	Ile	Lys	Gly	Lys	Tyr	Lys	Val	Tyr	Ile	Ile	Asp	Glu	Ala		
	115						120					125					
His	Met	Leu	Thr	Lys	Glu	Ala	Phe	Asn	Ala	Leu	Leu	Lys	Thr	Leu	Glu		
	130					135					140						
Glu	Pro	Pro	Pro	Arg	Thr	Val	Phe	Val	Leu	Cys	Thr	Thr	Glu	Tyr	Asp		
145					150					155					160		
Lys	Ile	Leu	Pro	Thr	Ile	Leu	Ser	Arg	Cys	Gln	Arg	Ile	Ile	Phe	Ser		
				165					170					175			
Lys	Val	Arg	Lys	Glu	Lys	Val	Ile	Glu	Tyr	Leu	Lys	Lys	Ile	Cys	Glu		
			180					185					190				
Lys	Glu	Gly	Ile	Glu	Cys	Glu	Glu	Gly	Ala	Leu	Glu	Val	Leu	Ala	His		
	195						200					205					
Ala	Ser	Glu	Gly	Cys	Met	Arg	Asp	Ala	Ala	Ser	Leu	Leu	Asp	Gln	Ala		
	210					215					220						
Ser	Val	Tyr	Gly	Glu	Gly	Arg	Val	Thr	Lys	Glu	Val	Val	Glu	Asn	Phe		
225					230					235				240			
Leu	Gly	Ile	Leu	Ser	Gln	Glu	Ser	Val	Arg	Ser	Phe	Leu	Lys	Leu	Leu		
			245						250					255			
Leu	Asn	Ser	Glu	Val	Asp	Glu	Ala	Ile	Lys	Phe	Leu	Arg	Glu	Leu	Ser		
		260						265					270				
Glu	Lys	Gly	Tyr	Asn	Leu	Thr	Lys	Phe	Trp	Glu	Met	Leu	Glu	Glu	Glu		
	275						280					285					
Val	Arg	Asn	Ala	Ile	Leu	Val	Lys	Ser	Leu	Lys	Asn	Pro	Glu	Ser	Val		
	290					295					300						
Val	Gln	Asn	Trp	Gln	Asp	Tyr	Glu	Asp	Phe	Lys	Asp	Tyr	Pro	Leu	Glu		
305				310						315				320			

Ala Leu Leu Tyr Val Glu Asn Leu Ile Asn Arg Gly Lys Val Glu Ala
325 330 335

Arg Thr Arg Glu Pro Leu Arg Ala Phe Glu Leu Ala Val Ile Lys Ser
340 345 350

Leu Ile Val Lys Asp Ile Ile Pro Val Ser Gln Leu Gly Ser Val Val
355 360 365

Lys Glu Thr Lys Lys Glu Glu Lys Lys Val Glu Val Lys Glu Glu Pro
370 375 380

Lys Val Lys Glu Glu Lys Pro Lys Glu Gln Glu Glu Asp Arg Phe Gln
385 390 395 400

Lys Val Leu Asn Ala Val Asp Gly Lys Ile Leu Lys Arg Ile Leu Glu
405 410 415

Gly Ala Lys Arg Glu Glu Arg Asp Gly Lys Ile Val Leu Lys Ile Glu
420 425 430

Ala Ser Tyr Leu Arg Thr Met Lys Lys Glu Phe Asp Ser Leu Lys Glu
435 440 445

Thr Phe Pro Phe Leu Glu Phe Glu Pro Val Glu Asp Lys Lys Lys Pro
450 455 460

Gln Lys Ser Ser Gly Thr Arg Leu Phe
465 470

<210> 121

<211> 1090

<212> DNA

<213> Aquifex aeolicus

<400> 121

atgCGcgTta aggtggacag ggaggagctt gaagaggttc ttaaaaaagc aagagaaagc 60
acggaaaaaa aagccgcact cccgatactc gcgaacttct tactctccgc aaaagaggaa 120
aacttaatcg taagggcaac ggacttggaa aactaccttg tagtctccgt aaagggggag 180
gttgaagagg aaggagaggt ttgCGtccac tctcaaaaac tctacgatat agtcaagaac 240
ttaaatccg cttacgttta ccttcatacg gaaggTgaaa aactCGtcat aacgggagga 300
aagagtacgt acaaaactcc gacagctccc gcggaggact ttcccgaatt tccagaaatc 360
gtagaaggag gagaaacact ttcgggaaac cttctcgtta acggaataga aaaggtagag 420
tacGCCatag cgaaggaaga agcgaacata gcccttcagg gaatgtatct gagaggatac 480
gaggacagaa ttcactttgt gttcggacgg tcacaggctt gcactttatg aacctctacg 540
taaacattga aaagagtga gacgagtctt ttgcttactt ctccactccc gagtggaaac 600

tcgccggttag ctccctggaag gagaattccc ggactacatg agtgtcatcc ctgaggagtt 660
 ttccggcgga gttctgtttg agacagagga agtcttaaag gttttaaaga ggttgaaggc 720
 ttttaagcgaa ggaaaagtgtt ttcccgtgaa gattacctta agcgaaaacc ttgccatctt 780
 tgagttcgcg gatccggagt tcggagaagc gagagaggaa attgaagtgg agtacacggg 840
 agagcccttt gagataggat tcaacggaaa taccttatgg aggcgcttga cgcctacgac 900
 agcgaaagag tgtggttcaa gttcacaacc cccgacacgg ccactttatt ggaggctgaa 960
 gattacgaaa aggaacctta caagtgcata ataatgccga tgagggtgta gccatgaaaa 1020
 aagctttaat cttttttattg agcttgagcc ttttaattcc tgcgttttagc gaagccaaac 1080
 ccaagtcttc 1090

<210> 122

<211> 363

<212> PRT

<213> Aquifex aeolicus

<400> 122

Met Arg Val Lys Val Asp Arg Glu Glu Leu Glu Glu Val Leu Lys Lys
 1 5 10 15

Ala Arg Glu Ser Thr Glu Lys Lys Ala Ala Leu Pro Ile Leu Ala Asn
 20 25 30

Phe Leu Leu Ser Ala Lys Glu Glu Asn Leu Ile Val Arg Ala Thr Asp
 35 40 45

Leu Glu Asn Tyr Leu Val Val Ser Val Lys Gly Glu Val Glu Glu Glu
 50 55 60

Gly Glu Val Cys Val His Ser Gln Lys Leu Tyr Asp Ile Val Lys Asn
 65 70 75 80

Leu Asn Ser Ala Tyr Val Tyr Leu His Thr Glu Gly Glu Lys Leu Val
 85 90 95

Ile Thr Gly Gly Lys Ser Thr Tyr Lys Leu Pro Thr Ala Pro Ala Glu
 100 105 110

Asp Phe Pro Glu Phe Pro Glu Ile Val Glu Gly Gly Glu Thr Leu Ser
 115 120 125

Gly Asn Leu Leu Val Asn Gly Ile Glu Lys Val Glu Tyr Ala Ile Ala
 130 135 140

Lys Glu Glu Ala Asn Ile Ala Leu Gln Gly Met Tyr Leu Arg Gly Tyr
 145 150 155 160

Glu Asp Arg Ile His Phe Val Gly Ser Asp Gly His Arg Leu Ala Leu

	165		170		175
Tyr Glu Pro Leu Gly Glu Phe Ser Lys Glu Leu Leu Ile Pro Arg Lys					
	180		185		190
Ser Leu Lys Val Leu Lys Lys Leu Ile Thr Gly Ile Glu Asp Val Asn					
	195		200		205
Ile Glu Lys Ser Glu Asp Glu Ser Phe Ala Tyr Phe Ser Thr Pro Glu					
	210		215		220
Trp Lys Leu Ala Val Arg Leu Leu Glu Gly Glu Phe Pro Asp Tyr Met					
	225		230		240
Ser Val Ile Pro Glu Glu Phe Ser Ala Glu Val Leu Phe Glu Thr Glu					
	245		250		255
Glu Val Leu Lys Val Leu Lys Arg Leu Lys Ala Leu Ser Glu Gly Lys					
	260		265		270
Val Phe Pro Val Lys Ile Thr Leu Ser Glu Asn Leu Ala Ile Phe Glu					
	275		280		285
Phe Ala Asp Pro Glu Phe Gly Glu Ala Arg Glu Glu Ile Glu Val Glu					
	290		295		300
Tyr Thr Gly Glu Pro Phe Glu Ile Gly Phe Asn Gly Lys Tyr Leu Met					
	305		310		320
Glu Ala Leu Asp Ala Tyr Asp Ser Glu Arg Val Trp Phe Lys Phe Thr					
	325		330		335
Thr Pro Asp Thr Ala Thr Leu Leu Glu Ala Glu Asp Tyr Glu Lys Glu					
	340		345		350
Pro Tyr Lys Cys Ile Ile Met Pro Met Arg Val					
	355		360		

<210> 123

<211> 1093

<212> DNA

<213> Aquifex aeolicus

<400> 123

gtggaaacca caatattcca gttccagaaa actttttttca caaaacctcc gaaggagagg 60
gtcttcgtcc ttcattggaga agagcagtat ctcataagaa cctttttgtc taagctgaag 120
gaaaagtacg gggagaatta cacggttctg tgggggggatg agataagcga ggaggaattc 180

tacactgccc tttccgagac cagtatatc ggcggttcaa aggaaaaagc ggtggtcatt 240
 tacaacttcg gggatttcct gaagaagctc ggaaggaaga aaaaggaaaa agaaaggctt 300
 ataaaagtc tcaaaaacgt aaagagtaac tacgtattta tagtgtacga tgcgaaactc 360
 cagaaacagg aactttcttc ggaacctctg aaatccgtag cgtctttcgg cggatatagt 420
 gtagcaaaca ggctgagcaa ggagaggata aaacagctcg tccttaagaa gttcaaagaa 480
 aaagggataa acgtagaaaa cgatgccctt gaataccttc tccagctcac gggttacaac 540
 ttgatggagc tcaaacttga ggttgaaaaa ctgatagatt acgcaagtga aaagaaaatt 600
 ttaacactcg atgaggtaaa gagagtagcc ttctcagtct cagaaaacgt aaacgtattt 660
 gagttcggtg atttactcct cttaaaagat tacgaaaagg ctcttaaagt tttggactcc 720
 ctcatttcct tcggaatata cccctccag attatgaaaa tcctgtcctc ctatgtctta 780
 aaactttaca ccctcaagag gcttgaagag aaggagagg acctgaataa ggcatggaa 840
 agcgtgggaa taaagaacaa ctttctcaag atgaagttca aatcttactt aaaggcaaac 900
 tctaaagagg acttgaagaa cctaactcct tccctccaga ggatagacgc tttttctaaa 960
 ctttactttc aggacacagt gcagttgctg gggatttctt gacctcaaga ctggagaggg 1020
 aagttgtgaa aaatacttct catgggtggat aatctttttt atgaagtttg cggtttgcgt 1080
 ttttcccggt tct 1093

<210> 124

<211> 350

<212> PRT

<213> Aquifex aeolicus

<400> 124

Val	Glu	Thr	Thr	Ile	Phe	Gln	Phe	Gln	Lys	Thr	Phe	Phe	Thr	Lys	Pro
1				5					10					15	
Pro	Lys	Glu	Arg	Val	Phe	Val	Leu	His	Gly	Glu	Glu	Gln	Tyr	Leu	Ile
			20					25						30	
Arg	Thr	Phe	Leu	Ser	Lys	Leu	Lys	Glu	Lys	Tyr	Gly	Glu	Asn	Tyr	Thr
		35					40					45			
Val	Leu	Trp	Gly	Asp	Glu	Ile	Ser	Glu	Glu	Glu	Phe	Tyr	Thr	Ala	Leu
	50					55					60				
Ser	Glu	Thr	Ser	Ile	Phe	Gly	Gly	Ser	Lys	Glu	Lys	Ala	Val	Val	Ile
	65				70				75					80	
Tyr	Asn	Phe	Gly	Asp	Phe	Leu	Lys	Lys	Leu	Gly	Arg	Lys	Lys	Lys	Glu
				85					90					95	
Lys	Glu	Arg	Leu	Ile	Lys	Val	Leu	Arg	Asn	Val	Lys	Ser	Asn	Tyr	Val
			100				105						110		
Phe	Ile	Val	Tyr	Asp	Ala	Lys	Leu	Gln	Lys	Gln	Glu	Leu	Ser	Ser	Glu
		115					120					125			

Pro Leu Lys Ser Val Ala Ser Phe Gly Gly Ile Val Val Ala Asn Arg
 130 135 140

Leu Ser Lys Glu Arg Ile Lys Gln Leu Val Leu Lys Lys Phe Lys Glu
 145 150 155 160

Lys Gly Ile Asn Val Glu Asn Asp Ala Leu Glu Tyr Leu Leu Gln Leu
 165 170 175

Thr Gly Tyr Asn Leu Met Glu Leu Lys Leu Glu Val Glu Lys Leu Ile
 180 185 190

Asp Tyr Ala Ser Glu Lys Lys Ile Leu Thr Leu Asp Glu Val Lys Arg
 195 200 205

Val Ala Phe Ser Val Ser Glu Asn Val Asn Val Phe Glu Phe Val Asp
 210 215 220

Leu Leu Leu Leu Lys Asp Tyr Glu Lys Ala Leu Lys Val Leu Asp Ser
 225 230 235 240

Leu Ile Ser Phe Gly Ile His Pro Leu Gln Ile Met Lys Ile Leu Ser
 245 250 255

Ser Tyr Ala Leu Lys Leu Tyr Thr Leu Lys Arg Leu Glu Glu Lys Gly
 260 265 270

Glu Asp Leu Asn Lys Ala Met Glu Ser Val Gly Ile Lys Asn Asn Phe
 275 280 285

Leu Lys Met Lys Phe Lys Ser Tyr Leu Lys Ala Asn Ser Lys Glu Asp
 290 295 300

Leu Lys Asn Leu Ile Leu Ser Leu Gln Arg Ile Asp Ala Phe Ser Lys
 305 310 315 320

Leu Tyr Phe Gln Asp Thr Val Gln Leu Leu Arg Asp Phe Leu Thr Ser
 325 330 335

Arg Leu Glu Arg Glu Val Val Lys Asn Thr Ser His Gly Gly
 340 345 350

<210> 125

<211> 1051

<212> DNA

<213> Aquifex aeolicus

<400> 125

```

atggaaaaag tttttttgga aaaactccag aaaaccttgc acatacccgaggactcctt 60
ttttacggca aagaaggaag cggaaagacg aaaacagctt ttgaatttgc aaaaggtatt 120
ttatgtaagg aaaacgtacc tggggatgcg gaagttgtcc ctcttgcaaa cacgtaaacy 180
agctggagga agccttcttt aaaggagaaa tagaagactt taaagtttat aagacaagga 240
cggtaaaaaag cacttcgttt accttatggg cgaacatccc gactttgtgg taataatccc 300
gagcggacat tacataaaga tagaacagat aagggaagtt aagaactttg cctatgtgaa 360
gcccgcacta agcaggagaa aagtaattat aatagacgac gccacgcga tgacctctca 420
ggcggcaaac gctcttttaa aggtattgga agagccacct gcggacacca ctttatctt 480
gaccacgaac aggcgttctg caatcctgcc gactatcctc tccagaactt ttcaagtgga 540
gttcaagggc ttttcagtaa aagaggttat ggaaatagcg aaagtagacg aggaaatagc 600
gaaactctct ggaggcagtc taaaaagggc tatcttacta aaggaaaaca aagatatcct 660
aaacaaagta aaggaattct tggaaaacga gccgttaaaa gtttacaagc ttgcaagtga 720
attcgaaaaag tgggaacctg aaaagcaaaa actcttcctt gaaattatgg aagaattggt 780
atctcaaaaa ttgaccgaag agaaaaaaga caattacacc taccttcttg atacgatcag 840
actctttaa gacggactcg caaggggtgt aaacgaacct ctgtggctgt ttacgttagc 900
cgttcaggcg gattaataaa ccgttattga ttccgtaaca tttaaacctt aatctaaatt 960
atgagagcct ttgaaggagg tctggtatgg aaaatttgaa gattagatat atagatacga 1020
ggaagatagg aaccgtgagc ggtgtaaaag t 1051

```

<210> 126

<211> 305

<212> PRT

<213> Aquifex aeolicus

<400> 126

```

Met Glu Lys Val Phe Leu Glu Lys Leu Gln Lys Thr Leu His Ile Pro
  1                      5                      10                      15

Gly Gly Leu Leu Phe Tyr Gly Lys Glu Gly Ser Gly Lys Thr Lys Thr
      20                      25                      30

Ala Phe Glu Phe Ala Lys Gly Ile Leu Cys Lys Glu Asn Val Pro Trp
      35                      40                      45

Gly Cys Gly Ser Cys Pro Ser Cys Lys His Val Asn Glu Leu Glu Glu
      50                      55                      60

Ala Phe Phe Lys Gly Glu Ile Glu Asp Phe Lys Val Tyr Lys Asp Lys
      65                      70                      75                      80

Asp Gly Lys Lys His Phe Val Tyr Leu Met Gly Glu His Pro Asp Phe
      85                      90                      95

Val Val Ile Ile Pro Ser Gly His Tyr Ile Lys Ile Glu Gln Ile Arg
      100                      105                      110

```


Glu Val Lys Asn Phe Ala Tyr Val Lys Pro Ala Leu Ser Arg Arg Lys
 115 120 125

Val Ile Ile Ile Asp Asp Ala His Ala Met Thr Ser Gln Ala Ala Asn
 130 135 140

Ala Leu Leu Lys Val Leu Glu Glu Pro Pro Ala Asp Thr Thr Phe Ile
 145 150 155 160

Leu Thr Thr Asn Arg Arg Ser Ala Ile Leu Pro Thr Ile Leu Ser Arg
 165 170 175

Thr Phe Gln Val Glu Phe Lys Gly Phe Ser Val Lys Glu Val Met Glu
 180 185 190

Ile Ala Lys Val Asp Glu Glu Ile Ala Lys Leu Ser Gly Gly Ser Leu
 195 200 205

Lys Arg Ala Ile Leu Leu Lys Glu Asn Lys Asp Ile Leu Asn Lys Val
 210 215 220

Lys Glu Phe Leu Glu Asn Glu Pro Leu Lys Val Tyr Lys Leu Ala Ser
 225 230 235 240

Glu Phe Glu Lys Trp Glu Pro Glu Lys Gln Lys Leu Phe Leu Glu Ile
 245 250 255

Met Glu Glu Leu Val Ser Gln Lys Leu Thr Glu Glu Lys Lys Asp Asn
 260 265 270

Tyr Thr Tyr Leu Leu Asp Thr Ile Arg Leu Phe Lys Asp Gly Leu Ala
 275 280 285

Arg Gly Val Asn Glu Pro Leu Trp Leu Phe Thr Leu Ala Val Gln Ala
 290 295 300

Asp
 305

<210> 127

<211> 630

<212> DNA

<213> Aquifex aeolicus

<400> 127

atgaacttcc tgaaaaagtt ccttttactg agaaaagctc aaaagtctcc ttacttcgaa 60
 gagttctacg aagaaatcga tttgaaccag aagggtgaaag atgcaagggt tgtagttttt 120

gactgcgaag ccacagaact cgacgtaaag aaggcaaaac tcctttcaat aggtgcgggt 180
 gaggttaaaa acctggaaat agacctctct aaatcttttt acgagatact caaaagtgac 240
 gagataaagg cggcgggagat acatggaata accaggggaag acgttgaaaa gtacggaaag 300
 gaaccaaagg aagtaataata cgactttctg aagtacataa aggggaagcgt tctcgttggc 360
 tactacgtga agtttgacgt ctactctgtt gagaagtact ccataaagta cttccagtat 420
 ccaatcatca actacaagtt agacctgttt agtttcgtga agagagagta ccagagtggc 480
 aggagtcttg acgaccttat gaaggaactc ggtgtagaaa taagggcaag gcacaacgcc 540
 cttgaagatg cctacataac cgctcttctt ttcctaaagt acgtttaccc gaacagggag 600
 tacagactaa aggatctccc gattttcctt 630

<210> 128

<211> 210

<212> PRT

<213> Aquifex aeolicus

<400> 128

Met Asn Phe Leu Lys Lys Phe Leu Leu Leu Arg Lys Ala Gln Lys Ser
 1 5 10 15

Pro Tyr Phe Glu Glu Phe Tyr Glu Glu Ile Asp Leu Asn Gln Lys Val
 20 25 30

Lys Asp Ala Arg Phe Val Val Phe Asp Cys Glu Ala Thr Glu Leu Asp
 35 40 45

Val Lys Lys Ala Lys Leu Leu Ser Ile Gly Ala Val Glu Val Lys Asn
 50 55 60

Leu Glu Ile Asp Leu Ser Lys Ser Phe Tyr Glu Ile Leu Lys Ser Asp
 65 70 75 80

Glu Ile Lys Ala Ala Glu Ile His Gly Ile Thr Arg Glu Asp Val Glu
 85 90 95

Lys Tyr Gly Lys Glu Pro Lys Glu Val Ile Tyr Asp Phe Leu Lys Tyr
 100 105 110

Ile Lys Gly Ser Val Leu Val Gly Tyr Tyr Val Lys Phe Asp Val Ser
 115 120 125

Leu Val Glu Lys Tyr Ser Ile Lys Tyr Phe Gln Tyr Pro Ile Ile Asn
 130 135 140

Tyr Lys Leu Asp Leu Phe Ser Phe Val Lys Arg Glu Tyr Gln Ser Gly
 145 150 155 160

Arg Ser Leu Asp Asp Leu Met Lys Glu Leu Gly Val Glu Ile Arg Ala

165

170

175

Arg His Asn Ala Leu Glu Asp Ala Tyr Ile Thr Ala Leu Leu Phe Leu
180 185 190

Lys Tyr Val Tyr Pro Asn Arg Glu Tyr Arg Leu Lys Asp Leu Pro Ile
195 200 205

Phe Leu
210

<210> 129

<211> 526

<212> DNA

<213> Aquifex aeolicus

<400> 129

atgctcaata aggtttttat aataggaaga cttacgggtg accccggttat aacttatcta 60
ccgagcggaa cgcccgtagt agagtttact ctggcttaca acagaaggta taaaaaccag 120
aacggtgaat ttcaggagga aagtcacttc tttgacgtaa aggcgtacgg aaaaatggct 180
gaagactggg ctacacgctt ctcgaaagga tacctcgtag tcgtagaggg aagactctcc 240
caggaaaagt gggagaaaga aggaaagaag ttctcaaagg tcaggataat agcggaaaac 300
gtaagattaa taaacaggcc gaaagggtgct gaacttcaag cagaagaaga ggaggaagtt 360
cctccattg aggaggaaat tgaaaaactc ggtaaagagg aagagaagcc ttttaccgat 420
gaagaggacg aaataccttt ttaattttga ggagggttaa gtatggtagt gagagctcct 480
aagaagaaag tttgtatgta ctgtgaacaa aagagagagc cagatt 526

<210> 130

<211> 147

<212> PRT

<213> Aquifex aeolicus

<400> 130

Met Leu Asn Lys Val Phe Ile Ile Gly Arg Leu Thr Gly Asp Pro Val
1 5 10 15

Ile Thr Tyr Leu Pro Ser Gly Thr Pro Val Val Glu Phe Thr Leu Ala
20 25 30

Tyr Asn Arg Arg Tyr Lys Asn Gln Asn Gly Glu Phe Gln Glu Glu Ser
35 40 45

His Phe Phe Asp Val Lys Ala Tyr Gly Lys Met Ala Glu Asp Trp Ala
50 55 60

Thr Arg Phe Ser Lys Gly Tyr Leu Val Leu Val Glu Gly Arg Leu Ser

65

70

75

80

Gln Glu Lys Trp Glu Lys Glu Gly Lys Lys Phe Ser Lys Val Arg Ile
85 90 95

Ile Ala Glu Asn Val Arg Leu Ile Asn Arg Pro Lys Gly Ala Glu Leu
100 105 110

Gln Ala Glu Glu Glu Glu Glu Val Pro Pro Ile Glu Glu Glu Ile Glu
115 120 125

Lys Leu Gly Lys Glu Glu Glu Lys Pro Phe Thr Asp Glu Glu Asp Glu
130 135 140

Ile Pro Phe
145

<210> 131

<211> 1472

<212> DNA

<213> *Aquifex aeolicus*

<400> 131

atgcaatttg tggataaaact tccctgtgac gaatccgccg agagggcggt tcttggcagt 60
atgcttgaag accccgaaaa catacctctg gtacttgaat accttaaaga agaagacttc 120
tgcatagacg agcacaagct acttttcagg gttcttaca acctctgggc cgagtacggc 180
aataagctcg atttcgtatt aataaaggat caccttgaaa agaaaaactt actccagaaa 240
atacctatag actggctcga agaactctac gaggaggcgg tatcccctga cacgcttgag 300
gaagtctgca aaatagtaaa acaacgttcc gcacagaggg cgataattca actcgggtata 360
gaactcattc acaaaggaaa ggaaaacaaa gactttcaca cattaatcga ggaagcccag 420
agcaggatat tttccatagc ggaaagtgtc acatctacgc agttttacca tgtgaaagac 480
gttgcggaag aagttataga actcatttat aaattcaaaa gctctgacag gctagtcacg 540
ggactcccaa gcggtttcac ggaactcgat cttaaagacga cgggattcca ccctggagac 600
ttaataatac tcgccgcaag acccggtatg gggaaaaccg cctttatgct ctccataatc 660
tacaatctcg caaaagacga gggaaaaccc tcagctgtat tttccttgga aatgagcaag 720
gaacagctcg ttatgagact cctctctatg atgtcggagg tcccactttt caagataagg 780
tctggaagta tatcgaatga agatttaaag aagcttgaag caagcgcaat agaactcgca 840
aagtacgaca tatacctcga cgacacaccc gctctcacta caacggattt aaggataagg 900
gcaagaaaagc tcagaaaagga aaaggaagtt gagttcgtgg cgggtggacta cttgcaactt 960
ctgagaccgc cagtccgaaa gagttcaaga caggaggaag tggcagaggt ttcaagaaac 1020
ttaaaagccc ttgcaaagga acttcacatt cccggttatgg cacttgcgca gctctcccg 1080
gaggtggaaa agaggagtga taaaagaccc cagcttgagg acctcagaga atccggacag 1140
atagaacagg acgcagacct aatccttttc ctccacagac ccgagtacta caagaaaaag 1200
ccaaatcccg aagagcaggg tatagcggaa gtgataatag ccaagcaaag gcaaggaccc 1260
acggacattg tgaagctcgc atttattaag gagtacacta agtttgcaaa cctagaagcc 1320
cttcctgaac aacctcctga agaagaggaa ctttccgaaa ttattgaaac acaggaggat 1380
gaaggattcg aagatattga cttctgaaaa ttaagggtttt ataattttat cttggctatc 1440

cggggtagct caatcggcag agcgggtggc tg

1472

<210> 132

<211> 438

<212> PRT

<213> Aquifex aeolicus

<400> 132

Met Gln Phe Val Asp Lys Leu Pro Cys Asp Glu Ser Ala Glu Arg Ala
1 5 10 15

Val Leu Gly Ser Met Leu Glu Asp Pro Glu Asn Ile Pro Leu Val Leu
20 25 30

Glu Tyr Leu Lys Glu Glu Asp Phe Cys Ile Asp Glu His Lys Leu Leu
35 40 45

Phe Arg Val Leu Thr Asn Leu Trp Ser Glu Tyr Gly Asn Lys Leu Asp
50 55 60

Phe Val Leu Ile Lys Asp His Leu Glu Lys Lys Asn Leu Leu Gln Lys
65 70 75 80

Ile Pro Ile Asp Trp Leu Glu Glu Leu Tyr Glu Glu Ala Val Ser Pro
85 90 95

Asp Thr Leu Glu Glu Val Cys Lys Ile Val Lys Gln Arg Ser Ala Gln
100 105 110

Arg Ala Ile Ile Gln Leu Gly Ile Thr Ser Thr Gln Phe Tyr His Val
115 120 125

Lys Asp Val Ala Glu Glu Val Ile Glu Leu Ile Tyr Lys Phe Lys Ser
130 135 140

Ser Asp Arg Leu Val Thr Gly Leu Pro Ser Gly Phe Thr Glu Leu Asp
145 150 155 160

Leu Lys Thr Thr Gly Phe His Pro Gly Asp Leu Ile Ile Leu Ala Ala
165 170 175

Arg Pro Gly Met Gly Lys Thr Ala Phe Met Leu Ser Ile Ile Tyr Asn
180 185 190

Leu Ala Lys Asp Glu Gly Lys Pro Ser Ala Val Phe Ser Leu Glu Met
195 200 205

Ser Lys Glu Gln Leu Val Met Arg Leu Leu Ser Met Met Ser Glu Val
 210 215 220

Pro Leu Phe Lys Ile Arg Ser Gly Ser Ile Ser Asn Glu Asp Leu Lys
 225 230 235 240

Lys Leu Glu Ala Ser Ala Ile Glu Leu Ala Lys Tyr Asp Ile Tyr Leu
 245 250 255

Asp Asp Thr Pro Ala Leu Thr Thr Thr Asp Leu Arg Ile Arg Ala Arg
 260 265 270

Lys Leu Arg Lys Glu Lys Glu Val Glu Phe Val Ala Val Asp Tyr Leu
 275 280 285

Gln Leu Leu Arg Pro Pro Val Arg Lys Ser Ser Arg Gln Glu Glu Val
 290 295 300

Ala Glu Val Ser Arg Asn Leu Lys Ala Leu Ala Lys Glu Leu His Ile
 305 310 315 320

Pro Val Met Ala Leu Ala Gln Leu Ser Arg Glu Val Glu Lys Arg Ser
 325 330 335

Asp Lys Arg Pro Gln Leu Ala Asp Leu Arg Glu Ser Gly Gln Ile Glu
 340 345 350

Gln Asp Ala Asp Leu Ile Leu Phe Leu His Arg Pro Glu Tyr Tyr Lys
 355 360 365

Lys Lys Pro Asn Pro Glu Glu Gln Gly Ile Ala Glu Val Ile Ile Ala
 370 375 380

Lys Gln Arg Gln Gly Pro Thr Asp Ile Val Lys Leu Ala Phe Ile Lys
 385 390 395 400

Glu Tyr Thr Lys Phe Ala Asn Leu Glu Ala Leu Pro Glu Gln Pro Pro
 405 410 415

Glu Glu Glu Glu Leu Ser Glu Ile Ile Glu Thr Gln Glu Asp Glu Gly
 420 425 430

Phe Glu Asp Ile Asp Phe
 435

<210> 133

<211> 1526

<212> DNA

<213> Aquifex aeolicus

<400> 133

atgtcctcgg acatagacga acttagacgg gaaatagata tagtagacgt catttccgaa 60
tacttaaaact tagagaaggt aggttccaat tacagaacga actgtccctt tcaccctgac 120
gatacaccct ccttttacgt gtctccaagt aaacaaatat tcaagtgttt cggttgccggg 180
gtagggggag acgcgataaa gttcggtttcc ctttacgagg acatctccta ttttgaagcc 240
gcccttgaac tcgcaaaacg ctacggaaaag aaattagacc ttgaaaagat atcaaaagac 300
gaaaaggtat acgtggctct tgacaggggt tgtgatttct acagggaaaag ctttctcaaa 360
aacagagagg caagttagta cgtaaagagt aggggaatag accctaaagt agcgaggaag 420
tttgatcttg ggtacgcacc ttccagtga gactcgtaa aagtcttaaa agagaacgat 480
cttttagagg cttaccttga aactaaaaac ctctttctc ctacgaaggg tgtttacagg 540
gatctctttc ttcggtgtgt cgtgatcccg ataaaggatc cgaggggaag agttataggt 600
ttcggtggaa ggaggatagt agaggacaaa tctccaagt acataaactc tccagacagc 660
agggtattta aaaaggggga gaacttattc ggtctttacg aggcaaagga gtatataaag 720
gaagaaggat ttgcgatact tgtggaaggg tactttgacc ttttgagact tttttccgag 780
ggaataagga acgttggtgc acccctcggg acagccctga cccaaaatca ggcaaactc 840
ctttccaagt tcacaaaaaa ggtctacatc ctttacgacg gagatgatgc gggaagaaaag 900
gctatgaaaa gtgccattcc cctactctc agtgcaggag tggaagtta tcccgttac 960
ctccccgaag gatacgatcc cgacgagttt ataaaggaat tcgggaaaga ggaattaaga 1020
agactgataa acagctcagg ggagctcttt gaaacgctca taaaaaccgc aagggaaaac 1080
ttagaggaga aaacgcgtga gttcaggtat tatctgggct ttatttccga tggagtaagg 1140
cgctttgctc tggcttcgga gtttcacacc aagtacaaaag ttcctatgga aattttatta 1200
atgaaaattg aaaaaaattc tcaagaaaaa gaaattaaac tctcctttaa ggaaaaaatc 1260
ttcctgaaag gactgataga attaaaacca aaaatagacc ttgaagtcct gaacttaagt 1320
cctgagttaa aggaactcgc agttaacgcc ttaaaggag aggagcattt acttccaaaa 1380
gaagttctcg agtaccaggt ggataacttg gagaaacttt ttaacaacat ccttagggat 1440
ttacaaaaat ctgggaaaaa gaggaagaaa agagggttga aaaatgtaaa tacttaatta 1500
actttaataa attttttagag ttagga 1526

<210> 134

<211> 498

<212> PRT

<213> Aquifex aeolicus

<400> 134

Met Ser Ser Asp Ile Asp Glu Leu Arg Arg Glu Ile Asp Ile Val Asp
1 5 10 15

Val Ile Ser Glu Tyr Leu Asn Leu Glu Lys Val Gly Ser Asn Tyr Arg
20 25 30

Thr Asn Cys Pro Phe His Pro Asp Asp Thr Pro Ser Phe Tyr Val Ser
35 40 45

Pro Ser Lys Gln Ile Phe Lys Cys Phe Gly Cys Gly Val Gly Gly Asp

50	55	60
Ala Ile Lys Phe Val Ser Leu Tyr Glu Asp Ile Ser Tyr Phe Glu Ala		
65	70	75 80
Ala Leu Glu Leu Ala Lys Arg Tyr Gly Lys Lys Leu Asp Leu Glu Lys		
	85	90 95
Ile Ser Lys Asp Glu Lys Val Tyr Val Ala Leu Asp Arg Val Cys Asp		
	100	105 110
Phe Tyr Arg Glu Ser Leu Leu Lys Asn Arg Glu Ala Ser Glu Tyr Val		
	115	120 125
Lys Ser Arg Gly Ile Asp Pro Lys Val Ala Arg Lys Phe Asp Leu Gly		
	130	135 140
Tyr Ala Pro Ser Ser Glu Ala Leu Val Lys Val Leu Lys Glu Asn Asp		
	145	150 155 160
Leu Leu Glu Ala Tyr Leu Glu Thr Lys Asn Leu Leu Ser Pro Thr Lys		
	165	170 175
Gly Val Tyr Arg Asp Leu Phe Leu Arg Arg Val Val Ile Pro Ile Lys		
	180	185 190
Asp Pro Arg Gly Arg Val Ile Gly Phe Gly Gly Arg Arg Ile Val Glu		
	195	200 205
Asp Lys Ser Pro Lys Tyr Ile Asn Ser Pro Asp Ser Arg Val Phe Lys		
	210	215 220
Lys Gly Glu Asn Leu Phe Gly Leu Tyr Glu Ala Lys Glu Tyr Ile Lys		
	225	230 235 240
Glu Glu Gly Phe Ala Ile Leu Val Glu Gly Tyr Phe Asp Leu Leu Arg		
	245	250 255
Leu Phe Ser Glu Gly Ile Arg Asn Val Val Ala Pro Leu Gly Thr Ala		
	260	265 270
Leu Thr Gln Asn Gln Ala Asn Leu Leu Ser Lys Phe Thr Lys Lys Val		
	275	280 285
Tyr Ile Leu Tyr Asp Gly Asp Asp Ala Gly Arg Lys Ala Met Lys Ser		
	290	295 300
Ala Ile Pro Leu Leu Leu Ser Ala Gly Val Glu Val Tyr Pro Val Tyr		

305 310 315 320
 Leu Pro Glu Gly Tyr Asp Pro Asp Glu Phe Ile Lys Glu Phe Gly Lys
 325 330 335
 Glu Glu Leu Arg Arg Leu Ile Asn Ser Ser Gly Glu Leu Phe Glu Thr
 340 345 350
 Leu Ile Lys Thr Ala Arg Glu Asn Leu Glu Glu Lys Thr Arg Glu Phe
 355 360 365
 Arg Tyr Tyr Leu Gly Phe Ile Ser Asp Gly Val Arg Arg Phe Ala Leu
 370 375 380
 Ala Ser Glu Phe His Thr Lys Tyr Lys Val Pro Met Glu Ile Leu Leu
 385 390 395 400
 Met Lys Ile Glu Lys Asn Ser Gln Glu Lys Glu Ile Lys Leu Ser Phe
 405 410 415
 Lys Glu Lys Ile Phe Leu Lys Gly Leu Ile Glu Leu Lys Pro Lys Ile
 420 425 430
 Asp Leu Glu Val Leu Asn Leu Ser Pro Glu Leu Lys Glu Leu Ala Val
 435 440 445
 Asn Ala Leu Asn Gly Glu Glu His Leu Leu Pro Lys Glu Val Leu Glu
 450 455 460
 Tyr Gln Val Asp Asn Leu Glu Lys Leu Phe Asn Asn Ile Leu Arg Asp
 465 470 475 480
 Leu Gln Lys Ser Gly Lys Lys Arg Lys Lys Arg Gly Leu Lys Asn Val
 485 490 495

Asn Thr

<210> 135

<211> 705

<212> DNA

<213> Aquifex aeolicus

<400> 135

atgcaagata ccgctacctg cagtatttgt caggggacgg gattcgtaaa gaccgaagac 60
 aacaaggtaa ggctctgcga atgcaggttc aagaaaaggg atgtaaacag ggaactaaac 120
 atcccaaaga ggtactggaa cgccaactta gacacttacc accccaagaa cgtatccag 180

aacagggcac ttttgacgat aagggtcttc gtccacaact tcaatcccga ggaagggaaa 240
 gggcttacct ttgtaggatc tcctggagtc ggcaaaactc accttgcggt tgcaacatta 300
 aaagcgattt atgagaagaa gggaatcaga ggatacttct tcgatacgaa ggatctaata 360
 ttcagggttaa aacacttaat ggacgagga aaggatacaa agttttttaa aactgtctta 420
 aactcaccgg ttttggttct cgacgacctc ggttctgaga ggctcagtga ctggcagagg 480
 gaactcatct cttacataat cacttacagg tataacaacc ttaagagcac gataataacc 540
 acgaattact cactccagag ggaagaagag agtagcgtga ggataagtgc ggatcttgca 600
 agcagactcg gagaaaacgt agtttcaaaa atttacgaga tgaacgagtt gctcgttata 660
 aagggttccg acctcaggaa gtctaaaaag ctatcaaccc catct 705

<210> 136

<211> 235

<212> PRT

<213> Aquifex aeolicus

<400> 136

Met Gln Asp Thr Ala Thr Cys Ser Ile Cys Gln Gly Thr Gly Phe Val
 1 5 10 15

Lys Thr Glu Asp Asn Lys Val Arg Leu Cys Glu Cys Arg Phe Lys Lys
 20 25 30

Arg Asp Val Asn Arg Glu Leu Asn Ile Pro Lys Arg Tyr Trp Asn Ala
 35 40 45

Asn Leu Asp Thr Tyr His Pro Lys Asn Val Ser Gln Asn Arg Ala Leu
 50 55 60

Leu Thr Ile Arg Val Phe Val His Asn Phe Asn Pro Glu Glu Gly Lys
 65 70 75 80

Gly Leu Thr Phe Val Gly Ser Pro Gly Val Gly Lys Thr His Leu Ala
 85 90 95

Val Ala Thr Leu Lys Ala Ile Tyr Glu Lys Lys Gly Ile Arg Gly Tyr
 100 105 110

Phe Phe Asp Thr Lys Asp Leu Ile Phe Arg Leu Lys His Leu Met Asp
 115 120 125

Glu Gly Lys Asp Thr Lys Phe Leu Lys Thr Val Leu Asn Ser Pro Val
 130 135 140

Leu Val Leu Asp Asp Leu Gly Ser Glu Arg Leu Ser Asp Trp Gln Arg
 145 150 155 160

Glu Leu Ile Ser Tyr Ile Ile Thr Tyr Arg Tyr Asn Asn Leu Lys Ser

165

170

175

Thr Ile Ile Thr Thr Asn Tyr Ser Leu Gln Arg Glu Glu Glu Ser Ser
180 185 190

Val Arg Ile Ser Ala Asp Leu Ala Ser Arg Leu Gly Glu Asn Val Val
195 200 205

Ser Lys Ile Tyr Glu Met Asn Glu Leu Leu Val Ile Lys Gly Ser Asp
210 215 220

Leu Arg Lys Ser Lys Lys Leu Ser Thr Pro Ser
225 230 235

<210> 137

<211> 4101

<212> DNA

<213> *Thermatoga maritima*

<400> 137

atgaaaaaga ttgaaaattt gaagtggaaa aatgtctcgt ttaaaagcct ggaaatagat 60
cccgatgcag gtgtggttct cgtttccgtg gaaaaattct ccgaagagat agaagacctt 120
gtgcgtttac tggagaagaa gacgcggtt cgagtcacgc tgaacgggtgt tcaaaaaagt 180
aacggggatc taaggggaaa gatactttcc cttctcaacg gtaatgtgcc ttacataaaa 240
gatgttgttt tcgaaggaaa caggctgatt ctgaaagtgc ttggagattt cgcgcgggac 300
aggatcgcct ccaaactcag aagcacgaaa aaacagctcg atgaactgct gcctcccgga 360
acagagatca tgctggaggt tgtggagcct ccggaagatc ttttgaaaaa ggaagtacca 420
caaccagaaa agagagaaga accaaaggtg gaagaattga agatcgagga tgaaaaccac 480
atctttggac agaaaccag aaagatcgct ttcaccccct caaaaatctt tgagtacaac 540
aaaaagacat cgggtgaagg caagatcttc aaaatagaga agatcgaggg gaaaagaacg 600
gtccttctga tttacctgac agacggagaa gattctctga tctgcaaagt cttcaacgac 660
gttgaaaagg tcgaagggaa agtatcgggt ggagacgtga tcgttgccac aggagacctc 720
cttctcgaaa acggggagcc caccctttac gtgaagggaa tcacaaaact tcccgaagcg 780
aaaaggatgg acaaactctc gggttaagagg gtggagctcc acgcccatac caagttcagc 840
gatcaggacg caataacaga tgtgaacgaa tatgtgaaac gagccaagga atggggcctt 900
cccgcgatag ccctcacgga tcatgggaac gttcaggcca taccttactt ctacgacgcg 960
gcgaaagaag ctggaataaa gccattttc ggtatcgaag cgtatctggt gagtacgtg 1020
gagcccgta taaggaatct ctccgacgat tcgacgtttg gagatgccac gttcgtcgtc 1080
ctcgacttcg agacgacggg tctcgaccgg cagggtggatg agatcatcga gataggagcg 1140
gtgaagatac aggttgacca gatagtggac gaggaccaca ctctcataaa gccttcagg 1200
gagatctcaa gaaaaagttc ggagatcacc ggaatcactc aagagatgct ggaaaacaag 1260
agaagcatcg aggaagtctt gccggagttc ctcgggtttc tggaagattc catcatcgta 1320
gcacacaacg ccaacttcga ctacagattt ctgaggctgt ggatcaaaaa agtgatggga 1380
ttggactggg aaagacccta catagatacg ctcgcccctc caaagtccct tctcaaactg 1440
agaagctact ctctggattc cgttgtggaa aagctcggat tgggtccctt ccggcaccac 1500
agggcccttg atgacgcgag ggtcaccgct cagggtttcc tcaggttcgt tgagatgatg 1560
aagaagatcg gtatcacgaa gctttcagaa atggagaagt tgaaggatac gatagactac 1620

```

accgcgttga aacccttcca ctgcacgato ctcgttcaga acaaaaaggg attgaaaaac 1680
ctatacaaac tggtttctga ttcctatata aagtacttct acggtgttcc gaggatcctc 1740
aaaagtgagc tcatcgagaa cagagaagga ctgctcgtgg gtagcgcgtg tatctccggt 1800
gagctcggac gtgccgccct cgaaggagcg agtgattcag aactcgaaga gatcgcgaag 1860
ttctacgact acatagaagt catgccgctc gacgttatag ccgaagatga agaagaccta 1920
gacagagaaa gactgaaaga agtgtaccga aaactctaca gaatagcgaa aaaattgaac 1980
aagttcgtcg tcatgaccgg tgatgttcat ttctctgata ccgaagatgc caggggcaga 2040
gctgcacttc tggcacctca gggaaacaga aacttcgaga atcagcccgc actctacctc 2100
agaacgaccg aagaaatgct cgagaaggcg atagagatat tcgaagatga agagatcgcg 2160
agggaaagtc tgatagagaa tccaacaga atagccgata tgatcgagga agtgcagccg 2220
ctcgagaaaa aacttcaccc gccgatcata gagaacgccg atgaaatagt gagaaacctc 2280
accatgaagc gggcgtagca gatctacggt gatccgcttc ccgaaatcgt ccagaagcgt 2340
gtggaaaagg aactgaacgc catcataaat catggatacg ccgttctcta tctcatcgct 2400
caggagctcg ttcagaaatc tatgagcgat gggtacgtgg ttggatccag aggatccgtc 2460
gggtcttcac tcgtggccaa tctctcggga ataacagagg tgaatcccct accaccacat 2520
tacaggtgtc cagagtgcaa atactttgaa gttgtcgaag acgacagata cggagcgggt 2580
tacgaccttc ccaacaagaa ctgtccaaga tgtggggctc ctctcagaaa agacggccac 2640
ggcataccgt ttgaaacggt catgggggttc gaggtgacaa aggtccccga catagatctc 2700
aacttctcag gagagtatca ggaacgtgct catcgttttg tggaagaact cttcggtaaa 2760
gaccacgtct atagggcggg aaccataaac accatcgcgg aaagaagtgc ggtgggttac 2820
gtgagaagct acgaagagaa aaccggaaag aagctcagaa aggcggaaat ggaaagactc 2880
gtttccatga tcacgggagt gaagagaacg acgggtcagc acccaggggg gctcatgata 2940
ataccgaaag acaagaagt ctacgatttc actcccatac agtatccagc caacgataga 3000
aacgcagggtg tgttcaccac gcacttcgca tacgagacga tccatgatga cctggtgaag 3060
atagatgcgc tcggccacga tgatcccact ttcatacaga tgcctcaagga cctcaccgga 3120
atcgatccca tgacgattcc catggatgac cccgatacgc tcgccatatt cagttctgtg 3180
aagcctcttg gtgtggatcc cggtgagctg gaaagcgatg tgggaacgta cggaattccg 3240
gagttcggaa ccgagtttgt gaggggaatg ctcgttgaaa cgagaccaa gagtttcgcc 3300
gagcttgtga gaatctcagg actgtcacac ggtacggacg tctggttgaa caacgcacgt 3360
gattggataa acctcggcta cgccaagctc tccgaggtta tctcgtgtag ggacgacatc 3420
atgaacttcc tcatacaca aggaatggaa ccgtcacttg ccttcaagat catggaaaac 3480
gtcaggaagg gaaagggtat cacagaagag atggagagcg agatgagaag gctgaagggt 3540
ccagaatggt tcatcgaatc ctgtaaaagg atcaaatatc tcttcccgaa agctcacgct 3600
gtggcttacg tgagtatggc cttcagaatt gcttacttca aggttcacta tctctctcag 3660
ttttacgcgg cgtacttcac gataaaagggt gatcagttcg atccggttct cgtactcagg 3720
ggaaaagaag ccataaagag gcgcttgaga gaactcaaag cgatgcctgc caaagacgcc 3780
cagaagaaaa acgaagtgag tgttctggag gttgccctgg aaatgatact gagagggttt 3840
tccttcttac cgcccgacat cttcaaatec gacgcgaaga aatttctgat agaaggaaac 3900
tcgctgagaa ttccgttcaa caaacttcca ggactgggtg acagcggttg cgagtcgata 3960
atcagagcca gggaagaaaa gccgttcaat tcgggtggaag atctcatgaa gaggaccaag 4020
gtcaacaaaa atcacataga gctgatgaaa agcctgggtg ttctcgggga ccttcagag 4080
acggaacagt tcacgctttt c 4101

```

<210> 138

<211> 1367

<212> PRT

<213> *Thermatoga maritima*

<400> 138

Met Lys Lys Ile Glu Asn Leu Lys Trp Lys Asn Val Ser Phe Lys Ser
1 5 10 15

Leu Glu Ile Asp Pro Asp Ala Gly Val Val Leu Val Ser Val Glu Lys
20 25 30

Phe Ser Glu Glu Ile Glu Asp Leu Val Arg Leu Leu Glu Lys Lys Thr
35 40 45

Arg Phe Arg Val Ile Val Asn Gly Val Gln Lys Ser Asn Gly Asp Leu
50 55 60

Arg Gly Lys Ile Leu Ser Leu Leu Asn Gly Asn Val Pro Tyr Ile Lys
65 70 75 80

Asp Val Val Phe Glu Gly Asn Arg Leu Ile Leu Lys Val Leu Gly Asp
85 90 95

Phe Ala Arg Asp Arg Ile Ala Ser Lys Leu Arg Ser Thr Lys Lys Gln
100 105 110

Leu Asp Glu Leu Leu Pro Pro Gly Thr Glu Ile Met Leu Glu Val Val
115 120 125

Glu Pro Pro Glu Asp Leu Leu Lys Lys Glu Val Pro Gln Pro Glu Lys
130 135 140

Arg Glu Glu Pro Lys Gly Glu Glu Leu Lys Ile Glu Asp Glu Asn His
145 150 155 160

Ile Phe Gly Gln Lys Pro Arg Lys Ile Val Phe Thr Pro Ser Lys Ile
165 170 175

Phe Glu Tyr Asn Lys Lys Thr Ser Val Lys Gly Lys Ile Phe Lys Ile
180 185 190

Glu Lys Ile Glu Gly Lys Arg Thr Val Leu Leu Ile Tyr Leu Thr Asp
195 200 205

Gly Glu Asp Ser Leu Ile Cys Lys Val Phe Asn Asp Val Glu Lys Val
210 215 220

Glu Gly Lys Val Ser Val Gly Asp Val Ile Val Ala Thr Gly Asp Leu
225 230 235 240

Leu Leu Glu Asn Gly Glu Pro Thr Leu Tyr Val Lys Gly Ile Thr Lys

001211 1964 112100

245	250	255
Leu Pro Glu Ala Lys Arg Met Asp Lys Ser Pro Val Lys Arg Val Glu		
260	265	270
Leu His Ala His Thr Lys Phe Ser Asp Gln Asp Ala Ile Thr Asp Val		
275	280	285
Asn Glu Tyr Val Lys Arg Ala Lys Glu Trp Gly Phe Pro Ala Ile Ala		
290	295	300
Leu Thr Asp His Gly Asn Val Gln Ala Ile Pro Tyr Phe Tyr Asp Ala		
305	310	315 320
Ala Lys Glu Ala Gly Ile Lys Pro Ile Phe Gly Ile Glu Ala Tyr Leu		
325	330	335
Val Ser Asp Val Glu Pro Val Ile Arg Asn Leu Ser Asp Asp Ser Thr		
340	345	350
Phe Gly Asp Ala Thr Phe Val Val Leu Asp Phe Glu Thr Thr Gly Leu		
355	360	365
Asp Pro Gln Val Asp Glu Ile Ile Glu Ile Gly Ala Val Lys Ile Gln		
370	375	380
Gly Gly Gln Ile Val Asp Glu Tyr His Thr Leu Ile Lys Pro Ser Arg		
385	390	395 400
Glu Ile Ser Arg Lys Ser Ser Glu Ile Thr Gly Ile Thr Gln Glu Met		
405	410	415
Leu Glu Asn Lys Arg Ser Ile Glu Glu Val Leu Pro Glu Phe Leu Gly		
420	425	430
Phe Leu Glu Asp Ser Ile Ile Val Ala His Asn Ala Asn Phe Asp Tyr		
435	440	445
Arg Phe Leu Arg Leu Trp Ile Lys Lys Val Met Gly Leu Asp Trp Glu		
450	455	460
Arg Pro Tyr Ile Asp Thr Leu Ala Leu Ala Lys Ser Leu Leu Lys Leu		
465	470	475 480
Arg Ser Tyr Ser Leu Asp Ser Val Val Glu Lys Leu Gly Leu Gly Pro		
485	490	495
Phe Arg His His Arg Ala Leu Asp Asp Ala Arg Val Thr Ala Gln Val		

500										505					510						
Phe	Leu	Arg	Phe	Val	Glu	Met	Met	Lys	Lys	Ile	Gly	Ile	Thr	Lys	Leu						
515					520					525											
Ser	Glu	Met	Glu	Lys	Leu	Lys	Asp	Thr	Ile	Asp	Tyr	Thr	Ala	Leu	Lys						
530					535					540											
Pro	Phe	His	Cys	Thr	Ile	Leu	Val	Gln	Asn	Lys	Lys	Gly	Leu	Lys	Asn						
545					550					555					560						
Leu	Tyr	Lys	Leu	Val	Ser	Asp	Ser	Tyr	Ile	Lys	Tyr	Phe	Tyr	Gly	Val						
565					570					575											
Pro	Arg	Ile	Leu	Lys	Ser	Glu	Leu	Ile	Glu	Asn	Arg	Glu	Gly	Leu	Leu						
580					585					590											
Val	Gly	Ser	Ala	Cys	Ile	Ser	Gly	Glu	Leu	Gly	Arg	Ala	Ala	Leu	Glu						
595					600					605											
Gly	Ala	Ser	Asp	Ser	Glu	Leu	Glu	Glu	Ile	Ala	Lys	Phe	Tyr	Asp	Tyr						
610					615					620											
Ile	Glu	Val	Met	Pro	Leu	Asp	Val	Ile	Ala	Glu	Asp	Glu	Glu	Asp	Leu						
625					630					635					640						
Asp	Arg	Glu	Arg	Leu	Lys	Glu	Val	Tyr	Arg	Lys	Leu	Tyr	Arg	Ile	Ala						
645					650					655											
Lys	Lys	Leu	Asn	Lys	Phe	Val	Val	Met	Thr	Gly	Asp	Val	His	Phe	Leu						
660					665					670											
Asp	Pro	Glu	Asp	Ala	Arg	Gly	Arg	Ala	Ala	Leu	Leu	Ala	Pro	Gln	Gly						
675					680					685											
Asn	Arg	Asn	Phe	Glu	Asn	Gln	Pro	Ala	Leu	Tyr	Leu	Arg	Thr	Thr	Glu						
690					695					700											
Glu	Met	Leu	Glu	Lys	Ala	Ile	Glu	Ile	Phe	Glu	Asp	Glu	Glu	Ile	Ala						
705					710					715					720						
Arg	Glu	Val	Val	Ile	Glu	Asn	Pro	Asn	Arg	Ile	Ala	Asp	Met	Ile	Glu						
725					730					735											
Glu	Val	Gln	Pro	Leu	Glu	Lys	Lys	Leu	His	Pro	Pro	Ile	Ile	Glu	Asn						
740					745					750											
Ala	Asp	Glu	Ile	Val	Arg	Asn	Leu	Thr	Met	Lys	Arg	Ala	Tyr	Glu	Ile						

00716964-112100

755		760		765
Tyr Gly Asp Pro Leu Pro Glu Ile Val Gln Lys Arg Val Glu Lys Glu				
770		775		780
Leu Asn Ala Ile Ile Asn His Gly Tyr Ala Val Leu Tyr Leu Ile Ala				
785		790		800
Gln Glu Leu Val Gln Lys Ser Met Ser Asp Gly Tyr Val Val Gly Ser				
	805		810	815
Arg Gly Ser Val Gly Ser Ser Leu Val Ala Asn Leu Leu Gly Ile Thr				
	820		825	830
Glu Val Asn Pro Leu Pro Pro His Tyr Arg Cys Pro Glu Cys Lys Tyr				
	835		840	845
Phe Glu Val Val Glu Asp Asp Arg Tyr Gly Ala Gly Tyr Asp Leu Pro				
	850		855	860
Asn Lys Asn Cys Pro Arg Cys Gly Ala Pro Leu Arg Lys Asp Gly His				
	865		870	880
Gly Ile Pro Phe Glu Thr Phe Met Gly Phe Glu Gly Asp Lys Val Pro				
		885	890	895
Asp Ile Asp Leu Asn Phe Ser Gly Glu Tyr Gln Glu Arg Ala His Arg				
	900		905	910
Phe Val Glu Glu Leu Phe Gly Lys Asp His Val Tyr Arg Ala Gly Thr				
	915		920	925
Ile Asn Thr Ile Ala Glu Arg Ser Ala Val Gly Tyr Val Arg Ser Tyr				
	930		935	940
Glu Glu Lys Thr Gly Lys Lys Leu Arg Lys Ala Glu Met Glu Arg Leu				
	945		950	955
Val Ser Met Ile Thr Gly Val Lys Arg Thr Thr Gly Gln His Pro Gly				
		965	970	975
Gly Leu Met Ile Ile Pro Lys Asp Lys Glu Val Tyr Asp Phe Thr Pro				
	980		985	990
Ile Gln Tyr Pro Ala Asn Asp Arg Asn Ala Gly Val Phe Thr Thr His				
	995		1000	1005
Phe Ala Tyr Glu Thr Ile His Asp Asp Leu Val Lys Ile Asp Ala Leu				

00746964-12100

1010	1015	1020
Gly His Asp Asp Pro Thr Phe Ile Lys Met Leu Lys Asp Leu Thr Gly		
1025	1030	1035 1040
Ile Asp Pro Met Thr Ile Pro Met Asp Asp Pro Asp Thr Leu Ala Ile		
1045	1050	1055
Phe Ser Ser Val Lys Pro Leu Gly Val Asp Pro Val Glu Leu Glu Ser		
1060	1065	1070
Asp Val Gly Thr Tyr Gly Ile Pro Glu Phe Gly Thr Glu Phe Val Arg		
1075	1080	1085
Gly Met Leu Val Glu Thr Arg Pro Lys Ser Phe Ala Glu Leu Val Arg		
1090	1095	1100
Ile Ser Gly Leu Ser His Gly Thr Asp Val Trp Leu Asn Asn Ala Arg		
1105	1110	1115 1120
Asp Trp Ile Asn Leu Gly Tyr Ala Lys Leu Ser Glu Val Ile Ser Cys		
1125	1130	1135
Arg Asp Asp Ile Met Asn Phe Leu Ile His Lys Gly Met Glu Pro Ser		
1140	1145	1150
Leu Ala Phe Lys Ile Met Glu Asn Val Arg Lys Gly Lys Gly Ile Thr		
1155	1160	1165
Glu Glu Met Glu Ser Glu Met Arg Arg Leu Lys Val Pro Glu Trp Phe		
1170	1175	1180
Ile Glu Ser Cys Lys Arg Ile Lys Tyr Leu Phe Pro Lys Ala His Ala		
1185	1190	1195 1200
Val Ala Tyr Val Ser Met Ala Phe Arg Ile Ala Tyr Phe Lys Val His		
1205	1210	1215
Tyr Pro Leu Gln Phe Tyr Ala Ala Tyr Phe Thr Ile Lys Gly Asp Gln		
1220	1225	1230
Phe Asp Pro Val Leu Val Leu Arg Gly Lys Glu Ala Ile Lys Arg Arg		
1235	1240	1245
Leu Arg Glu Leu Lys Ala Met Pro Ala Lys Asp Ala Gln Lys Lys Asn		
1250	1255	1260
Glu Val Ser Val Leu Glu Val Ala Leu Glu Met Ile Leu Arg Gly Phe		

1265 1270 1275 1280
 Ser Phe Leu Pro Pro Asp Ile Phe Lys Ser Asp Ala Lys Lys Phe Leu
 1285 1290 1295
 Ile Glu Gly Asn Ser Leu Arg Ile Pro Phe Asn Lys Leu Pro Gly Leu
 1300 1305 1310
 Gly Asp Ser Val Ala Glu Ser Ile Ile Arg Ala Arg Glu Glu Lys Pro
 1315 1320 1325
 Phe Thr Ser Val Glu Asp Leu Met Lys Arg Thr Lys Val Asn Lys Asn
 1330 1335 1340
 His Ile Glu Leu Met Lys Ser Leu Gly Val Leu Gly Asp Leu Pro Glu
 1345 1350 1355 1360
 Thr Glu Gln Phe Thr Leu Phe
 1365

<210> 139
 <211> 567
 <212> DNA
 <213> *Thermatoga maritima*

<400> 139
 gtgctcgcca tgatatggaa cgacaccggt ttttgcgtcg tagacacaga aaccacggga 60
 accgatccct ttgccggaga ccggatagtt gaaatagccg ctgttcctgt cttcaagggg 120
 aagatctaca gaaacaaagc gtttcactct ctcgtgaatc ccagaataag aatccctgcg 180
 ctgattcaga aagttcacgg tatcagcaac atggacatcg tggaagcgcc agacatggac 240
 acagtttacg atcttttcag ggattacgtg aagggaacgg tgctcgtgtt tcacaacgcc 300
 aacttcgacc tcactttttct ggatatgatg gcaaaggaaa cgggaaactt tccaataacg 360
 aatccctaca tcgacacact cgatctttca gaagagatct ttggaaggcc tcattctctc 420
 aaatggctct ccgaaagact tggaataaaa accacgatac ggcaccgtgc tcttccagat 480
 gccctgggtga ccgcaagagt ttttgtgaag cttgttgaat ttcttgggtga aaacaggggtc 540
 aacgaattca tacgtggaaa acggggg 567

<210> 140
 <211> 189
 <212> PRT
 <213> *Thermatoga maritima*

<400> 140
 Met Leu Ala Met Ile Trp Asn Asp Thr Val Phe Cys Val Val Asp Thr
 1 5 10 15

Glu Thr Thr Gly Thr Asp Pro Phe Ala Gly Asp Arg Ile Val Glu Ile
20 25 30

Ala Ala Val Pro Val Phe Lys Gly Lys Ile Tyr Arg Asn Lys Ala Phe
35 40 45

His Ser Leu Val Asn Pro Arg Ile Arg Ile Pro Ala Leu Ile Gln Lys
50 55 60

Val His Gly Ile Ser Asn Met Asp Ile Val Glu Ala Pro Asp Met Asp
65 70 75 80

Thr Val Tyr Asp Leu Phe Arg Asp Tyr Val Lys Gly Thr Val Leu Val
85 90 95

Phe His Asn Ala Asn Phe Asp Leu Thr Phe Leu Asp Met Met Ala Lys
100 105 110

Glu Thr Gly Asn Phe Pro Ile Thr Asn Pro Tyr Ile Asp Thr Leu Asp
115 120 125

Leu Ser Glu Glu Ile Phe Gly Arg Pro His Ser Leu Lys Trp Leu Ser
130 135 140

Glu Arg Leu Gly Ile Lys Thr Thr Ile Arg His Arg Ala Leu Pro Asp
145 150 155 160

Ala Leu Val Thr Ala Arg Val Phe Val Lys Leu Val Glu Phe Leu Gly
165 170 175

Glu Asn Arg Val Asn Glu Phe Ile Arg Gly Lys Arg Gly
180 185

<210> 141

<211> 1434

<212> DNA

<213> *Thermatoga maritima*

<400> 141

gtggaagttc tttacaggaa gtacaggcca aagacttttt ctgaggttgt caatcaggat 60
catgtgaaga aggcaataat cgggtgctatt cagaagaaca gcgtggccca cggatacata 120
ttcgccggtc cgaggggaac ggggaagact actcttgcca gaattctcgc aaaatccctg 180
aactgtgaga acagaaaggg agttgaaccc tgcaattcct gcagagcctg cagagagata 240
gacgaggga ccttcatgga cgtgatagag ctcgacgcgg cctccaacag aggaatagac 300
gagatcagaa gaatcagaga cgccgttga tacaggccga tggaaggtaa atacaaagtc 360
tacataatag acgaagtcca catgctcacg aaagaagcct tcaacgcgct cctcaaaaca 420
ctcgaagaac ctccttccca cgtcgtgttc gtgctggcaa cgacaaacct tgagaaggtt 480

```

cctccacga ttatctcgag atgtcaggtt ttcgagttca gaaacattcc cgacgagctc 540
atcgaaaaga ggctccagga agttgctggag gctgaaggaa tagagataga cagggaagct 600
ctgagcttca tcgcaaaaag agcctctgga ggcttgagag acgcgctcac catgctcgag 660
cagggtgtgga agttctcggg aggaaagata gatctcgaga cggtagacag ggcgctcggg 720
ttgataccga tacaggttgt tcgctgattac gtgaacgcta tcttttctgg tgatgtgaaa 780
agggctcttca ccgttctcga cgacgtctat tacagcggga aggactacga ggtgctcatt 840
caggaagcag tcgaggatct ggtcgaagac ctggaaaggg agagaggggt ttaccaggtt 900
tcagcgaacg atatagttca ggtttcgaga caacttctga atcttctgag agagataaaag 960
ttcgccgaag aaaaacgact cgtctgtaaa gtgggttcgg cttacatagc gacgaggttc 1020
tccaccacaa acgttcagga aaacgatgtc agagaaaaaa acgataattc aaatgtacag 1080
cagaaagaag agaagaaaga aacggtgaag gcaaaagaag aaaaacagga agacagcgag 1140
ttcgagaaac gcttcaaaga actcatggaa gaactgaaag aaaagggcga tctctctatc 1200
tttgctcgtc tcagcctctc agaggtgcag tttgacggag aaaaggtgat tatttctttt 1260
gattcatcga aagctatgca ttacgagttg atgaagaaaa aactgcctga gctggaaaac 1320
atthtttcta gaaaactcgg gaaaaaagta gaagttgaac ttcgactgat gggaaaagaa 1380
gaaacaatcg agaaggtttc tcagaagatc ctgagattgt ttgaacagga ggga 1434

```

<210> 142

<211> 478

<212> PRT

<213> *Thermatoga maritima*

<400> 142

```

Met Glu Val Leu Tyr Arg Lys Tyr Arg Pro Lys Thr Phe Ser Glu Val
  1                      5                      10                      15

```

```

Val Asn Gln Asp His Val Lys Lys Ala Ile Ile Gly Ala Ile Gln Lys
          20                      25                      30

```

```

Asn Ser Val Ala His Gly Tyr Ile Phe Ala Gly Pro Arg Gly Thr Gly
          35                      40                      45

```

```

Lys Thr Thr Leu Ala Arg Ile Leu Ala Lys Ser Leu Asn Cys Glu Asn
          50                      55                      60

```

```

Arg Lys Gly Val Glu Pro Cys Asn Ser Cys Arg Ala Cys Arg Glu Ile
          65                      70                      75                      80

```

```

Asp Glu Gly Thr Phe Met Asp Val Ile Glu Leu Asp Ala Ala Ser Asn
          85                      90                      95

```

```

Arg Gly Ile Asp Glu Ile Arg Arg Ile Arg Asp Ala Val Gly Tyr Arg
          100                      105                      110

```

```

Pro Met Glu Gly Lys Tyr Lys Val Tyr Ile Ile Asp Glu Val His Met
          115                      120                      125

```

Leu Thr Lys Glu Ala Phe Asn Ala Leu Leu Lys Thr Leu Glu Glu Pro
 130 135 140
 Pro Ser His Val Val Phe Val Leu Ala Thr Thr Asn Leu Glu Lys Val
 145 150 155 160
 Pro Pro Thr Ile Ile Ser Arg Cys Gln Val Phe Glu Phe Arg Asn Ile
 165 170 175
 Pro Asp Glu Leu Ile Glu Lys Arg Leu Gln Glu Val Ala Glu Ala Glu
 180 185 190
 Gly Ile Glu Ile Asp Arg Glu Ala Leu Ser Phe Ile Ala Lys Arg Ala
 195 200 205
 Ser Gly Gly Leu Arg Asp Ala Leu Thr Met Leu Glu Gln Val Trp Lys
 210 215 220
 Phe Ser Glu Gly Lys Ile Asp Leu Glu Thr Val His Arg Ala Leu Gly
 225 230 235 240
 Leu Ile Pro Ile Gln Val Val Arg Asp Tyr Val Asn Ala Ile Phe Ser
 245 250 255
 Gly Asp Val Lys Arg Val Phe Thr Val Leu Asp Asp Val Tyr Tyr Ser
 260 265 270
 Gly Lys Asp Tyr Glu Val Leu Ile Gln Glu Ala Val Glu Asp Leu Val
 275 280 285
 Glu Asp Leu Glu Arg Glu Arg Gly Val Tyr Gln Val Ser Ala Asn Asp
 290 295 300
 Ile Val Gln Val Ser Arg Gln Leu Leu Asn Leu Leu Arg Glu Ile Lys
 305 310 315 320
 Phe Ala Glu Glu Lys Arg Leu Val Cys Lys Val Gly Ser Ala Tyr Ile
 325 330 335
 Ala Thr Arg Phe Ser Thr Thr Asn Val Gln Glu Asn Asp Val Arg Glu
 340 345 350
 Lys Asn Asp Asn Ser Asn Val Gln Gln Lys Glu Glu Lys Lys Glu Thr
 355 360 365
 Val Lys Ala Lys Glu Glu Lys Gln Glu Asp Ser Glu Phe Glu Lys Arg
 370 375 380

<212> PRT

<213> *Thermatoga maritima*

<400> 144

Met Lys Val Thr Val Thr Thr Leu Glu Leu Lys Asp Lys Ile Thr Ile
1 5 10 15

Ala Ser Lys Ala Leu Ala Lys Lys Ser Val Lys Pro Ile Leu Ala Gly
20 25 30

Phe Leu Phe Glu Val Lys Asp Gly Asn Phe Tyr Ile Cys Ala Thr Asp
35 40 45

Leu Glu Thr Gly Val Lys Ala Thr Val Asn Ala Ala Glu Ile Ser Gly
50 55 60

Glu Ala Arg Phe Val Val Pro Gly Asp Val Ile Gln Lys Met Val Lys
65 70 75 80

Val Leu Pro Asp Glu Ile Thr Glu Leu Ser Leu Glu Gly Asp Ala Leu
85 90 95

Val Ile Ser Ser Gly Ser Thr Val Phe Arg Ile Thr Thr Met Pro Ala
100 105 110

Asp Glu Phe Pro Glu Ile Thr Pro Ala Glu Ser Gly Ile Thr Phe Glu
115 120 125

Val Asp Thr Ser Leu Leu Glu Glu Met Val Glu Lys Val Ile Phe Ala
130 135 140

Ala Ala Lys Asp Glu Phe Met Arg Asn Leu Asn Gly Val Phe Trp Glu
145 150 155 160

Leu His Lys Asn Leu Leu Arg Leu Val Ala Ser Asp Gly Phe Arg Leu
165 170 175

Ala Leu Ala Glu Glu Gln Ile Glu Asn Glu Glu Glu Ala Ser Phe Leu
180 185 190

Leu Ser Leu Lys Ser Met Lys Glu Val Gln Asn Val Leu Asp Asn Thr
195 200 205

Thr Glu Pro Thr Ile Thr Val Arg Tyr Asp Gly Arg Arg Val Ser Leu
210 215 220

Ser Thr Asn Asp Val Glu Thr Val Met Arg Val Val Asp Ala Glu Phe
225 230 235 240

Pro Asp Tyr Lys Arg Val Ile Pro Glu Thr Phe Lys Thr Lys Val Val
245 250 255

Val Ser Arg Lys Glu Leu Arg Glu Ser Leu Lys Arg Val Met Val Ile
260 265 270

Ala Ser Lys Gly Ser Glu Ser Val Lys Phe Glu Ile Glu Glu Asn Val
275 280 285

Met Arg Leu Val Ser Lys Ser Pro Asp Tyr Gly Glu Val Val Asp Glu
290 295 300

Val Glu Val Gln Lys Glu Gly Glu Asp Leu Val Ile Ala Phe Asn Pro
305 310 315 320

Lys Phe Ile Glu Asp Val Leu Lys His Ile Glu Thr Glu Glu Ile Glu
325 330 335

Met Asn Phe Val Asp Ser Thr Ser Pro Cys Gln Ile Asn Pro Leu Asp
340 345 350

Ile Ser Gly Tyr Leu Tyr Ile Val Met Pro Ile Arg Leu Ala
355 360 365

<210> 145

<211> 972

<212> DNA

<213> *Thermatoga maritima*

<400> 145

```
atgccagtca cgtttctcac aggtactgca gaaactcaga aggaagaatt gataaagaaa 60
ctcctgaagg atggtaacgt ggagtacata aggatccatc cggaggatcc cgacaagatc 120
gatttcataa ggtctttact caggacaaag acgatctttt ccaacaagac gatcattgac 180
atcgtcaatt tcgatgagtg gaaagcacag gagcagaagc gtctcgttga acttttgaaa 240
aacgtaccgg aagacgttca tatcttcac cgttctcaaa aaacaggtgg aaagggagta 300
gcgctggagc ttccgaagcc atgggaaacg gacaagtggc ttgagtggat agaaaagcgc 360
ttcagggaga atggtttgct catcgataaa gatgcccttc agctgttttt ctccaagggt 420
ggaacgaacg acctgatcat agaaagggag attgaaaaac tgaaagctta ttccgaggac 480
agaaagataa cggtagaaga cgtggaagag gtcgttttta cctatcagac tccgggatac 540
gatgattttt gctttgctgt ttccgaagga aaaaggaagc tcgctcactc tcttctgtcg 600
cagctgtgga aaaccacaga gtccgtggtg attgccactg tccttgcgaa tcacttcttg 660
gatctcttca aaatcctcgt tcttgtagaca aagaaaagat actacacctg gcctgatgtg 720
tccaggggtg ccaaagagct gggaattccc gttcctcgtg tggtcgtttt cctcggtttc 780
tcctttaaga cctggaaatt caagtgatg aaccacctcc tctactacga tgtgaagaag 840
gttagaaaga tactgagga tctctacgat ctggacagag ccgtgaaaag cgaagaagat 900
ccaaaaccgt tcttcacga gttcatagaa gaggtggcac tggatgtata ttctcttcag 960
```


<210> 146

<211> 324

<212> PRT

<213> *Thermatoga maritima*

<400> 146

Met Pro Val Thr Phe Leu Thr Gly Thr Ala Glu Thr Gln Lys Glu Glu
 1 5 10 15

Leu Ile Lys Lys Leu Leu Lys Asp Gly Asn Val Glu Tyr Ile Arg Ile
 20 25 30

His Pro Glu Asp Pro Asp Lys Ile Asp Phe Ile Arg Ser Leu Leu Arg
 35 40 45

Thr Lys Thr Ile Phe Ser Asn Lys Thr Ile Ile Asp Ile Val Asn Phe
 50 55 60

Asp Glu Trp Lys Ala Gln Glu Gln Lys Arg Leu Val Glu Leu Leu Lys
 65 70 75 80

Asn Val Pro Glu Asp Val His Ile Phe Ile Arg Ser Gln Lys Thr Gly
 85 90 95

Gly Lys Gly Val Ala Leu Glu Leu Pro Lys Pro Trp Glu Thr Asp Lys
 100 105 110

Trp Leu Glu Trp Ile Glu Lys Arg Phe Arg Glu Asn Gly Leu Leu Ile
 115 120 125

Asp Lys Asp Ala Leu Gln Leu Phe Phe Ser Lys Val Gly Thr Asn Asp
 130 135 140

Leu Ile Ile Glu Arg Glu Ile Glu Lys Leu Lys Ala Tyr Ser Glu Asp
 145 150 155 160

Arg Lys Ile Thr Val Glu Asp Val Glu Glu Val Val Phe Thr Tyr Gln
 165 170 175

Thr Pro Gly Tyr Asp Asp Phe Cys Phe Ala Val Ser Glu Gly Lys Arg
 180 185 190

Lys Leu Ala His Ser Leu Leu Ser Gln Leu Trp Lys Thr Thr Glu Ser
 195 200 205

Val Val Ile Ala Thr Val Leu Ala Asn His Phe Leu Asp Leu Phe Lys
 210 215 220

Ile Leu Val Leu Val Thr Lys Lys Arg Tyr Tyr Thr Trp Pro Asp Val
 225 230 235 240

Ser Arg Val Ser Lys Glu Leu Gly Ile Pro Val Pro Arg Val Ala Arg
 245 250 255

Phe Leu Gly Phe Ser Phe Lys Thr Trp Lys Phe Lys Val Met Asn His
 260 265 270

Leu Leu Tyr Tyr Asp Val Lys Lys Val Arg Lys Ile Leu Arg Asp Leu
 275 280 285

Tyr Asp Leu Asp Arg Ala Val Lys Ser Glu Glu Asp Pro Lys Pro Phe
 290 295 300

Phe His Glu Phe Ile Glu Glu Val Ala Leu Asp Val Tyr Ser Leu Gln
 305 310 315 320

Arg Asp Glu Glu

<210> 147
 <211> 936
 <212> DNA
 <213> *Thermatoga maritima*

<400> 147
 atgaacgatt tgatcagaaa gtacgctaaa gatcaactgg aaactttgaa aaggatcata 60
 gaaaaagtctg aaggaatatc catcctcata aatggagaag atctctcgta tccgagagaa 120
 gtatcccttg aacttcccga gtacgtggag aaatttcccc cgaaggcctc ggatgttctg 180
 gagatagatc ccgaggggga gaacataggc atagacgaca tcagaacgat aaaggacttc 240
 ctgaactaca gccccgagct ctacacgaga aagtacgtga tagtccacga ctgtgaaaga 300
 atgaccagc aggcggcgaa cgcgtttctg aaggcccttg aagaaccacc agaatacgct 360
 gtgatcgctt tgaacactcg ccgctggcat tatctactgc cgacgataaa gagccgagtg 420
 ttcagagtgg ttgtgaacgt tccaaaggag ttcagagatc tcgtgaaaga gaaaatagga 480
 gatctctggg aggaacttcc acttcttgag agagacttca aaacggctct cgaagcctac 540
 aaacttggtg cggaaaaact ttctggattg atggaaagtc tcaaagtttt ggagacggaa 600
 aaactcttga aaaaggtcct ttcaaaaggc ctcgaagggt atctcgcatg tagggagctc 660
 ctggagagat tttcaaaggt ggaatcgaag gaattctttg cgctttttga tcaggtgact 720
 aacacgataa caggaaaaga cgcgtttctt ttgatccaga gactgacaag aatcattctc 780
 cacgaaaaca catgggaaag cgttgaagat caaaaaagcg tgtctttcct cgattcaatt 840
 ctcagggtga agatagcgaa tctgaacaac aaactcactc tgatgaacat cctcgcgata 900
 cacagagaga gaaagagagg tgtcaacgct tggagc 936

<210> 148
 <211> 311
 <212> PRT
 <213> *Thermatoga maritima*

<400> 148
 Met Asn Asp Leu Ile Arg Lys Tyr Ala Lys Asp Gln Leu Glu Thr Leu
 1 5 10 15
 Lys Arg Ile Ile Glu Lys Ser Glu Gly Ile Ser Ile Leu Ile Asn Gly
 20 25 30
 Glu Asp Leu Ser Tyr Pro Arg Glu Val Ser Leu Glu Leu Pro Glu Tyr
 35 40 45
 Val Glu Lys Phe Pro Pro Lys Ala Ser Asp Val Leu Glu Ile Asp Pro
 50 55 60
 Glu Gly Glu Asn Ile Gly Ile Asp Asp Ile Arg Thr Ile Lys Asp Phe
 65 70 75 80
 Leu Asn Tyr Ser Pro Glu Leu Tyr Thr Arg Lys Tyr Val Ile Val His
 85 90 95
 Asp Cys Glu Arg Met Thr Gln Gln Ala Ala Asn Ala Phe Leu Lys Ala
 100 105 110
 Leu Glu Glu Pro Pro Glu Tyr Ala Val Ile Val Leu Asn Thr Arg Arg
 115 120 125
 Trp His Tyr Leu Leu Pro Thr Ile Lys Ser Arg Val Phe Arg Val Val
 130 135 140
 Val Asn Val Pro Lys Glu Phe Arg Asp Leu Val Lys Glu Lys Ile Gly
 145 150 155 160
 Asp Leu Trp Glu Glu Leu Pro Leu Leu Glu Arg Asp Phe Lys Thr Ala
 165 170 175
 Leu Glu Ala Tyr Lys Leu Gly Ala Glu Lys Leu Ser Gly Leu Met Glu
 180 185 190
 Ser Leu Lys Val Leu Glu Thr Glu Lys Leu Leu Lys Lys Val Leu Ser
 195 200 205
 Lys Gly Leu Glu Gly Tyr Leu Ala Cys Arg Glu Leu Leu Glu Arg Phe
 210 215 220

Ile Ala Val Asp Arg Val Pro Arg Lys Asn Ala Pro Asp Asp Ala Gln
 35 40 45

Thr Thr Asp Phe Phe Arg Ile Val Thr Phe Gly Arg Leu Ala Glu Phe
 50 55 60

Ala Arg Thr Tyr Leu Thr Lys Gly Arg Leu Val Leu Val Glu Gly Glu
 65 70 75 80

Met Arg Met Arg Arg Trp Glu Thr Pro Thr Gly Glu Lys Arg Val Ser
 85 90 95

Pro Glu Val Val Ala Asn Val Val Arg Phe Met Asp Arg Lys Pro Ala
 100 105 110

Glu Thr Val Ser Glu Thr Glu Glu Glu Leu Glu Ile Pro Glu Glu Asp
 115 120 125

Phe Ser Ser Asp Thr Phe Ser Glu Asp Glu Pro Pro Phe
 130 135 140

<210> 151

<211> 1353

<212> DNA

<213> *Thermatoga maritima*

<400> 151

atgcgtgttc cccgcacaa cttagaggcc gaagttgctg tgctcggaag catattgata 60
 gatccgctcg taataaacga cgttcttgaa attttgagcc acgaagattt ctatctgaaa 120
 aaacaccaac acatcttcag agcgatggaa gagctttacg acgaaggaaa accggtggac 180
 gtggtttccg tctgtgacaa gcttcaaagc atgggaaaac tcgaggaagt aggtggagat 240
 ctggaagtgg ccagctcgc tgaggctgtg ccagttctg cacacgcact tcactacgcg 300
 gagatcgtca aggaaaaatc cattctgagg aaactcattg agatctccag aaaaatctca 360
 gaaagtgcct acatggaaga agatgtggag atcctgctcg acaacgcaga aaagatgatc 420
 ttcgagatct cagagatgaa aacgacaaaa tcctacgata atctgagagg catcatgcac 480
 cgggtgtttg aaaacctgga gaacttcagg gaaagagcca accttataga acccggtgtg 540
 ctcataacgg gactaccaac gggattcaaa agtctggaca aacagaccac agggttccac 600
 agctccgatac tggtgataat agcagcgaga cctccatgg gaaaaacctc cttcgactc 660
 tcaatagcga ggaacatggc tgtcaatttc gaaatccccg tcggaatatt cagtctcgag 720
 atgtccaagg aacagctcgc tcaaagacta ctcagcatgg agtccggtgt ggatctttac 780
 agcatcagaa caggatacct ggatcaggag aagtgggaaa gactcacaat agcggcttct 840
 aaactctaca aagcacccat agttgtggac gatgagtcac tctcgcgatcc gcgatcgttg 900
 agggcaaaaag cgagaaggat gaaaaaagaa tacgatgtaa aagccatttt tgtcgactat 960
 ctccagctca tgcacctgaa aggaagaaaa gaaagcagac agcaggagat atccgagatc 1020
 tcgagatctc tgaagctcct tgcgagggaa ctcgacatag tggatgtagc gctttcacag 1080
 ctttcgaggg ccgtagaaca gagagaagac aaaagaccga ggctgagtga cctcagggaa 1140
 tccggtgcga tagaacagga cgcagacaca gtcattctca tctacaggga ggaatattac 1200

aggagcaaaa aatccaaaga ggaaagcaag cttcacgaac ctcacgaage tgaaatcata 1260
 ataggtaaac agagaaacgg tcccgttgga acgatcactc tgatcttcga cccagaacg 1320
 gttacgttcc atgaagtcga tgtggtgcat tca 1353

<210> 152

<211> 451

<212> PRT

<213> *Thermatoga maritima*

<400> 152

Met Arg Val Pro Pro His Asn Leu Glu Ala Glu Val Ala Val Leu Gly
 1 5 10 15

Ser Ile Leu Ile Asp Pro Ser Val Ile Asn Asp Val Leu Glu Ile Leu
 20 25 30

Ser His Glu Asp Phe Tyr Leu Lys Lys His Gln His Ile Phe Arg Ala
 35 40 45

Met Glu Glu Leu Tyr Asp Glu Gly Lys Pro Val Asp Val Val Ser Val
 50 55 60

Cys Asp Lys Leu Gln Ser Met Gly Lys Leu Glu Glu Val Gly Gly Asp
 65 70 75 80

Leu Glu Val Ala Gln Leu Ala Glu Ala Val Pro Ser Ser Ala His Ala
 85 90 95

Leu His Tyr Ala Glu Ile Val Lys Glu Lys Ser Ile Leu Arg Lys Leu
 100 105 110

Ile Glu Ile Ser Arg Lys Ile Ser Glu Ser Ala Tyr Met Glu Glu Asp
 115 120 125

Val Glu Ile Leu Leu Asp Asn Ala Glu Lys Met Ile Phe Glu Ile Ser
 130 135 140

Glu Met Lys Thr Thr Lys Ser Tyr Asp His Leu Arg Gly Ile Met His
 145 150 155 160

Arg Val Phe Glu Asn Leu Glu Asn Phe Arg Glu Arg Ala Asn Leu Ile
 165 170 175

Glu Pro Gly Val Leu Ile Thr Gly Leu Pro Thr Gly Phe Lys Ser Leu
 180 185 190

Asp Lys Gln Thr Thr Gly Phe His Ser Ser Asp Leu Val Ile Ile Ala

195	200	205
Ala Arg Pro Ser Met Gly Lys Thr Ser Phe Ala Leu Ser Ile Ala Arg		
210	215	220
Asn Met Ala Val Asn Phe Glu Ile Pro Val Gly Ile Phe Ser Leu Glu		
225	230	235 240
Met Ser Lys Glu Gln Leu Ala Gln Arg Leu Leu Ser Met Glu Ser Gly		
	245	250 255
Val Asp Leu Tyr Ser Ile Arg Thr Gly Tyr Leu Asp Gln Glu Lys Trp		
	260	265 270
Glu Arg Leu Thr Ile Ala Ala Ser Lys Leu Tyr Lys Ala Pro Ile Val		
	275	280 285
Val Asp Asp Glu Ser Leu Leu Asp Pro Arg Ser Leu Arg Ala Lys Ala		
	290	295 300
Arg Arg Met Lys Lys Glu Tyr Asp Val Lys Ala Ile Phe Val Asp Tyr		
305	310	315 320
Leu Gln Leu Met His Leu Lys Gly Arg Lys Glu Ser Arg Gln Gln Glu		
	325	330 335
Ile Ser Glu Ile Ser Arg Ser Leu Lys Leu Leu Ala Arg Glu Leu Asp		
	340	345 350
Ile Val Val Ile Ala Leu Ser Gln Leu Ser Arg Ala Val Glu Gln Arg		
	355	360 365
Glu Asp Lys Arg Pro Arg Leu Ser Asp Leu Arg Glu Ser Gly Ala Ile		
	370	375 380
Glu Gln Asp Ala Asp Thr Val Ile Phe Ile Tyr Arg Glu Glu Tyr Tyr		
385	390	395 400
Arg Ser Lys Lys Ser Lys Glu Glu Ser Lys Leu His Glu Pro His Glu		
	405	410 415
Ala Glu Ile Ile Ile Gly Lys Gln Arg Asn Gly Pro Val Gly Thr Ile		
	420	425 430
Thr Leu Ile Phe Asp Pro Arg Thr Val Thr Phe His Glu Val Asp Val		
	435	440 445
Val His Ser		

450

<210> 153
 <211> 1695
 <212> DNA
 <213> *Thermatoga maritima*

<400> 153
 gtgattcctc gagaggatcat cgaggaaata aaagaaaagg ttgacatcgt agaggtcatt 60
 tccgagtacg tgaatcttac ccgggtaggt tctcctaca gggctctctg tccctttcat 120
 tcagaaacca atccttcttt ctacgttcat ccgggtttga agatatacca ttgtttcggc 180
 tggcgtgcga gtggagacgt catcaaattt cttcaagaaa tgggaaggat cagtttccag 240
 gaagcgctgg aaagacttgc caaaagagct gggattgatc tttctctcta cagaacagaa 300
 gggacttctg aatacggaaa atacattcgt ttgtacgaag aaacgtggaa aaggtagctc 360
 aaagagctgg agaaatcgaa agaggcaaaa gactatttaa aaagcagagg cttctctgaa 420
 gaagatatag caaagtccg ctttgggtac gtccccaaga gatccagcat ctctatagaa 480
 gttgcagaag gcatgaacat aacactggaa gaacttgtca gatacggat cgcgctgaaa 540
 aagggtgatc gattcgttga tagattcgaa ggaagaatcg ttgttccaat aaagaacgac 600
 agtggtcata ttgtggcctt tgggtggcgt gctctcggca acgaagaacc gaagtatttg 660
 aactctccag agaccaggta tttttcgaag aagaagacc ttttctctt cgatgaggcg 720
 aaaaaagtg caaaagaggt tgggtttttc gtcatcaccc aaggctactt cgacgcgctc 780
 gcattcagaa aggatggaat accaacggcg gtcgctgttc ttggggcgag tctttcaaga 840
 gaggcgatcc taaaactttc ggcgtattcg aaaaacgtca tactgtgttt cgataatgac 900
 aaagcaggct tcagagccac tctcaaattc ctcgaggatc tcctagacta cgaattcaac 960
 gtgcttgttg caacccctc tctttacaaa gaccagatg aactctttca gaaagaagga 1020
 gaaggttcat tgaaaaagat gctgaaaaac tcgcgttcgt tcgaatattt tctggtgacg 1080
 gctggtgagg tcttctttga caggaacagc ccgcgggtg tgagatccta ctttctttc 1140
 ctcaaagggt ggggtccaaa gatgagaagg aaaggatatt tgaaacacat agaaaatctc 1200
 gtgaatgagg tttcatcttc tctccagata ccagaaaacc agattttgaa cttttttgaa 1260
 agcgacaggt ctaacactat gcctgttcat gagaccaagt cgtcaaaggt ttacgatgag 1320
 gggagaggac tggcttattt gtttttgaac tacgaggatt tgagggaaaa gattctggaa 1380
 ctggacttag aggtactgga agataaaaa gcgaggaggt ttttcaagag agtctcactg 1440
 ggagaagatt tgaacaaagt catagaaaac ttcccaaaag agctgaaaga ctggattttt 1500
 gagacaatag aaagcattcc tctccaaag gatcccgaga aattcctcgg tgacctctcc 1560
 gaaaagttga aaatccgacg gatagagaga cgtatcgag aaatagatga tatgataaag 1620
 aaagcttcaa acgatgaaga aaggcgtctt cttctctcta tgaaagtgga tctcctcaga 1680
 aaaataaaga ggagg 1695

<210> 154
 <211> 565
 <212> PRT
 <213> *Thermatoga maritima*

<400> 154
 Met Ile Pro Arg Glu Val Ile Glu Glu Ile Lys Glu Lys Val Asp Ile
 1 5 10 15

Val	Glu	Val	Ile	Ser	Glu	Tyr	Val	Asn	Leu	Thr	Arg	Val	Gly	Ser	Ser	20	25	30
Tyr	Arg	Ala	Leu	Cys	Pro	Phe	His	Ser	Glu	Thr	Asn	Pro	Ser	Phe	Tyr	35	40	45
Val	His	Pro	Gly	Leu	Lys	Ile	Tyr	His	Cys	Phe	Gly	Cys	Gly	Ala	Ser	50	55	60
Gly	Asp	Val	Ile	Lys	Phe	Leu	Gln	Glu	Met	Glu	Gly	Ile	Ser	Phe	Gln	65	70	75
Glu	Ala	Leu	Glu	Arg	Leu	Ala	Lys	Arg	Ala	Gly	Ile	Asp	Leu	Ser	Leu	85	90	95
Tyr	Arg	Thr	Glu	Gly	Thr	Ser	Glu	Tyr	Gly	Lys	Tyr	Ile	Arg	Leu	Tyr	100	105	110
Glu	Glu	Thr	Trp	Lys	Arg	Tyr	Val	Lys	Glu	Leu	Glu	Lys	Ser	Lys	Glu	115	120	125
Ala	Lys	Asp	Tyr	Leu	Lys	Ser	Arg	Gly	Phe	Ser	Glu	Glu	Asp	Ile	Ala	130	135	140
Lys	Phe	Gly	Phe	Gly	Tyr	Val	Pro	Lys	Arg	Ser	Ser	Ile	Ser	Ile	Glu	145	150	155
Val	Ala	Glu	Gly	Met	Asn	Ile	Thr	Leu	Glu	Glu	Leu	Val	Arg	Tyr	Gly	165	170	175
Ile	Ala	Leu	Lys	Lys	Gly	Asp	Arg	Phe	Val	Asp	Arg	Phe	Glu	Gly	Arg	180	185	190
Ile	Val	Val	Pro	Ile	Lys	Asn	Asp	Ser	Gly	His	Ile	Val	Ala	Phe	Gly	195	200	205
Gly	Arg	Ala	Leu	Gly	Asn	Glu	Glu	Pro	Lys	Tyr	Leu	Asn	Ser	Pro	Glu	210	215	220
Thr	Arg	Tyr	Phe	Ser	Lys	Lys	Lys	Thr	Leu	Phe	Leu	Phe	Asp	Glu	Ala	225	230	235
Lys	Lys	Val	Ala	Lys	Glu	Val	Gly	Phe	Phe	Val	Ile	Thr	Glu	Gly	Tyr	245	250	255
Phe	Asp	Ala	Leu	Ala	Phe	Arg	Lys	Asp	Gly	Ile	Pro	Thr	Ala	Val	Ala	260	265	270

Val Leu Gly Ala Ser Leu Ser Arg Glu Ala Ile Leu Lys Leu Ser Ala
 275 280 285
 Tyr Ser Lys Asn Val Ile Leu Cys Phe Asp Asn Asp Lys Ala Gly Phe
 290 295 300
 Arg Ala Thr Leu Lys Ser Leu Glu Asp Leu Leu Asp Tyr Glu Phe Asn
 305 310 315 320
 Val Leu Val Ala Thr Pro Ser Pro Tyr Lys Asp Pro Asp Glu Leu Phe
 325 330 335
 Gln Lys Glu Gly Glu Gly Ser Leu Lys Lys Met Leu Lys Asn Ser Arg
 340 345 350
 Ser Phe Glu Tyr Phe Leu Val Thr Ala Gly Glu Val Phe Phe Asp Arg
 355 360 365
 Asn Ser Pro Ala Gly Val Arg Ser Tyr Leu Ser Phe Leu Lys Gly Trp
 370 375 380
 Val Gln Lys Met Arg Arg Lys Gly Tyr Leu Lys His Ile Glu Asn Leu
 385 390 395 400
 Val Asn Glu Val Ser Ser Ser Leu Gln Ile Pro Glu Asn Gln Ile Leu
 405 410 415
 Asn Phe Phe Glu Ser Asp Arg Ser Asn Thr Met Pro Val His Glu Thr
 420 425 430
 Lys Ser Ser Lys Val Tyr Asp Glu Gly Arg Gly Leu Ala Tyr Leu Phe
 435 440 445
 Leu Asn Tyr Glu Asp Leu Arg Glu Lys Ile Leu Glu Leu Asp Leu Glu
 450 455 460
 Val Leu Glu Asp Lys Asn Ala Arg Glu Phe Phe Lys Arg Val Ser Leu
 465 470 475 480
 Gly Glu Asp Leu Asn Lys Val Ile Glu Asn Phe Pro Lys Glu Leu Lys
 485 490 495
 Asp Trp Ile Phe Glu Thr Ile Glu Ser Ile Pro Pro Pro Lys Asp Pro
 500 505 510
 Glu Lys Phe Leu Gly Asp Leu Ser Glu Lys Leu Lys Ile Arg Arg Ile
 515 520 525

Glu Arg Arg Ile Ala Glu Ile Asp Asp Met Ile Lys Lys Ala Ser Asn
530 535 540

Asp Glu Glu Arg Arg Leu Leu Leu Ser Met Lys Val Asp Leu Leu Arg
545 550 555 560

Lys Ile Lys Arg Arg
565

<210> 155

<211> 804

<212> DNA

<213> *Thermus thermophilus*

<400> 155

```
atggctctac acccggtcca ccctggggca ataatcgggc acgaggccgt tctcgccctc 60
cttccccgcc tcaccgcca gacctgctc ttctccggcc ccgaggggt ggggcggcgc 120
accgtggccc gctggtacgc ctgggggctc aaccgcggt tcccccgcc ctccctgggg 180
gagcaccggg acgtcctcga ggtggggccc aaggccggg acctccggg ccgggcccag 240
gtgcggctgg aggaggtggc gccctcttg gagtgggtgt ccagccacc ccgggagcgg 300
gtgaaggtgg ccatcctgga ctgggcccac ctctcaccg aggcggccgc caacgccctc 360
ctcaagctcc tggaggagcc cccttcctac gcccgcatcg tctcatcgc cccaagccgc 420
gccaccctcc tccccaccct ggccctcccg gccacggagg tggcattcgc cccgtgccc 480
gaggaggccc tgcgcgccct caccaggac ccggagctcc tccgctacgc cgccggggcc 540
ccgggcccgc tccttagggc cctccaggac ccggaggggt accgggccc catggccagg 600
gcgcaaaggg tcctgaaagc cccgccctg gagcgctcg ctttgcttcg ggagcttttg 660
gccgaggagg agggggtcca cgccctccac gccgtcctaa agcgcccgga gcacctcctt 720
gccctggagc gggcgcgggg ggccctggag gggtagctga gccccgagct ggtcctcgcc 780
cggctggcct tagacttaga gaca 804
```

<210> 156

<211> 268

<212> PRT

<213> *Thermus thermophilus*

<400> 156

Met Ala Leu His Pro Ala His Pro Gly Ala Ile Ile Gly His Glu Ala
1 5 10 15

Val Leu Ala Leu Leu Pro Arg Leu Thr Ala Gln Thr Leu Leu Phe Ser
20 25 30

Gly Pro Glu Gly Val Gly Arg Arg Thr Val Ala Arg Trp Tyr Ala Trp
35 40 45

Gly Leu Asn Arg Gly Phe Pro Pro Pro Ser Leu Gly Glu His Pro Asp
 50 55 60
 Val Leu Glu Val Gly Pro Lys Ala Arg Asp Leu Arg Gly Arg Ala Glu
 65 70 75 80
 Val Arg Leu Glu Glu Val Ala Pro Leu Leu Glu Trp Cys Ser Ser His
 85 90 95
 Pro Arg Glu Arg Val Lys Val Ala Ile Leu Asp Ser Ala His Leu Leu
 100 105 110
 Thr Glu Ala Ala Ala Asn Ala Leu Leu Lys Leu Leu Glu Glu Pro Pro
 115 120 125
 Ser Tyr Ala Arg Ile Val Leu Ile Ala Pro Ser Arg Ala Thr Leu Leu
 130 135 140
 Pro Thr Leu Ala Ser Arg Ala Thr Glu Val Ala Phe Ala Pro Val Pro
 145 150 155 160
 Glu Glu Ala Leu Arg Ala Leu Thr Gln Asp Pro Glu Leu Leu Arg Tyr
 165 170 175
 Ala Ala Gly Ala Pro Gly Arg Leu Leu Arg Ala Leu Gln Asp Pro Glu
 180 185 190
 Gly Tyr Arg Ala Arg Met Ala Arg Ala Gln Arg Val Leu Lys Ala Pro
 195 200 205
 Pro Leu Glu Arg Leu Ala Leu Leu Arg Glu Leu Leu Ala Glu Glu Glu
 210 215 220
 Gly Val His Ala Leu His Ala Val Leu Lys Arg Pro Glu His Leu Leu
 225 230 235 240
 Ala Leu Glu Arg Ala Arg Glu Ala Leu Glu Gly Tyr Val Ser Pro Glu
 245 250 255
 Leu Val Leu Ala Arg Leu Ala Leu Asp Leu Glu Thr
 260 265

<210> 157
 <211> 729
 <212> DNA
 <213> Thermus thermophilus

<400> 157

```

atgctggacc tgagggaggt gggggaggcg gagtgggaagg ccctaaagcc ccttttggaa 60
agcgtgcccc agggcgctccc cgtcctcctc ctggacccta agccaagccc ctcccgggcg 120
gccttctacc ggaaccggga aaggcgggac ttccccaccc ccaaggggaa ggacctggtg 180
cggcacctgg aaaaccgggc caagcgctg gggctcaggc tcccgggagg ggtggcccag 240
tacctggcct ccctggaggg ggacctcgag gccctggagc gggagctgga gaagcttgcc 300
ctcctctccc caccctcac cctggagaag gtggagaagg tggtagccct gagggcccc 360
ctcacgggct ttgacctggt gcgctccgtc ctggagaagg accccaagga ggccctcctg 420
cgcctaggcg gcctcaagga ggagggggag gagccctca ggctcctcgg ggccctctcc 480
tggcagttcg ccctcctcgc ccgggccttc ttctcctcc gggaaaaccc caggcccaag 540
gaggaggacc tcgcccgcct cgaggccac ccctacgccg ccgcgcgcgc cctggaggcg 600
gcgaagcgcc tcacggaaga ggccctcaag gaggccttg acgcctcat ggaggcgga 660
aagagggcca agggggggaa agaccgtgg ctgccttg aggcggcggt cctccgcctc 720
gcccgttga 729

```

<210> 158

<211> 292

<212> PRT

<213> *Thermus thermophilus*

<400> 158

```

Met Val Ile Ala Phe Thr Gly Asp Pro Phe Leu Ala Arg Glu Ala Leu
  1             5             10             15

Leu Glu Glu Ala Arg Leu Arg Gly Leu Ser Arg Phe Thr Glu Pro Thr
      20             25             30

Pro Glu Ala Leu Ala Gln Ala Leu Ala Pro Gly Leu Phe Gly Gly Gly
      35             40             45

Gly Ala Met Leu Asp Leu Arg Glu Val Gly Glu Ala Glu Trp Lys Ala
      50             55             60

Leu Lys Pro Leu Leu Glu Ser Val Pro Glu Gly Val Pro Val Leu Leu
      65             70             75             80

Leu Asp Pro Lys Pro Ser Pro Ser Arg Ala Ala Phe Tyr Arg Asn Arg
      85             90             95

Glu Arg Arg Asp Phe Pro Thr Pro Lys Gly Lys Asp Leu Val Arg His
      100            105            110

Leu Glu Asn Arg Ala Lys Arg Leu Gly Leu Arg Leu Pro Gly Gly Val
      115            120            125

Ala Gln Tyr Leu Ala Ser Leu Glu Gly Asp Leu Glu Ala Leu Glu Arg
      130            135            140

```

Glu Leu Glu Lys Leu Ala Leu Leu Ser Pro Pro Leu Thr Leu Glu Lys
145 150 155 160

Val Glu Lys Val Val Ala Leu Arg Pro Pro Leu Thr Gly Phe Asp Leu
165 170 175

Val Arg Ser Val Leu Glu Lys Asp Pro Lys Glu Ala Leu Leu Arg Leu
180 185 190

Gly Gly Leu Lys Glu Glu Gly Glu Glu Pro Leu Arg Leu Leu Gly Ala
195 200 205

Leu Ser Trp Gln Phe Ala Leu Leu Ala Arg Ala Phe Phe Leu Leu Arg
210 215 220

Glu Asn Pro Arg Pro Lys Glu Glu Asp Leu Ala Arg Leu Glu Ala His
225 230 235 240

Pro Tyr Ala Ala Arg Arg Ala Leu Glu Ala Ala Lys Arg Leu Thr Glu
245 250 255

Glu Ala Leu Lys Glu Ala Leu Asp Ala Leu Met Glu Ala Glu Lys Arg
260 265 270

Ala Lys Gly Gly Lys Asp Pro Trp Leu Ala Leu Glu Ala Ala Val Leu
275 280 285

Arg Leu Ala Arg
290

<210> 159

<211> 37

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 159

gtgtgtcata tgagtaagga ttctgtccac cttcacc

37

<210> 160

<211> 34

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 160

gtgtgtggat ccggggacta ctcggaagta aggg

34

<210> 161

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 161

gtgtgtcata tggaaaccac aatattccag ttccag

36

<210> 162

<211> 39

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 162

gtgtgtggat ccttatccac catgagaagt atttttcac

39

<210> 163

<211> 41

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 163

gtgtgtcata tggaaaaagt tttttttgga aaaaactcca g

41

<210> 164

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 168

tgtgtctcga gtcattggcta caccctcatc ggcat

35

<210> 169

<211> 47

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 169

gtgtgtcata tgctcaataa gggtttttata ataggaagac ttacggg

47

<210> 170

<211> 39

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 170

gtgtggatcc ttaaaaaggt atttcgtcct cttcatcgg

39

<210> 171

<211> 807

<212> DNA

<213> Thermus thermophilus

<400> 171

atggctcgag gcctgaaccg cgttttcctc atcggcgccc tcgccacccg gccggacatg 60
cgctacaccc cggcgggggt cgccattttg gacctgaccc tcgccggtca ggacctgctt 120
ctttccgata acggggggga accggaggtg tcctggtacc accgggtgag gctcttaggc 180
cgccaggcgg agatgtgggg cgacctcttg gaccaagggc agctcgtctt cgtggagggc 240
cgcctggagt accgccagt ggaagaggag ggggagaagc ggagcgagct ccagatccgg 300
gccgacttcc ggaccccctg gacgaccggg ggaagaagcg ggcgaggagc agccggggcc 360
agcccaggct ccgcgccgcc ctgaaccagg tcttcctcat gggcaacctg accggggacc 420
cggaactccg ctacaccccc cagggcaccg cgggtggccc gctgggcctg gcggtgaacg 480
agcgcgcgca gggggcggag gagcgcaccc acttcgtgga gggttcaggcc tggcgcgacc 540
tggcggagtg ggccgccgag ctgaggaagg gcgacggcct tttcgtgatc ggcaggttgg 600

Gly Leu Phe Val Ile Gly Arg Leu Val Asn Asp Ser Trp Thr Ser Ser
 195 200 205

Ser Gly Glu Arg Arg Phe Gln Thr Arg Val Glu Ala Leu Arg Leu Glu
 210 215 220

Arg Pro Thr Arg Gly Pro Ala Gln Ala Cys Pro Gly Arg Arg Asn Arg
 225 230 235 240

Ser Arg Glu Val Gln Thr Gly Gly Val Asp Ile Asp Glu Gly Leu Glu
 245 250 255

Asp Phe Pro Pro Glu Glu Asp Leu Pro Phe
 260 265

<210> 173

<211> 992

<212> DNA

<213> *Bacillus stearothermophilus*

<400> 173

aattccgaca tttcaattga atcggtttatt ccgcttgaaa aagaaggcaa gttgctcggtt 60
 gatgtgaaaa gaccggggag catcgactg caggcgcgct ttttctctga aatcgtgaaa 120
 aaactgccgc aacaaacggt ggaaatcgaa acggaagaca actttttgac gatcatccgc 180
 tcggggcact cagaattccg cctcaatggg ctaaacgccg acgaatatcc gcgcctgccg 240
 caaattgaag aagaaaacgt gtttcaaata ccggctgatt tattgaaaac cgtgattcgg 300
 caaacggtgt tcgccgtttc tacatcgga acgcgcccaa tcttgacagg tgtcaactgg 360
 aaagtgaac atggcgagct tgtctgcaca gcgaccgaca gtcacgctt agccatgcgc 420
 aaagtgaaaa ttgagtcgga aaatgaagta tcatacaacg tcgtcatccc tggaaaaagt 480
 cttaatgagc tcagcaaaat tttggatgac ggcaaccacc cgggtggacat cgtcatgaca 540
 gccaatcaag tgctatttaa ggccgagcac cttctcttct tttcccggtt gcttgacggc 600
 aactatccgg agacggcccg cttgattcca acagaaagca aaacgaccat gatcgtcaat 660
 gcaaaagagt ttctgcaggc aatcgaccga gcgtccttgc ttgctcgaga aggaaggaaac 720
 aacgttgtag aactgacgac gcttcctgga ggaatgctcg aaatttcttc gatttctccg 780
 agatcgggaa agtgacggag cagctgcaaa cggagtctct tgaaggggaa gagttgaaca 840
 tttegttcag cgcgaaatat atgatggacg cgttgccggc gcttgatgga acagacattt 900
 caaatcagct tcaactggggc catgcggccg ttctgtttgc gcccgcttca accgattcga 960
 tgcttcagct cattttgccg gtgagaacat at 992

<210> 174

<211> 334

<212> PRT

<213> *Bacillus stearothermophilus*

<400> 174

Asn Ser Asp Ile Ser Ile Ile Glu Ser Phe Ile Pro Leu Glu Lys Glu

007716964-112100

1	5	10	15
Gly Lys Leu Leu Val Asp Val Lys Arg Pro Gly Ser Ile Val Leu Gln	20	25	30
Ala Arg Phe Phe Ser Glu Ile Val Lys Lys Leu Pro Gln Gln Thr Val	35	40	45
Glu Ile Glu Thr Glu Asp Asn Phe Leu Thr Ile Ile Arg Ser Gly His	50	55	60
Ser Glu Phe Arg Leu Asn Gly Leu Asn Ala Asp Glu Tyr Pro Arg Leu	65	70	75
Pro Gln Ile Glu Glu Glu Asn Val Phe Gln Ile Pro Ala Asp Leu Leu	85	90	95
Lys Thr Val Ile Arg Gln Thr Val Phe Ala Val Ser Thr Ser Glu Thr	100	105	110
Arg Pro Ile Leu Thr Gly Val Asn Trp Lys Val Glu His Gly Glu Leu	115	120	125
Val Cys Thr Ala Thr Asp Ser His Arg Leu Ala Met Arg Lys Val Lys	130	135	140
Ile Ile Glu Ser Glu Asn Glu Val Ser Tyr Asn Val Val Ile Pro Gly	145	150	155
Lys Ser Leu Asn Glu Leu Ser Lys Ile Ile Leu Asp Asp Gly Asn His	165	170	175
Pro Val Asp Ile Val Met Thr Ala Asn Gln Val Leu Phe Lys Ala Glu	180	185	190
His Leu Leu Phe Phe Ser Arg Leu Leu Asp Gly Asn Tyr Pro Glu Thr	195	200	205
Ala Arg Leu Ile Pro Thr Glu Ser Lys Thr Thr Met Ile Val Asn Ala	210	215	220
Lys Glu Phe Leu Gln Ala Ile Asp Arg Ala Ser Leu Leu Ala Arg Glu	225	230	235
Gly Arg Asn Asn Val Val Lys Leu Thr Thr Leu Pro Gly Gly Met Leu	245	250	255
Glu Ile Ser Ser Ile Ser Pro Glu Ile Gly Lys Val Thr Glu Gln Leu			

260 265 270

Gln Thr Glu Ser Leu Glu Gly Glu Glu Leu Asn Ile Ser Phe Ser Ala
 275 280 285

Lys Tyr Met Met Asp Ala Leu Arg Ala Leu Asp Gly Thr Asp Ile Gln
 290 295 300

Ile Ser Phe Thr Gly Ala Met Arg Pro Phe Leu Leu Arg Pro Leu His
 305 310 315 320

Thr Asp Ser Met Leu Gln Leu Ile Leu Pro Val Arg Thr Tyr
 325 330

<210> 175
 <211> 492
 <212> DNA
 <213> Bacillus stearothermophilus

<400> 175

atgattaacc gcgtcatttt ggtcggcagg ttaacgagag atccggagtt gcgttacact 60
 ccaagcggag tggtgtgtgc cacgtttacg ctgcgggtca accgtccgtt tacaaatcag 120
 cagggcgagc gggaaacgga ttttattcaa tgtgtcgttt ggcgccgcca ggcggaaaac 180
 gtcgccaact ttttgaaaaa ggggagcttg gctggtgtcg atggccgact gcaaaccgcg 240
 agctatgaaa atcaagaagg tggcggtgtg tacgtgacgg aagtgggtggc tgatagcgtc 300
 caatttcttg agccgaaagg aacgagcgag cagcgagggg cgacagcagg cggctactat 360
 ggggatccat tcccattcgg gcaagatcag aaccaccaat atccgaacga aaaagggttt 420
 ggccgcacgc atgacgatcc tttcgccaat gacggccagc cgatcgatat ttctgatgat 480
 gatttgccgt tt 492

<210> 176
 <211> 164
 <212> PRT
 <213> Bacillus stearothermophilus

<400> 176

Met Ile Asn Arg Val Ile Leu Val Gly Arg Leu Thr Arg Asp Pro Glu
 1 5 10 15

Leu Arg Tyr Thr Pro Ser Gly Val Ala Val Ala Thr Phe Thr Leu Ala
 20 25 30

Val Asn Arg Pro Phe Thr Asn Gln Ser Tyr Glu Asn Gln Glu Gly Arg
 35 40 45

Arg Val Tyr Val Thr Glu Val Val Ala Asp Ser Val Gln Phe Leu Glu

50 55 60
 Pro Lys Gly Thr Ser Glu Gln Arg Gly Ala Thr Ala Gly Gly Tyr Tyr
 65 70 75 80
 Gln Gly Glu Arg Glu Thr Asp Phe Ile Gln Cys Val Val Trp Arg Arg
 85 90 95
 Gln Ala Glu Asn Val Ala Asn Phe Leu Lys Lys Gly Ser Leu Ala Gly
 100 105 110
 Val Asp Gly Arg Leu Gln Thr Arg Gly Asp Pro Phe Pro Phe Gly Gln
 115 120 125
 Asp Gln Asn His Gln Tyr Pro Asn Glu Lys Gly Phe Gly Arg Ile Asp
 130 135 140
 Asp Asp Pro Phe Ala Asn Asp Gly Gln Pro Ile Asp Ile Ser Asp Asp
 145 150 155 160
 Asp Leu Pro Phe

 <210> 177
 <211> 1044
 <212> DNA
 <213> *Bacillus stearothermophilus*

 <400> 177
 atgctggaac gcgtatgggg aaacattgaa aaacggcggt tttctcccct ttatttatta 60
 tacggcaatg agccgttttt attaacggaa acgtatgagc gattggtgaa cgcagcgctt 120
 ggccccgagg agcgggagtg gaacttggct gtgtacgact gcgaggaaac gccgatcgag 180
 gcggcgcttg aggaggccga gacggtgccg tttttcggcg agcggcggtg cattctcatc 240
 aagcatccat atttttttac gtctgaaaaa gagaaggaga tcgaacatga tttggcgaag 300
 ctggaggcgt acttgaaggc gccgtcgccg ttttcgatcg tcgtcttttt cgcgccgtac 360
 gagaagcttg atgagcgaaa aaaaattacg aagctcgcca aagagcaaag cgaagtcgtc 420
 atcgccgccc cgctcgccga agcggagctg cgtgcctggg tgcggcgccg catcgagagc 480
 caaggggctc aagcaagcga cgaggcgatt gatgtcctgt tgcggcgggc cgggacgcag 540
 ctttcgcgct tggcgaatga aatcgataaa ttggccctgt ttgccggatc gggcggaacc 600
 atcgaggcgg cggcggttga gcggcttgct gcccgcacgc cggaagaaaa cgtatttggtg 660
 cttgtcgagc aagtggcgaa gcgcgacatt ccagcagcgt tgcagacgtt ttatgatctg 720
 cttgaaaaca atgaagagcc gatcaaaatt ttggcggttg tcgccgccca tttccgcttg 780
 ctttcgcaag tgaaatggct tgccctccta ggctacggac aggcgcaaat tgctgcgggc 840
 ctcaaggtgc acccgttccg cgtcaagctc gctcttgctc aagcggcccc cttcgtgac 900
 ggagagcttg ctgaggcgat caacgagctc gctgacgccg attacgaagt gaaaagcggg 960
 gcggtcgatc gccggttggc cgttgagctg cttctgatgc gctggggcgc ccgcccggcg 1020
 caagcggggc gccacggccg gcgg 1044

<210> 178
 <211> 348
 <212> PRT
 <213> Bacillus stearothermophilus

<400> 178

Met	Leu	Glu	Arg	Val	Trp	Gly	Asn	Ile	Glu	Lys	Arg	Arg	Phe	Ser	Pro
1				5					10					15	
Leu	Tyr	Leu	Leu	Tyr	Gly	Asn	Glu	Pro	Phe	Leu	Leu	Thr	Glu	Thr	Tyr
			20					25					30		
Glu	Arg	Leu	Val	Asn	Ala	Ala	Leu	Gly	Pro	Glu	Glu	Arg	Glu	Trp	Asn
		35					40					45			
Leu	Ala	Val	Tyr	Asp	Cys	Glu	Glu	Thr	Pro	Ile	Glu	Ala	Ala	Leu	Glu
	50					55					60				
Glu	Ala	Glu	Thr	Val	Pro	Phe	Phe	Gly	Glu	Arg	Arg	Val	Ile	Leu	Ile
65					70					75					80
Lys	His	Pro	Tyr	Phe	Phe	Thr	Ser	Glu	Lys	Glu	Lys	Glu	Ile	Glu	His
				85					90					95	
Asp	Leu	Ala	Lys	Leu	Glu	Ala	Tyr	Leu	Lys	Ala	Pro	Ser	Pro	Phe	Ser
			100					105					110		
Ile	Val	Val	Phe	Phe	Ala	Pro	Tyr	Glu	Lys	Leu	Asp	Glu	Arg	Lys	Lys
		115					120					125			
Ile	Thr	Lys	Leu	Ala	Lys	Glu	Gln	Ser	Glu	Val	Val	Ile	Ala	Ala	Pro
	130					135					140				
Leu	Ala	Glu	Ala	Glu	Leu	Arg	Ala	Trp	Val	Arg	Arg	Arg	Ile	Glu	Ser
145					150					155					160
Gln	Gly	Ala	Gln	Ala	Ser	Asp	Glu	Ala	Ile	Asp	Val	Leu	Leu	Arg	Arg
				165					170					175	
Ala	Gly	Thr	Gln	Leu	Ser	Ala	Leu	Ala	Asn	Glu	Ile	Asp	Lys	Leu	Ala
			180					185					190		
Leu	Phe	Ala	Gly	Ser	Gly	Gly	Thr	Ile	Glu	Ala	Ala	Ala	Val	Glu	Arg
		195					200					205			
Leu	Val	Ala	Arg	Thr	Pro	Glu	Glu	Asn	Val	Phe	Val	Leu	Val	Glu	Gln

210 215 220
 Val Ala Lys Arg Asp Ile Pro Ala Ala Leu Gln Thr Phe Tyr Asp Leu
 225 230 235 240
 Leu Glu Asn Asn Glu Glu Pro Ile Lys Ile Leu Ala Leu Leu Ala Ala
 245 250 255
 His Phe Arg Leu Leu Ser Gln Val Lys Trp Leu Ala Ser Leu Gly Tyr
 260 265 270
 Gly Gln Ala Gln Ile Ala Ala Ala Leu Lys Val His Pro Phe Arg Val
 275 280 285
 Lys Leu Ala Leu Ala Gln Ala Ala Arg Phe Ala Asp Gly Glu Leu Ala
 290 295 300
 Glu Ala Ile Asn Glu Leu Ala Asp Ala Asp Tyr Glu Val Lys Ser Gly
 305 310 315 320
 Ala Val Asp Arg Arg Leu Ala Val Glu Leu Leu Leu Met Arg Trp Gly
 325 330 335
 Ala Arg Pro Ala Gln Ala Gly Arg His Gly Arg Arg
 340 345

<210> 179

<211> 757

<212> DNA

<213> *Bacillus stearothermophilus*

<400> 179

atgcgatggg aacagctagc gaaacgccag ccggtggtgg cgaaaatgct gcaaagcggc 60
 ttggaaaaag ggcgatttc tcatgcgtac ttgtttgagg ggcagcgggg gacgggcaaa 120
 aaagcggcca gtttgttgtt ggcgaaacgt ttgttttgtc tgtccccaat cggagtttcc 180
 ccgtgtctag agtgccgcaa ctgccggcgc atcgactccg gcaaccaccc tgacgtccgg 240
 gtgatcggcc cagatggagg atcaatcaaa aaggaacaaa tcgaatggct gcagcaagag 300
 ttctcgaaaa cagcggtcga gtcggataaa aaaatgtaca tcgttgagca cgccgatcaa 360
 atgacgacaa gcgctgccaa cagccttctg aaatttttgg aagagccgca tccggggacg 420
 gtggcggtat tgctgactga gcaataccac cgcctgctag ggacgatcgt ttcccgtgt 480
 caagtgcctt cgttccggcc gttgccgccg gcagagctcg cccagggact tgtcgaggag 540
 cacgtgccgt tgccgttggc gctgttggct gcccatattga caaacagctt cgaggaagca 600
 ctggcgcttg ccaaagatag ttggttgcc gaggcgcgaa cattagtgt acaatggtat 660
 gagatgctgg gcaagccgga gctgcagctt ttgtttttca tccacgaccg cttgtttccg 720
 ctttttttgg aaagccatca gcttgacctt ggacttg 757

<210> 180
 <211> 252
 <212> PRT
 <213> *Bacillus stearothermophilus*

<400> 180

Met	Arg	Trp	Glu	Gln	Leu	Ala	Lys	Arg	Gln	Pro	Val	Val	Ala	Lys	Met
1				5					10					15	
Leu	Gln	Ser	Gly	Leu	Glu	Lys	Gly	Arg	Ile	Ser	His	Ala	Tyr	Leu	Phe
			20					25					30		
Glu	Gly	Gln	Arg	Gly	Thr	Gly	Lys	Lys	Ala	Ala	Ser	Leu	Leu	Leu	Ala
		35					40					45			
Lys	Arg	Leu	Phe	Cys	Leu	Ser	Pro	Ile	Gly	Val	Ser	Pro	Cys	Leu	Glu
	50					55					60				
Cys	Arg	Asn	Cys	Arg	Arg	Ile	Asp	Ser	Gly	Asn	His	Pro	Asp	Val	Arg
	65				70					75					80
Val	Ile	Gly	Pro	Asp	Gly	Gly	Ser	Ile	Lys	Lys	Glu	Gln	Ile	Glu	Trp
				85					90					95	
Leu	Gln	Gln	Glu	Phe	Ser	Lys	Thr	Ala	Val	Glu	Ser	Asp	Lys	Lys	Met
			100					105					110		
Tyr	Ile	Val	Glu	His	Ala	Asp	Gln	Met	Thr	Thr	Ser	Ala	Ala	Asn	Ser
		115					120					125			
Leu	Leu	Lys	Phe	Leu	Glu	Glu	Pro	His	Pro	Gly	Thr	Val	Ala	Val	Leu
	130					135					140				
Leu	Thr	Glu	Gln	Tyr	His	Arg	Leu	Leu	Gly	Thr	Ile	Val	Ser	Arg	Cys
145					150					155					160
Gln	Val	Leu	Ser	Phe	Arg	Pro	Leu	Pro	Pro	Ala	Glu	Leu	Ala	Gln	Gly
				165					170					175	
Leu	Val	Glu	Glu	His	Val	Pro	Leu	Pro	Leu	Ala	Leu	Leu	Ala	Ala	His
			180				185						190		
Leu	Thr	Asn	Ser	Phe	Glu	Glu	Ala	Leu	Ala	Leu	Ala	Lys	Asp	Ser	Trp
		195					200					205			
Phe	Ala	Glu	Ala	Arg	Thr	Leu	Val	Leu	Gln	Trp	Tyr	Glu	Met	Leu	Gly
	210					215					220				

Lys Pro Glu Leu Gln Leu Leu Phe Phe Ile His Asp Arg Leu Phe Pro
 225 230 235 240

His Phe Leu Glu Ser His Gln Leu Asp Leu Gly Leu
 245 250

<210> 181
 <211> 1677
 <212> DNA
 <213> *Bacillus stearothermophilus*

<400> 181
 gtggcataacc aagcggttata tcgcgtgttt cggccgcagc gctttgcgga catggtcggc 60
 caagaacacg tgaccaagac gttgcaaagc gccctgcttc aacataaaat atcgcacgct 120
 tacttatttt cgggcccgcg cggtagagga aaaacgagcg cagcgaaaat ttctcgccaag 180
 gcggtcaact gtgaacaggc gccagcggcg gagccatgca atgagtgtcc agcttgccctc 240
 ggcattacga atggaacggt tcccgatgtg ctggaaattg acgctgcttc caacaaccgc 300
 gtcgatgaaa ttcgtgatat ccgtgagaag gtgaaatttg cgccaacgtc ggcccgtctac 360
 aaagtgtata tcatcgacga ggtgcatatg ctgtcgatcg gtgcgtttta cgcgctgttg 420
 aaaacggttg aggagccgcc gaaacacgtc attttcattt tggccacgac cgagccgcac 480
 aaaattccgg cgacgatcat ttcccgtgct caacggttcg attttcgcc catcccgtt 540
 caggcgatcg tttcacggct aaagtacgtc gcaagcgccc aagggtgtcg ggcgtcagat 600
 gaggcattgt ccgccatcgc ccgtgctgca gacgggggga tgcgcgatgc gctcagcttg 660
 ctgtatcaag ccatttcgtt cagcgacggg aaacttcggc tcgacgacgt gctggcgatg 720
 accggggctg catcatttgc cgccttatcg agcttcacg aagccatcca ccgcaaagat 780
 acagcggcgg ttcttcagca cttggaaacg atgatggcg aagggaaga tccgcatcgt 840
 ttggttgaag acttgatttt gtactatcgc gatttattgc tgtacaaaac cgctccctat 900
 gtggaggagg cgattcaaat tgctgtcgtt gacgaagcgt tcacttcact gtcggaaatg 960
 attccggttt ccaatttata cgaggccatc gagttgctga acaaaagcca gcaagagatg 1020
 aagtggacaa accaccgcg ccttctgttg gaagtggcg ttgtgaaact ttgccatcca 1080
 tcagccgccg ccccgctcgt gtcggcttcc gagttggaac cgttgataaa gcggattgaa 1140
 acgctggagg cggaattgcg gcgcctgaag gaacaaccgc ctgcccctcc gtcgaccgcc 1200
 gcgcgggtga aaaaactgtc caaaccgatg aaaacgggg gatataaagc cccggttggc 1260
 cgcatttacg agctgttgaa acaggcgacg catgaagatt tagctttggt gaaaggatgc 1320
 tgggcggatg tgctcgacac gttgaaacgg cagcataaag tgctgcacgc tgccttgctg 1380
 caagagagcg agccggttg agcgagcgcc tcagcgtttg tattaaaatt caaatacgaa 1440
 atccactgca aaatggcgac cgatcccaca agttcgggtc aagaaaacgt cgaagcgatt 1500
 ttgtttgagc tgacaaaccg ccgctttgaa atggttagcca ttccggaggg agaatgggga 1560
 aaaataagag aagagttcat ccgcaataag gacgccatgg tggaaaaaag cgaagaagat 1620
 ccgttaatcg ccgaagcgaa gcggctgttt ggcggaagagc tgatcgaaat taaagaa 1677

<210> 182
 <211> 559
 <212> PRT
 <213> *Bacillus stearothermophilus*

<400> 182

Val Ala Tyr Gln Ala Leu Tyr Arg Val Phe Arg Pro Gln Arg Phe Ala
1 5 10 15

Asp Met Val Gly Gln Glu His Val Thr Lys Thr Leu Gln Ser Ala Leu
20 25 30

Leu Gln His Lys Ile Ser His Ala Tyr Leu Phe Ser Gly Pro Arg Gly
35 40 45

Thr Gly Lys Thr Ser Ala Ala Lys Ile Phe Ala Lys Ala Val Asn Cys
50 55 60

Glu Gln Ala Pro Ala Ala Glu Pro Cys Asn Glu Cys Pro Ala Cys Leu
65 70 75 80

Gly Ile Thr Asn Gly Thr Val Pro Asp Val Leu Glu Ile Asp Ala Ala
85 90 95

Ser Asn Asn Arg Val Asp Glu Ile Arg Asp Ile Arg Glu Lys Val Lys
100 105 110

Phe Ala Pro Thr Ser Ala Arg Tyr Lys Val Tyr Ile Ile Asp Glu Val
115 120 125

His Met Leu Ser Ile Gly Ala Phe Asn Ala Leu Leu Lys Thr Leu Glu
130 135 140

Glu Pro Pro Lys His Val Ile Phe Ile Leu Ala Thr Thr Glu Pro His
145 150 155 160

Lys Ile Pro Ala Thr Ile Ile Ser Arg Cys Gln Arg Phe Asp Phe Arg
165 170 175

Arg Ile Pro Leu Gln Ala Ile Val Ser Arg Leu Lys Tyr Val Ala Ser
180 185 190

Ala Gln Gly Val Glu Ala Ser Asp Glu Ala Leu Ser Ala Ile Ala Arg
195 200 205

Ala Ala Asp Gly Gly Met Arg Asp Ala Leu Ser Leu Leu Asp Gln Ala
210 215 220

Ile Ser Phe Ser Asp Gly Lys Leu Arg Leu Asp Asp Val Leu Ala Met
225 230 235 240

Thr Gly Ala Ala Ser Phe Ala Ala Leu Ser Ser Phe Ile Glu Ala Ile
245 250 255

His Arg Lys Asp Thr Ala Ala Val Leu Gln His Leu Glu Thr Met Met
260 265 270

Ala Gln Gly Lys Asp Pro His Arg Leu Val Glu Asp Leu Ile Leu Tyr
275 280 285

Tyr Arg Asp Leu Leu Leu Tyr Lys Thr Ala Pro Tyr Val Glu Gly Ala
290 295 300

Ile Gln Ile Ala Val Val Asp Glu Ala Phe Thr Ser Leu Ser Glu Met
305 310 315 320

Ile Pro Val Ser Asn Leu Tyr Glu Ala Ile Glu Leu Leu Asn Lys Ser
325 330 335

Gln Gln Glu Met Lys Trp Thr Asn His Pro Arg Leu Leu Leu Glu Val
340 345 350

Ala Leu Val Lys Leu Cys His Pro Ser Ala Ala Ala Pro Ser Leu Ser
355 360 365

Ala Ser Glu Leu Glu Pro Leu Ile Lys Arg Ile Glu Thr Leu Glu Ala
370 375 380

Glu Leu Arg Arg Leu Lys Glu Gln Pro Pro Ala Pro Pro Ser Thr Ala
385 390 395 400

Ala Pro Val Lys Lys Leu Ser Lys Pro Met Lys Thr Gly Gly Tyr Lys
405 410 415

Ala Pro Val Gly Arg Ile Tyr Glu Leu Leu Lys Gln Ala Thr His Glu
420 425 430

Asp Leu Ala Leu Val Lys Gly Cys Trp Ala Asp Val Leu Asp Thr Leu
435 440 445

Lys Arg Gln His Lys Val Ser His Ala Ala Leu Leu Gln Glu Ser Glu
450 455 460

Pro Val Ala Ala Ser Ala Ser Ala Phe Val Leu Lys Phe Lys Tyr Glu
465 470 475 480

Ile His Cys Lys Met Ala Thr Asp Pro Thr Ser Ser Val Lys Glu Asn
485 490 495

Val Glu Ala Ile Leu Phe Glu Leu Thr Asn Arg Arg Phe Glu Met Val
500 505 510

Ala Ile Pro Glu Gly Glu Trp Gly Lys Ile Arg Glu Glu Phe Ile Arg
515 520 525

Asn Lys Asp Ala Met Val Glu Lys Ser Glu Glu Asp Pro Leu Ile Ala
530 535 540

Glu Ala Lys Arg Leu Phe Gly Glu Glu Leu Ile Glu Ile Lys Glu
545 550 555

<210> 183

<211> 4301

<212> DNA

<213> *Bacillus stearothermophilus*

<400> 183

```

atggtgacaa aagagcaaaa agagcgggtt ctcacctgc ttgagcagct gaagatgacg 60
tcggacgaat ggatgccgca ttttcgtgag gcagccattc gcaaagtcgt gatcgataaa 120
gaggagaaaa gctggcattt ttattttcag ttcgacaacg tgctgccggt tcatgtatac 180
aaaacgtttg ccgatcggct gcagacggcg ttccgccata tcgccgccgt ccgccatacg 240
atggagggtcg aagcgccgcg cgtaactgag gcggatgtgc aggcgtattg gccgctttgc 300
cttgccgagc tgcaagaagg catgtcgccg cttgtcgatt ggctcagccg gcagacgcct 360
gagctgaaaag gaaacaagct gcttgtcggt gcccgccatg aagcggaagc gctggcgatc 420
aaacggcgggt tcgcaaaaaa aatcgctgat gtgtacgctt cgtttgggtt tccccccctt 480
cagcttgacg tcagcgtcga gccgtccaag caagaaatgg aacagttttt ggcgcaaaaa 540
cagcaagagg acgaagagcg agcgcttgct gtactgaccg atttagcgag ggaagaagaa 600
aaggccgctg ctgcgcgccg gtccgggtccg cttgtcatcg gctatccgat ccgcgacgag 660
gagccggtgc ggccggttga aacgatcgtc gaagaagagc ggcgcgctcg tgtgcaaggc 720
tatgtatttg acgccgaagt gagcgaatta aaaagcggcc gcacgctggt gacctgaaa 780
atcacagatt acacgaactc gattttagtc aaaatgttct cgcgcgacaa agaggacgcc 840
gagcttatga gcggcgtaaa aaaaggcatg tgggtgaaag tgcgcggcag cgtgcaaaac 900
gatacgttcg tccgtgattt ggtcatcatc gccaacgatt tgaacgaaat cgccgcaaac 960
gaacggcaag atacggcgcc ggaaggggaa aagagggtcg agctccattt gcataccccg 1020
atgagccaaa tggacgcggt cacctcggtg acaaaactca ttgagcaagc gaaaaaatgg 1080
gggcatccgg cgatcgccgt caccgaccat gccgttggtc agtcgtttcc ggaggcctac 1140
agcgcgcgca aaaaacacgg catgaaggct atttacggcc ttgaggcgaa catcgtcgac 1200
gatggcggtc cgatcgctta caatgagacg caccgcgctc tttcgagga aacgtacgtc 1260
gtctttgacg tcgagacgac gggcctgtcg gctgtgtaca atacgatcat tgagctggcg 1320
gcgggtgaaag tgaaagacgg cgagatcatc gaccgattca tgtcgtttgc caacctgga 1380
catccgttgt cggtgacaac gatggagctg actgggatca ccgatgagat ggtgaaagac 1440
gccccgaagc cggacgaggt gctagcccggt tttgttgact gggccggcga tgcgacgctt 1500
gttgcccaca acgccagctt tgacatcggt tttttaaacg cgggcctcgc tcgcatgggg 1560
cgcggcaaaa tcgcgaatcc agtcatcgat acgctcgagc tggcccgttt tttatacccg 1620
gatttgaaaa accatcggtt caatacattg tgcaaaaaat ttgacattga attgacgcag 1680
catcaccgcg ccatctacga cgcgaggcgg accgggcatt tgcttatgcy gctgttgaag 1740
gaagcggaag agcgcggcgt actgtttcat gacgaattaa acagccgcac gcacagcgaa 1800
gcgtcctatc ggcttgcgcg cccgttccat gtgacgctgt tggcgcaaaa cgagactgga 1860

```

```

ttgaaaaatt tgttcaagct tgtgtcattg tgcacattc aatattttca ccgtgtgccg 1920
cgcatcccg cgtccgtgct cgtcaagcac cgcgacggcc tgcttgtcgg ctcgggctgc 1980
gacaaaggag agctgtttga caacttgatc caaaaggcgc cggaagaagt cgaagacatc 2040
gcccgttttt acgattttct tgaagtgcac ccgccggacg tgtacaagcc gtcacatcgag 2100
atggattatg tgaaagacga agagatgatc aaaaacatca tccgcagcat cgtcgccctt 2160
ggtgagaagc ttgacatccc ggttgtcgcc actggcaacg tccattactt gaaccagaa 2220
gataaaatth accggaaaat cttaatccat tgcgaaggcg gggcgaatcc gctcaaccgc 2280
catgaactgc cggatgtata tttccgtacg acgaatgaaa tgcttgactg cttctcgttt 2340
ttagggccgg aaaaagcgaa ggaaatcgct gttgacaaca cgcaaaaaat cgcttcgtta 2400
atcggcgatg tcaagccgat caaagatgag ctgtatacgc cgcgcattga agggcgggac 2460
gaggaaatca gggaaatgag ctaccggcgg gcgaaggaaa tttacggcga cccgttgccg 2520
aaacttgttg aagagcggct tgagaaggag ctaaaaagca tcatcggcca tggctttgcc 2580
gtcattttat tgatctcgca caagcttgtg aaaaaatcgc tcgatgacgg ctaccttgtc 2640
gggtcgcgcg gatcggtcgg ctcgctggtt gtcgacgacg tgacggaaat caccgaggtc 2700
aatccgctgc cgcgcatta cgtttgcccg aactgcaagc attcggagtt ctttaacgac 2760
ggttcagtcg gctcaggggt tgatttgccg gataaaaact gccgcgatg tgggacgaaa 2820
tacaagaaaag acgggcacga catcccgttt gagacgtttc tcggctttaa aggcgacaaa 2880
gtgccggata tcgacttgaa cttttccggc gaataccagc cgcgcgccc aactatacgc 2940
aaagtgtctg ttggcgaaga caacgtctac cgcgcgggga cgattggcac ggtcgctgac 3000
aaaacggcgt acggatttgt caaagcgtat gcgagcgacc ataacttaga gtcgcggcgc 3060
gcggaaatcg acggctcgcg gctggctgca ccggggtgaa gcggacgacc gggcagcatc 3120
cgggcggcat catcgctcgc ccggattata tggaaattta cgattttacg ccgattcaat 3180
atccggccga tgacacgtcc tctgaatggc ggacgacca tttcgacttc cattcgatcc 3240
acgacaatth gttgaagctc gatattctcg ggcacgacga tccgacggtc attcgcatgc 3300
tgcaagatth aagcggcatc gatccgaaaa cgatcccgac cgacgaccgc gatgtgatgg 3360
gcattttcag cagcaccgag ccgcttggcg ttacgcggga gcaaatcatg tgcaatgtcg 3420
gcacgatcgg cattccggag tttggcacgc gcttcgttcg gcaaatgttg gaagagacaa 3480
ggccaaaaac gttttccgaa ctogtgcaaa tttccggctt gtcgcacggc accgatgtgt 3540
ggctcggcaa cgcgcaagag ctcatcaaaa acggcacgtg tacgttatcg gaagtcacg 3600
gctgcccgca cgacattatg gtctatttga tttaccgcgg gctcgagccg tcgctcgctt 3660
ttaaatacat ggaatccgtg cgcaaaggaa aaggcttaac gccggagttt gaagcagaaa 3720
tgcgcaaaaa tgacgtgccg gagtggtaca tcgattcatg caaaaaaatc aagtacatgt 3780
tcccgaagc gcacgccgcc gcctacgtgt taatggcggg gcgcacgcc tactttaagg 3840
tgacccatcc gcttttgtat tacgcgtcgt actttacggg gcgggcggag gactttgacc 3900
ttgacgccat gatcaaagga tccccgccca ttcgcaagcg gattgaggaa atcaacgcc 3960
aaggcattca ggcgacggcg aaagaaaaaa gcttgctcac ggttcttgag gtggccttag 4020
agatgtgcga gcgcggcttt tcctttaaaa atatcgattt gtaccgctcg caggcgacgg 4080
aattcgatc tgacggcaat tctctcattc cgcggttcaa cgccattccg gggcttggga 4140
cgaacgtggc gcaggcgatc gtgcgcgcc cgcaggaagg cgagtttttg tcgaaggagg 4200
atthtgaaca gcgcggcaaa ttgtcgaaaa cgctgctcga gtatctagaa agccgcggct 4260
gccttgactc gcttcagac cataaccagc tgctcgctgt t 4301

```

<210> 184

<211> 1433

<212> PRT

<213> *Bacillus stearothermophilus*

<400> 184

Met Val Thr Lys Glu Gln Lys Glu Arg Phe Leu Ile Leu Leu Glu Gln
1 5 10 15

Leu Lys Met Thr Ser Asp Glu Trp Met Pro His Phe Arg Glu Ala Ala
20 25 30

Ile Arg Lys Val Val Ile Asp Lys Glu Glu Lys Ser Trp His Phe Tyr
35 40 45

Phe Gln Phe Asp Asn Val Leu Pro Val His Val Tyr Lys Thr Phe Ala
50 55 60

Asp Arg Leu Gln Thr Ala Phe Arg His Ile Ala Ala Val Arg His Thr
65 70 75 80

Met Glu Val Glu Ala Pro Arg Val Thr Glu Ala Asp Val Gln Ala Tyr
85 90 95

Trp Pro Leu Cys Leu Ala Glu Leu Gln Glu Gly Met Ser Pro Leu Val
100 105 110

Asp Trp Leu Ser Arg Gln Thr Pro Glu Leu Lys Gly Asn Lys Leu Leu
115 120 125

Val Val Ala Arg His Glu Ala Glu Ala Leu Ala Ile Lys Arg Arg Phe
130 135 140

Ala Lys Lys Ile Ala Asp Val Tyr Ala Ser Phe Gly Phe Pro Pro Leu
145 150 155 160

Gln Leu Asp Val Ser Val Glu Pro Ser Lys Gln Glu Met Glu Gln Phe
165 170 175

Leu Ala Gln Lys Gln Gln Glu Asp Glu Glu Arg Ala Leu Ala Val Leu
180 185 190

Thr Asp Leu Ala Arg Glu Glu Glu Lys Ala Ala Ser Ala Pro Pro Ser
195 200 205

Gly Pro Leu Val Ile Gly Tyr Pro Ile Arg Asp Glu Glu Pro Val Arg
210 215 220

Arg Leu Glu Thr Ile Val Glu Glu Glu Arg Arg Val Val Val Gln Gly
225 230 235 240

Tyr Val Phe Asp Ala Glu Val Ser Glu Leu Lys Ser Gly Arg Thr Leu
245 250 255

Leu Thr Met Lys Ile Thr Asp Tyr Thr Asn Ser Ile Leu Val Lys Met
 260 265 270
 Phe Ser Arg Asp Lys Glu Asp Ala Glu Leu Met Ser Gly Val Lys Lys
 275 280 285
 Gly Met Trp Val Lys Val Arg Gly Ser Val Gln Asn Asp Thr Phe Val
 290 295 300
 Arg Asp Leu Val Ile Ile Ala Asn Asp Leu Asn Glu Ile Ala Ala Asn
 305 310 315 320
 Glu Arg Gln Asp Thr Ala Pro Glu Gly Glu Lys Arg Val Glu Leu His
 325 330 335
 Leu His Thr Pro Met Ser Gln Met Asp Ala Val Thr Ser Val Thr Lys
 340 345 350
 Leu Ile Glu Gln Ala Lys Lys Trp Gly His Pro Ala Ile Ala Val Thr
 355 360 365
 Asp His Ala Val Val Gln Ser Phe Pro Glu Ala Tyr Ser Ala Ala Lys
 370 375 380
 Lys His Gly Met Lys Val Ile Tyr Gly Leu Glu Ala Asn Ile Val Asp
 385 390 395 400
 Asp Gly Val Pro Ile Ala Tyr Asn Glu Thr His Arg Arg Leu Ser Glu
 405 410 415
 Glu Thr Tyr Val Val Phe Asp Val Glu Thr Thr Gly Leu Ser Ala Val
 420 425 430
 Tyr Asn Thr Ile Ile Glu Leu Ala Ala Val Lys Val Lys Asp Gly Glu
 435 440 445
 Ile Ile Asp Arg Phe Met Ser Phe Ala Asn Pro Gly His Pro Leu Ser
 450 455 460
 Val Thr Thr Met Glu Leu Thr Gly Ile Thr Asp Glu Met Val Lys Asp
 465 470 475 480
 Ala Pro Lys Pro Asp Glu Val Leu Ala Arg Phe Val Asp Trp Ala Gly
 485 490 495
 Asp Ala Thr Leu Val Ala His Asn Ala Ser Phe Asp Ile Gly Phe Leu
 500 505 510

Asn	Ala	Gly	Leu	Ala	Arg	Met	Gly	Arg	Gly	Lys	Ile	Ala	Asn	Pro	Val	515	520	525
Ile	Asp	Thr	Leu	Glu	Leu	Ala	Arg	Phe	Leu	Tyr	Pro	Asp	Leu	Lys	Asn	530	535	540
His	Arg	Leu	Asn	Thr	Leu	Cys	Lys	Lys	Phe	Asp	Ile	Glu	Leu	Thr	Gln	545	550	555
His	His	Arg	Ala	Ile	Tyr	Asp	Ala	Glu	Ala	Thr	Gly	His	Leu	Leu	Met	565	570	575
Arg	Leu	Leu	Lys	Glu	Ala	Glu	Glu	Arg	Gly	Ile	Leu	Phe	His	Asp	Glu	580	585	590
Leu	Asn	Ser	Arg	Thr	His	Ser	Glu	Ala	Ser	Tyr	Arg	Leu	Ala	Arg	Pro	595	600	605
Phe	His	Val	Thr	Leu	Leu	Ala	Gln	Asn	Glu	Thr	Gly	Leu	Lys	Asn	Leu	610	615	620
Phe	Lys	Leu	Val	Ser	Leu	Ser	His	Ile	Gln	Tyr	Phe	His	Arg	Val	Pro	625	630	635
Arg	Ile	Pro	Arg	Ser	Val	Leu	Val	Lys	His	Arg	Asp	Gly	Leu	Leu	Val	645	650	655
Gly	Ser	Gly	Cys	Asp	Lys	Gly	Glu	Leu	Phe	Asp	Asn	Leu	Ile	Gln	Lys	660	665	670
Ala	Pro	Glu	Glu	Val	Glu	Asp	Ile	Ala	Arg	Phe	Tyr	Asp	Phe	Leu	Glu	675	680	685
Val	His	Pro	Pro	Asp	Val	Tyr	Lys	Pro	Leu	Ile	Glu	Met	Asp	Tyr	Val	690	695	700
Lys	Asp	Glu	Glu	Met	Ile	Lys	Asn	Ile	Ile	Arg	Ser	Ile	Val	Ala	Leu	705	710	715
Gly	Glu	Lys	Leu	Asp	Ile	Pro	Val	Val	Ala	Thr	Gly	Asn	Val	His	Tyr	725	730	735
Leu	Asn	Pro	Glu	Asp	Lys	Ile	Tyr	Arg	Lys	Ile	Leu	Ile	His	Ser	Gln	740	745	750
Gly	Gly	Ala	Asn	Pro	Leu	Asn	Arg	His	Glu	Leu	Pro	Asp	Val	Tyr	Phe	755	760	765

Arg	Thr	Thr	Asn	Glu	Met	Leu	Asp	Cys	Phe	Ser	Phe	Leu	Gly	Pro	Glu	
770						775					780					
Lys	Ala	Lys	Glu	Ile	Val	Val	Asp	Asn	Thr	Gln	Lys	Ile	Ala	Ser	Leu	
785					790					795					800	
Ile	Gly	Asp	Val	Lys	Pro	Ile	Lys	Asp	Glu	Leu	Tyr	Thr	Pro	Arg	Ile	
				805					810					815		
Glu	Gly	Ala	Asp	Glu	Glu	Ile	Arg	Glu	Met	Ser	Tyr	Arg	Arg	Ala	Lys	
			820					825					830			
Glu	Ile	Tyr	Gly	Asp	Pro	Leu	Pro	Lys	Leu	Val	Glu	Glu	Arg	Leu	Glu	
		835					840					845				
Lys	Glu	Leu	Lys	Ser	Ile	Ile	Gly	His	Gly	Phe	Ala	Val	Ile	Tyr	Leu	
		850				855					860					
Ile	Ser	His	Lys	Leu	Val	Lys	Lys	Ser	Leu	Asp	Asp	Gly	Tyr	Leu	Val	
865					870					875					880	
Gly	Ser	Arg	Gly	Ser	Val	Gly	Ser	Ser	Phe	Val	Ala	Thr	Met	Thr	Glu	
				885					890					895		
Ile	Thr	Glu	Val	Asn	Pro	Leu	Pro	Pro	His	Tyr	Val	Cys	Pro	Asn	Cys	
			900					905					910			
Lys	His	Ser	Glu	Phe	Phe	Asn	Asp	Gly	Ser	Val	Gly	Ser	Gly	Phe	Asp	
		915					920					925				
Leu	Pro	Asp	Lys	Asn	Cys	Pro	Arg	Cys	Gly	Thr	Lys	Tyr	Lys	Lys	Asp	
		930				935					940					
Gly	His	Asp	Ile	Pro	Phe	Glu	Thr	Phe	Leu	Gly	Phe	Lys	Gly	Asp	Lys	
945					950					955					960	
Val	Pro	Asp	Ile	Asp	Leu	Asn	Phe	Ser	Gly	Glu	Tyr	Gln	Pro	Arg	Ala	
				965					970					975		
His	Asn	Tyr	Thr	Lys	Val	Leu	Phe	Gly	Glu	Asp	Asn	Val	Tyr	Arg	Ala	
			980					985					990			
Gly	Thr	Ile	Gly	Thr	Val	Ala	Asp	Lys	Thr	Ala	Tyr	Gly	Phe	Val	Lys	
		995				1000						1005				
Ala	Tyr	Ala	Ser	Asp	His	Asn	Leu	Glu	Leu	Arg	Gly	Ala	Glu	Ile	Asp	
1010						1015					1020					

<223> Description of Artificial Sequence: PCR primer

<400> 187

ggcccccttg gccttctcgg cctccat

27

<210> 188

<211> 331

<212> DNA

<213> *Thermus thermophilus*

<400> 188

agactcgagg ccctggagcg ggagctggag aagcttgccc tcctctcccc acccctcacc 60
ctggagaagg tggagaaggt ggtggccctg agggccccc tcacgggctt tgacctggtg 120
cgctccgtcc tggagaagga cccaaggag gccctcctgc gcctcaggcg cctcaggagg 180
gagggggagg agccctcag gctcctcggg gccctctcct ggcagttcgc cctcctcgcc 240
cgggccttct tcctcctcgg ggaaaacccc agggccaagg aggaggacct cggccgcctc 300
gaggcccacc cctacgccgc caagaaggcc a 331

<210> 189

<211> 110

<212> PRT

<213> *Thermus thermophilus*

<400> 189

Arg Leu Glu Ala Leu Glu Arg Glu Leu Glu Lys Leu Ala Leu Leu Ser
1 5 10 15

Pro Pro Leu Thr Leu Glu Lys Val Glu Lys Val Val Ala Leu Arg Pro
20 25 30

Pro Leu Thr Gly Phe Asp Leu Val Arg Ser Val Leu Glu Lys Asp Pro
35 40 45

Lys Glu Ala Leu Leu Arg Leu Arg Arg Leu Arg Glu Glu Gly Glu Glu
50 55 60

Pro Leu Arg Leu Leu Gly Ala Leu Ser Trp Gln Phe Ala Leu Leu Ala
65 70 75 80

Arg Ala Phe Phe Leu Leu Arg Glu Asn Pro Arg Pro Lys Glu Glu Asp
85 90 95

Leu Ala Arg Leu Glu Ala His Pro Tyr Ala Ala Lys Lys Ala
100 105 110

<210> 190
<211> 31
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR primer

<400> 190
gtgggtgtcta gacatcataa cggttctggc a 31

<210> 191
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR Primer

<400> 191
gagggccacc accttctcca ctttctc 27

<210> 192
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR Primer

<400> 192
ctccgtcctg gagaaggacc ccaag 25

<210> 193
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR primer

<220>
<221> primer_bind
<222> (15)
<223> S at position 15 can be either C or G

<220>
 <221> primer_bind
 <222> (27)
 <223> S at position 27 can be either C or G

<400> 193
 cgccaattca acgcscctct caagacsct

29

<210> 194
 <211> 31
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: PCR primer

<400> 194
 gacacttaac atatgggtcat cgccttcacc g

31

<210> 195
 <211> 38
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: PCR primer

<400> 195
 gtgtgtgaat tcgggtcaac gggcgaggcg gaggaccg

38

<210> 196
 <211> 10
 <212> PRT
 <213> Deinococcus radiodurans

<400> 196
 Val Ile Leu Asn Pro Gly Ser Val Gly Gln
 1 5 10

<210> 197
 <211> 10
 <212> PRT
 <213> Methanococcus jannaschii

<400> 197

Tyr Leu Ile Asn Pro Gly Ser Val Gly Gln
1 5 10

<210> 198

<211> 10

<212> PRT

<213> *Thermotoga maritima*

<400> 198

Leu Val Leu Asn Pro Gly Ser Ala Gly Arg
1 5 10

<210> 199

<211> 28

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR primer

<400> 199

ctggtgaacc cgggctcgt gggccagc

28

<210> 200

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: polypeptide

<400> 200

Leu Leu Val Asn Pro Gly Ser Val Gly Gln
1 5 10

<210> 201

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR primer

<400> 201
ctcgaggagc ttgaggaggg tgttggc

27

<210> 202
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: polypeptide

<400> 202
Ala Asn Thr Leu Leu Lys Leu Leu Glu
1 5

<210> 203
<211> 32
<212> PRT
<213> Deinococcus radiodurans

<400> 203
Gly Phe Gly Gly Val Gln Leu His Ala Ala His Gly Tyr Leu Leu Ser
1 5 10 15
Gln Phe Leu Ser Pro Arg His Asn Val Arg Glu Asp Glu Tyr Gly Gly
20 25 30

<210> 204
<211> 32
<212> PRT
<213> Caenorhabditis elegans

<400> 204
Gly Phe Asp Gly Ile Gln Leu His Gly Ala His Gly Tyr Leu Leu Ser
1 5 10 15
Gln Phe Thr Ser Pro Thr Thr Asn Lys Arg Val Asp Lys Tyr Gly Gly
20 25 30

<210> 205
 <211> 32
 <212> PRT
 <213> *Pseudomonas aeruginosa*

<400> 205
 Gly Phe Ser Gly Val Glu Ile His Ala Ala His Gly Tyr Leu Leu Ser
 1 5 10 15
 Gln Phe Leu Ser Pro Leu Ser Asn Arg Arg Ser Asp Ala Trp Gly Gly
 20 25 30

<210> 206
 <211> 32
 <212> PRT
 <213> *Archaeoglobus fulgidus*

<400> 206
 Gly Phe Asp Ala Val Gln Leu His Ala Ala His Gly Tyr Leu Leu Ser
 1 5 10 15
 Glu Phe Ile Ser Pro His Val Asn Arg Arg Lys Asp Glu Tyr Gly Gly
 20 25 30

<210> 207
 <211> 30
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: PCR primer

<400> 207
 catcctggac tcggcccacc tcctcaccga 30

<210> 208
 <211> 9

<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: polypeptide

<400> 208
Ile Leu Asp Ser Ala His Leu Leu Thr
1 5

<210> 209
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR primer

<400> 209
gaggaggtag ccgtgggccg cgtggagctc cac 33

<210> 210
<211> 11
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: polypeptide

<400> 210
Val Glu Leu His Ala Ala His Gly Tyr Leu Leu
1 5 10

<210> 211
<211> 32
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR primer

<400> 211
ggctttccca tatggctcta caccggctc ac 32

<210> 212

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR primer

<400> 212

gcgtggatcc acggtcattgt ctctaagtc

29